

**Clinical trial results:****A Phase 3, Double-Blind, Randomized, Vehicle-Controlled, Efficacy and Safety Study of Ruxolitinib Cream Followed by an Extension Period in Participants With Vitiligo****Summary**

EudraCT number	2019-000846-37
Trial protocol	FR PL DE BG ES IT
Global end of trial date	21 October 2021

**Results information**

Result version number	v2 (current)
This version publication date	11 October 2022
First version publication date	05 May 2022
Version creation reason	

**Trial information****Trial identification**

Sponsor protocol code	INCB 18424-306
-----------------------	----------------

**Additional study identifiers**

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT04052425
WHO universal trial number (UTN)	-

Notes:

**Sponsors**

Sponsor organisation name	Incyte Corporation
Sponsor organisation address	1801 Augustine Cutoff Drive, Wilmington, United States, 19803
Public contact	Study Director, Incyte Corporation, 1 18554633463, medinfo@incyte.com
Scientific contact	Study Director, Incyte Corporation, 1 18554633463, medinfo@incyte.com

Notes:

**Paediatric regulatory details**

Is trial part of an agreed paediatric investigation plan (PIP)	Yes
EMA paediatric investigation plan number(s)	EMA-002618-PIP02-20
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	21 October 2021
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	21 October 2021
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

The purpose of this study was to evaluate the efficacy and safety of ruxolitinib cream in adolescent and adult participants with non-segmental vitiligo with facial involvement for whom total body involved vitiligo area (facial and nonfacial) did not exceed 10% body surface area (BSA).

Protection of trial subjects:

This study was performed in accordance with ethical principles that have their origin in the Declaration of Helsinki and conducted in adherence to the study Protocol, applicable Good Clinical Practices, and applicable laws and country-specific regulations in which the study was being conducted.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	20 September 2019
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	United States: 204
Country: Number of subjects enrolled	Bulgaria: 17
Country: Number of subjects enrolled	Canada: 16
Country: Number of subjects enrolled	France: 19
Country: Number of subjects enrolled	Germany: 6
Country: Number of subjects enrolled	Italy: 1
Country: Number of subjects enrolled	Poland: 63
Country: Number of subjects enrolled	Spain: 4
Worldwide total number of subjects	330
EEA total number of subjects	110

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23	0

months)	
Children (2-11 years)	0
Adolescents (12-17 years)	36
Adults (18-64 years)	265
From 65 to 84 years	29
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

This study was conducted at 45 study centers in North America and Europe.

### Pre-assignment

Screening details:

A total of 330 participants were randomized into the study. All randomized participants (Intent-to-Treat Population) applied study drug at least once (Safety Population), and 283 participants applied ruxolitinib cream at least once during the Treatment-Extension (TE) Period (TE Evaluable Population).

### Period 1

Period 1 title	24-Week Double-blind Period
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator, Monitor, Data analyst, Assessor

### Arms

Are arms mutually exclusive?	Yes
------------------------------	-----

<b>Arm title</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID
------------------	---

Arm description:

Participants applied ruxolitinib 1.5% cream twice daily (BID) for 24 weeks.

Arm type	Experimental
Investigational medicinal product name	ruxolitinib
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Cream
Routes of administration	Topical use

Dosage and administration details:

1.5% cream twice daily

<b>Arm title</b>	Double-Blind Period: Vehicle cream BID
------------------	--

Arm description:

Participants applied matching vehicle cream BID for 24 weeks.

Arm type	Placebo
Investigational medicinal product name	Vehicle
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Cream
Routes of administration	Topical use

Dosage and administration details:

twice daily

Number of subjects in period 1	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID
	Started	221
Completed	193	90
Not completed	28	19
Physician decision	1	-
Consent withdrawn by subject	9	10
Adverse event, non-fatal	-	1
Discontinued Treatment Due to COVID-19 Pandemic	3	-
Lost to follow-up	14	7
Protocol deviation	1	-
Lack of efficacy	-	1

## Period 2

Period 2 title	28-Week Treatment-Extension Period
Is this the baseline period?	No
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator, Monitor

## Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID

### Arm description:

Participants who completed the Week 24 assessments with no safety concerns could continue into the 28-week Treatment-Extension Period. Participants who applied ruxolitinib cream 1.5% BID during the Double-Blind Period continued to apply ruxolitinib cream 1.5% BID for an additional 28 weeks in the Treatment-Extension Period.

Arm type	Experimental
Investigational medicinal product name	ruxolitinib
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Cream
Routes of administration	Topical use

### Dosage and administration details:

1.5% cream twice daily

<b>Arm title</b>	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID
------------------	--

### Arm description:

Participants who completed the Week 24 assessments with no safety concerns could continue into the 28-week Treatment-Extension Period. Participants who applied vehicle cream BID during the Double-Blind Period applied ruxolitinib cream 1.5% BID for 28 weeks in the Treatment-Extension Period.

Arm type	Experimental
----------	--------------

Investigational medicinal product name	Vehicle
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Cream
Routes of administration	Topical use

Dosage and administration details:

twice daily

Investigational medicinal product name	ruxolitinib
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Cream
Routes of administration	Topical use

Dosage and administration details:

1.5% cream twice daily

Number of subjects in period 2	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID
Started	193	90
Completed	174	80
Not completed	19	10
Consent withdrawn by subject	10	7
Physician decision	-	1
Adverse event, non-fatal	1	-
Participant Moved	1	1
Lost to follow-up	5	1
Sponsor Opinion Due to Safety Reason	1	-
Lack of efficacy	1	-

## Baseline characteristics

### Reporting groups

Reporting group title	Double-Blind Period: Ruxolitinib cream 1.5% BID
Reporting group description: Participants applied ruxolitinib 1.5% cream twice daily (BID) for 24 weeks.	
Reporting group title	Double-Blind Period: Vehicle cream BID
Reporting group description: Participants applied matching vehicle cream BID for 24 weeks.	

Reporting group values	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID	Total
Number of subjects	221	109	330
Age categorical Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	0	0	0
Infants and toddlers (28 days-23 months)	0	0	0
Children (2-11 years)	0	0	0
Adolescents (12-17 years)	25	11	36
Adults (18-64 years)	180	85	265
From 65-84 years	16	13	29
85 years and over	0	0	0
Age Continuous Units: years			
arithmetic mean	40.5	39.7	
standard deviation	± 15.44	± 16.71	-
Sex: Female, Male Units: participants			
Female	136	50	186
Male	85	59	144
Race, Customized Units: Subjects			
White	180	96	276
Black/African American	11	4	15
Asian	5	4	9
American Indian/Alaska Native	1	0	1
Not Reported	16	3	19
Latino	3	1	4
Hispanic or Latino	1	0	1
Iranian	1	0	1
Indian	1	1	2
Hispanic	1	0	1
Black-Hispanic	1	0	1
Ethnicity, Customized Units: Subjects			

Hispanic or Latino	53	20	73
Not Hispanic or Latino	151	86	237
Not Reported	15	3	18
Unknown	1	0	1
Captured as "Other"	1	0	1
<b>Face Vitiligo Area Scoring Index (F-VASI)</b>			
F-VASI was measured by the percentage of vitiligo involvement (percentage of body surface area [BSA]; assessed by the Investigator) and the degree of depigmentation: 0% (no depigmentation), 10% (specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), 100% (no pigment). F-VASI was derived by multiplying the vitiligo involvement values by the percentage of affected skin for each facial site and summing all values (range: 0-3; higher values=worse outcome).			
Units: scores on a scale			
arithmetic mean	0.932	0.999	
standard deviation	± 0.5813	± 0.5942	-
<b>Facial Body Surface Area (F-BSA) Involvement</b>			
F-BSA involvement was the proportion of the facial body surface area with vitiligo. The area "Face" was defined as including the area on the forehead to the original hairline, on the cheek to the jawline vertically to the jawline and laterally from the corner of the mouth to the tragus. The area "Face" did not include surface area of the lips, scalp, ears, or neck, but included the nose and eyelids.			
Units: percentage of facial surface area			
arithmetic mean	1.05	1.15	
standard deviation	± 0.692	± 0.710	-
<b>Total Body Vitiligo Area Scoring Index (T-VASI)</b>			
T-VASI was measured by the percentage of vitiligo involvement from all body regions (percentage of BSA; Investigator assessed) and the degree of depigmentation: 0% (no depigmentation), 10% (specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), 100% (no pigment). T-VASI was derived by multiplying the vitiligo involvement values by the percentage of affected skin for each site and summing all values (possible range: 0-100; higher values=worse outcome).			
Units: scores on a scale			
arithmetic mean	6.489	6.424	
standard deviation	± 2.0228	± 1.9241	-
<b>Total Body Surface Area (T-BSA) Involvement</b>			
T-BSA involvement was the proportion of the body surface area with vitiligo. The body was divided into the following 6 separate and mutually exclusive sites: (1) head/neck, (2) hands, (3) upper extremities (excluding hands), (4) trunk, (5) lower extremities (excluding feet), and (6) feet.			
Units: percentage of total body surface area			
arithmetic mean	7.28	7.22	
standard deviation	± 2.033	± 2.008	-

## End points

### End points reporting groups

Reporting group title	Double-Blind Period: Ruxolitinib cream 1.5% BID
Reporting group description:	Participants applied ruxolitinib 1.5% cream twice daily (BID) for 24 weeks.
Reporting group title	Double-Blind Period: Vehicle cream BID
Reporting group description:	Participants applied matching vehicle cream BID for 24 weeks.
Reporting group title	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID
Reporting group description:	Participants who completed the Week 24 assessments with no safety concerns could continue into the 28-week Treatment-Extension Period. Participants who applied ruxolitinib cream 1.5% BID during the Double-Blind Period continued to apply ruxolitinib cream 1.5% BID for an additional 28 weeks in the Treatment-Extension Period.
Reporting group title	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID
Reporting group description:	Participants who completed the Week 24 assessments with no safety concerns could continue into the 28-week Treatment-Extension Period. Participants who applied vehicle cream BID during the Double-Blind Period applied ruxolitinib cream 1.5% BID for 28 weeks in the Treatment-Extension Period.
Subject analysis set title	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID
Subject analysis set type	Intention-to-treat
Subject analysis set description:	Participants who completed the Week 24 assessments with no safety concerns could continue into the 28-week Treatment-Extension Period. Participants who applied ruxolitinib cream 1.5% BID during the Double-Blind Period continued to apply ruxolitinib cream 1.5% BID for an additional 28 weeks in the Treatment-Extension Period.
Subject analysis set title	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID
Subject analysis set type	Intention-to-treat
Subject analysis set description:	Participants who completed the Week 24 assessments with no safety concerns could continue into the 28-week Treatment-Extension Period. Participants who applied vehicle cream BID during the Double-Blind Period applied ruxolitinib cream 1.5% BID for 28 weeks in the Treatment-Extension Period.

### Primary: Percentage of participants achieving a $\geq 75\%$ Improvement from Baseline in the Face Vitiligo Area Scoring Index (F-VASI75) Score at Week 24

End point title	Percentage of participants achieving a $\geq 75\%$ Improvement from Baseline in the Face Vitiligo Area Scoring Index (F-VASI75) Score at Week 24
End point description:	An F-VASI75 responder achieved at least 75% improvement from Baseline in F-VASI, measured by the percentage of vitiligo involvement (percentage of body surface area [BSA]) and the degree of depigmentation: 0% (no depigmentation), 10% (only specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), or 100% (no pigment). The percentage of BSA (hand unit) vitiligo involvement was estimated to the nearest 0.1% by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate the percentage of BSA vitiligo involvement. F-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site on the face and summing the values of all sites (possible range: 0-3; lower scores indicate increased improvement).
End point type	Primary
End point timeframe:	Baseline; Week 24

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage of participants				
number (not applicable)	29.8	7.4		

## Statistical analyses

<b>Statistical analysis title</b>	exact logistic regression
Comparison groups	Double-Blind Period: Vehicle cream BID v Double-Blind Period: Ruxolitinib cream 1.5% BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.0001 <sup>[1]</sup>
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	5.28
Confidence interval	
level	95 %
sides	2-sided
lower limit	2.341
upper limit	11.903

Notes:

[1] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

## Secondary: Percentage of participants achieving a $\geq$ 50% Improvement from Baseline in the Face Vitiligo Area Scoring Index (F-VASI50) Score at Week 24

End point title	Percentage of participants achieving a $\geq$ 50% Improvement from Baseline in the Face Vitiligo Area Scoring Index (F-VASI50) Score at Week 24
-----------------	---

End point description:

An F-VASI50 responder achieved at least 50% improvement from Baseline in F-VASI, measured by the percentage of vitiligo involvement (percentage of BSA) and the degree of depigmentation: 0% (no depigmentation), 10% (only specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), or 100% (no pigment). The percentage of BSA (hand unit) vitiligo involvement was estimated to the nearest 0.1% by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate the percentage of BSA vitiligo involvement. F-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site on the face and summing the values of all sites (possible range: 0-3; lower scores indicate increased improvement).

End point type	Secondary
End point timeframe:	
Baseline; Week 24	

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage of participants				
number (not applicable)	51.2	16.9		

## Statistical analyses

<b>Statistical analysis title</b>	exact logistic regression
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.0001 [2]
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	5.18
Confidence interval	
level	95 %
sides	2-sided
lower limit	2.831
upper limit	9.482

Notes:

[2] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

## Secondary: Percentage of participants achieving a $\geq$ 90% Improvement from Baseline in the Face Vitiligo Area Scoring Index (F-VASI90) Score at Week 24

End point title	Percentage of participants achieving a $\geq$ 90% Improvement from Baseline in the Face Vitiligo Area Scoring Index (F-VASI90) Score at Week 24
-----------------	---

End point description:

An F-VASI90 responder achieved at least 90% improvement from Baseline in F-VASI, measured by the percentage of vitiligo involvement (percentage of BSA) and the degree of depigmentation: 0% (no depigmentation), 10% (only specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), or 100% (no pigment). The percentage of BSA (hand unit) vitiligo involvement was estimated to the nearest 0.1% by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate the percentage of BSA vitiligo involvement. F-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site on the face and summing the values of all sites (possible range: 0-3; lower scores indicate increased improvement).

End point type	Secondary
End point timeframe:	
Baseline; Week 24	

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage of participants				
number (not applicable)	15.3	2.2		

## Statistical analyses

<b>Statistical analysis title</b>	exact logistic regression
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.0038 [3]
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	8.49
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.997
upper limit	36.048

Notes:

[3] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

## Secondary: Percentage of participants achieving a $\geq$ 50% Improvement from Baseline in the Total Body Vitiligo Area Scoring Index (T-VASI50) Score at Week 24

End point title	Percentage of participants achieving a $\geq$ 50% Improvement from Baseline in the Total Body Vitiligo Area Scoring Index (T-VASI50) Score at Week 24
-----------------	---

End point description:

A T-VASI50 responder achieved at least 50% improvement from Baseline in T-VASI, calculated with contributions from 6 sites. The percentage of vitiligo involvement was estimated in hand units (percentage of BSA estimated to the nearest 0.1%) by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate percent BSA vitiligo involvement. The degree of depigmentation for each site was estimated to the nearest percentage: 0% (no depigmentation present), 10% (only specks of depigmentation present), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment present), 100% (no pigment present). T-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site and summing the values (range: 0-100; lower scores indicate increased improvement).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 24

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage of participants				
number (not applicable)	20.6	5.1		

### Statistical analyses

<b>Statistical analysis title</b>	exact logistic regression
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.002 <sup>[4]</sup>
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	4.93
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.795
upper limit	13.566

Notes:

[4] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

### Secondary: Percentage of participants achieving a Vitiligo Noticeability Scale (VNS) of 4 or 5 at Week 24

End point title	Percentage of participants achieving a Vitiligo Noticeability Scale (VNS) of 4 or 5 at Week 24
-----------------	--

End point description:

The VNS is a patient-reported measure of vitiligo treatment success that is rated on a 5-point scale. The Baseline facial photograph was shown to the participants for reference, and a mirror was provided for the participants to assess the vitiligo on their face. The participant was asked to respond to the following query: Compared with before treatment, how noticeable is the vitiligo now? Responses: (1) more noticeable, (2) as noticeable, (3) slightly less noticeable, (4) a lot less noticeable, and (5) no longer noticeable.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 24

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage of participants				
number (not applicable)	24.5	3.3		

## Statistical analyses

<b>Statistical analysis title</b>	exact logistic regression
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.0002 <sup>[5]</sup>
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	9.53
Confidence interval	
level	95 %
sides	2-sided
lower limit	2.9
upper limit	31.29

Notes:

[5] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

## Secondary: Percentage change from Baseline in Facial Body Surface Area (F-BSA) at Week 24

End point title	Percentage change from Baseline in Facial Body Surface Area (F-BSA) at Week 24
-----------------	--

End point description:

F-BSA involvement was the proportion of the facial body surface area with vitiligo. The area "Face" was defined as including the area on the forehead to the original hairline, on the cheek to the jawline vertically to the jawline and laterally from the corner of the mouth to the tragus. The area "Face" did not include surface area of the lips, scalp, ears, or neck, but included the nose and eyelids. Body surface area assessment was performed by the Palmar Method. Body surface area was estimated to the nearest 0.1%. The approximate size of the participant's entire palmar surface (i.e., the palm plus 5 digits) was considered as 1% BSA, and the approximate size of the participant's thumb was considered as 0.1% BSA. Percentage change = ([post-Baseline (BL) value minus BL value]/BL value) X 100.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 24

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage change				
least squares mean (standard error)	-28.9 (± 2.22)	-9.5 (± 3.25)		

## Statistical analyses

<b>Statistical analysis title</b>	ANCOVA
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.0001 <sup>[6]</sup>
Method	ANCOVA
Parameter estimate	least squares mean difference
Point estimate	-19.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	-27.05
upper limit	-11.64
Variability estimate	Standard error of the mean
Dispersion value	3.93

Notes:

[6] - Response Variable = Treatment + Stratification Factors (Skin Type Fitzpatrick scale Type I, II versus Type III, IV, V, and VI, Region North America/Europe) + Baseline

## Secondary: Number of participants with treatment-emergent adverse events (TEAEs) during the Double-Blind Period

End point title	Number of participants with treatment-emergent adverse events (TEAEs) during the Double-Blind Period
-----------------	--

End point description:

An adverse event (AE) was defined as any untoward medical occurrence associated with the use of a drug in humans, whether or not considered drug-related. An AE could have been any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of study treatment. A TEAE was defined as any AE reported for the first time or the worsening of a pre-existing event after the first application of study drug.

End point type	Secondary
----------------	-----------

End point timeframe:

from the time of Informed Consent Form signing until at least 30 days after the last application of study drug (up to Week 24)

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: participants	101	42		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Number of participants with treatment-emergent adverse events (TEAEs) during the Treatment-Extension Period

End point title	Number of participants with treatment-emergent adverse events (TEAEs) during the Treatment-Extension Period
-----------------	---

End point description:

An AE was defined as any untoward medical occurrence associated with the use of a drug in humans, whether or not considered drug-related. An AE could have been any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of study treatment. A TEAE was defined as any AE reported for the first time or the worsening of a pre-existing event after the first application of study drug.

End point type	Secondary
----------------	-----------

End point timeframe:

from the completion of the Week 24 assessments until at least 30 days after the last application of study drug (up to Week 52 + 30 days)

<b>End point values</b>	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	193	90		
Units: participants	65	31		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of participants achieving a $\geq 25\%$ Improvement in the Face Vitiligo Area Scoring Index (F-VASI25) Score at Week 24

End point title	Percentage of participants achieving a $\geq 25\%$ Improvement in the Face Vitiligo Area Scoring Index (F-VASI25) Score at Week 24
-----------------	--

End point description:

An F-VASI25 responder achieved at least 25% improvement from Baseline in F-VASI, measured by the

percentage of vitiligo involvement (percentage of BSA) and the degree of depigmentation: 0% (no depigmentation), 10% (only specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), or 100% (no pigment). The percentage of BSA (hand unit) vitiligo involvement was estimated to the nearest 0.1% by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate the percentage of BSA vitiligo involvement. F-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site on the face and summing the values of all sites (possible range: 0-3; lower scores indicate increased improvement).

End point type	Secondary
End point timeframe:	
Baseline; Week 24	

End point values	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage of participants				
number (not applicable)	69.8	30.0		

## Statistical analyses

<b>Statistical analysis title</b>	exact logistic regression
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	superiority
P-value	< 0.0001 [7]
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	5.56
Confidence interval	
level	95 %
sides	2-sided
lower limit	3.226
upper limit	9.578

Notes:

[7] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

## Secondary: Percentage of participants achieving a $\geq$ %25, $\geq$ %50, $\geq$ 75%, and $\geq$ 90% Improvement in the Face Vitiligo Area Scoring Index (F-VASI25/50/75/90) Score at Week 52

End point title	Percentage of participants achieving a $\geq$ %25, $\geq$ %50, $\geq$ 75%, and $\geq$ 90% Improvement in the Face Vitiligo Area Scoring Index (F-VASI25/50/75/90) Score at Week 52
-----------------	--

---

**End point description:**

An F-VASI25/50/75/90 responder achieved at least 25/50/75/90% improvement from Baseline in F-VASI, measured by the percentage of vitiligo involvement (percentage of BSA) and the degree of depigmentation: 0% (no depigmentation), 10% (only specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), or 100% (no pigment). The percentage of BSA (hand unit) vitiligo involvement was estimated to the nearest 0.1% by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate the percentage of BSA vitiligo involvement. F-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site on the face and summing the values of all sites (possible range: 0-3; lower scores indicate increased improvement).

---

End point type	Secondary
----------------	-----------

---

**End point timeframe:**Baseline; Week 52

---

<b>End point values</b>	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	173	82		
Units: percentage of participants				
number (not applicable)				
F-VASI25	89.6	74.4		
F-VASI50	75.1	56.1		
F-VASI75	52.6	26.8		
F-VASI90	32.9	12.2		

---

**Statistical analyses**No statistical analyses for this end point

---

---

**Secondary: Percentage change from Baseline in F-VASI at Week 24**

---

End point title	Percentage change from Baseline in F-VASI at Week 24
-----------------	--

---

**End point description:**

F-VASI was measured by the percentage of vitiligo involvement (percentage of BSA) and the degree of depigmentation: 0% (no depigmentation), 10% (only specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), or 100% (no pigment). The percentage of BSA (hand unit) vitiligo involvement was estimated to the nearest 0.1% by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate the percentage of BSA vitiligo involvement. F-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site on the face and summing the values of all sites (possible range: 0-3; lower scores indicate increased improvement). Percentage change =  $([\text{post-BL value} - \text{BL value}] / \text{BL value}) \times 100$ .

---

End point type	Secondary
----------------	-----------

---

**End point timeframe:**Baseline; Week 24

---

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	195	90		
Units: percentage change				
least squares mean (standard error)	-47.79 (± 2.43)	-17.18 (± 3.53)		

### Statistical analyses

<b>Statistical analysis title</b>	mixed-effect model; repeated measurement
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	285
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.0001
Method	mixed-effect model; repeated measurement
Parameter estimate	least squares mean difference
Point estimate	-30.61
Confidence interval	
level	95 %
sides	2-sided
lower limit	-39.03
upper limit	-22.19
Variability estimate	Standard error of the mean
Dispersion value	4.28

### Secondary: Percentage change from Baseline in F-VASI at Week 52

End point title	Percentage change from Baseline in F-VASI at Week 52
End point description:	<p>F-VASI was measured by the percentage of vitiligo involvement (percentage of BSA) and the degree of depigmentation: 0% (no depigmentation), 10% (only specks of depigmentation), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment), or 100% (no pigment). The percentage of BSA (hand unit) vitiligo involvement was estimated to the nearest 0.1% by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate the percentage of BSA vitiligo involvement. F-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site on the face and summing the values of all sites (possible range: 0-3; lower scores indicate increased improvement). Percentage change = ([post-BL value minus BL value]/BL value) X 100.</p>
End point type	Secondary
End point timeframe:	Baseline; Week 52

<b>End point values</b>	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	173	82		
Units: percentage change				
arithmetic mean (standard deviation)	-67.24 (± 33.660)	-52.98 (± 30.174)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage change from Baseline in F-BSA at Week 52

End point title	Percentage change from Baseline in F-BSA at Week 52
-----------------	---

End point description:

F-BSA involvement was the proportion of the facial body surface area with vitiligo. The area "Face" was defined as including the area on the forehead to the original hairline, on the cheek to the jawline vertically to the jawline and laterally from the corner of the mouth to the tragus. The area "Face" did not include surface area of the lips, scalp, ears, or neck, but included the nose and eyelids. Body surface area assessment was performed by the Palmar Method. Body surface area was estimated to the nearest 0.1%. The approximate size of the participant's entire palmar surface (i.e., the palm plus 5 digits) was considered as 1% BSA, and the approximate size of the participant's thumb was considered as 0.1% BSA. Percentage change = ([post-BL value minus BL value]/BL value) X 100.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 52

<b>End point values</b>	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	173	82		
Units: percentage change				
arithmetic mean (standard deviation)	-44.87 (± 43.954)	-32.40 (± 30.068)		

## Statistical analyses

**Secondary: Percentage change from Baseline in T-VASI at Week 24**

End point title	Percentage change from Baseline in T-VASI at Week 24
End point description:	T-VASI was calculated with contributions from 6 sites. The percentage of vitiligo involvement was estimated in hand units (percentage of BSA estimated to the nearest 0.1%) by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate percent BSA vitiligo involvement. The degree of depigmentation for each site was estimated to the nearest percentage: 0% (no depigmentation present), 10% (only specks of depigmentation present), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment present), 100% (no pigment present). T-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site and summing the values (range: 0-100; lower scores indicate increased improvement). Percentage change = ([post-BL value minus BL value]/BL value) X 100.
End point type	Secondary
End point timeframe:	Baseline; Week 24

End point values	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	195	90		
Units: percentage change				
least squares mean (standard error)	-27.60 ( $\pm$ 1.81)	-10.62 ( $\pm$ 2.64)		

**Statistical analyses**

<b>Statistical analysis title</b>	mixed-effect model; repeated measurement
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	285
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.0001
Method	mixed-effect model; repeated measurement
Parameter estimate	least squares mean difference
Point estimate	-16.98
Confidence interval	
level	95 %
sides	2-sided
lower limit	-23.28
upper limit	-10.68
Variability estimate	Standard error of the mean
Dispersion value	3.2

---

**Secondary: Percentage change from Baseline in T-VASI at Week 52**

---

End point title	Percentage change from Baseline in T-VASI at Week 52
-----------------	--

End point description:

T-VASI was calculated with contributions from 6 sites. The percentage of vitiligo involvement was estimated in hand units (percentage of BSA estimated to the nearest 0.1%) by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate percent BSA vitiligo involvement. The degree of depigmentation for each site was estimated to the nearest percentage: 0% (no depigmentation present), 10% (only specks of depigmentation present), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment present), 100% (no pigment present). T-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site and summing the values (range: 0-100; lower scores indicate increased improvement). Percentage change =  $([\text{post-BL value minus BL value}]/\text{BL value}) \times 100$ .

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 52

---

End point values	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	173	82		
Units: percentage change				
arithmetic mean (standard deviation)	-49.23 ( $\pm$ 26.366)	-29.85 ( $\pm$ 37.832)		

---

**Statistical analyses**

---

No statistical analyses for this end point

---

---

**Secondary: Percentage change from Baseline in T-BSA at Week 24**

---

End point title	Percentage change from Baseline in T-BSA at Week 24
-----------------	---

End point description:

T-BSA involvement was the proportion of the body surface area with vitiligo. Body surface area assessment was performed by the Palmar Method. Body surface area was estimated to the nearest 0.1%. The approximate size of the participant's entire palmar surface (i.e., the palm plus 5 digits) was considered as 1% BSA, and the approximate size of the participant's thumb was considered as 0.1% BSA. Percentage change =  $([\text{post-BL value minus BL value}]/\text{BL value}) \times 100$ .

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 24

---

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	195	90		
Units: percentage change				
least squares mean (standard error)	-13.08 (± 1.40)	-4.02 (± 2.05)		

### Statistical analyses

<b>Statistical analysis title</b>	mixed-effect model; repeated measurement
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	285
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.0003
Method	mixed-effect model; repeated measurement
Parameter estimate	least squares mean difference
Point estimate	-9.07
Confidence interval	
level	95 %
sides	2-sided
lower limit	-13.96
upper limit	-4.18
Variability estimate	Standard error of the mean
Dispersion value	2.49

### Secondary: Percentage change from Baseline in T-BSA at Week 52

End point title	Percentage change from Baseline in T-BSA at Week 52
End point description:	T-BSA involvement was the proportion of the body surface area with vitiligo. Body surface area assessment was performed by the Palmar Method. Body surface area was estimated to the nearest 0.1%. The approximate size of the participant's entire palmar surface (i.e., the palm plus 5 digits) was considered as 1% BSA, and the approximate size of the participant's thumb was considered as 0.1% BSA. Percentage change = ([post-BL value minus BL value]/BL value) X 100.
End point type	Secondary
End point timeframe:	
Baseline; Week 52	

<b>End point values</b>	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	173	82		
Units: percentage change				
arithmetic mean (standard deviation)	-27.39 (± 25.705)	-11.83 (± 34.654)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of participants achieving a $\geq 25\%$ , $\geq 75\%$ , and $\geq 90\%$ Improvement in the Total Body Vitiligo Area Scoring Index (T-VASI25/75/90) Score at Week 24

End point title	Percentage of participants achieving a $\geq 25\%$ , $\geq 75\%$ , and $\geq 90\%$ Improvement in the Total Body Vitiligo Area Scoring Index (T-VASI25/75/90) Score at Week 24
-----------------	--

End point description:

A T-VASI25/75/90 responder achieved at least 25/75/90% improvement from Baseline in T-VASI, calculated with contributions from 6 sites. The percentage of vitiligo involvement was estimated in hand units (percentage of BSA estimated to nearest 0.1%) by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate percent BSA vitiligo involvement. The degree of depigmentation for each site was estimated to the nearest percentage: 0% (no depigmentation present), 10% (only specks of depigmentation present), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment present), 100% (no pigment present). T-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site and summing the values (range: 0-100; lower scores indicate increased improvement).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 24

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	221	109		
Units: percentage of participants				
number (not applicable)				
T-VASI25	48.8	23.8		
T-VASI75	4.1	1.8		
T-VASI90	0.5	0.0		

## Statistical analyses

<b>Statistical analysis title</b>	T-VASI25; exact logistic regression
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	< 0.0001 [8]
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	3.04
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.746
upper limit	5.307

Notes:

[8] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

<b>Statistical analysis title</b>	T-VASI75; exact logistic regression
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.2921 [9]
Method	exact logistic regression
Parameter estimate	Odds ratio (OR)
Point estimate	2.3
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.489
upper limit	10.823

Notes:

[9] - The model included the treatment group (1.5% BID and vehicle) and stratification factors (skin type and region).

<b>Statistical analysis title</b>	T-VASI90
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID

Number of subjects included in analysis	330
Analysis specification	Pre-specified
Analysis type	
Parameter estimate	Odds ratio (OR)
Point estimate	0.49
Confidence interval	
level	95 %
sides	2-sided
lower limit	-9999
upper limit	9999

**Secondary: Percentage of participants achieving a  $\geq 25\%$ ,  $\geq 50\%$ ,  $75\%$ , and  $\geq 90\%$  Improvement in the Total Body Vitiligo Area Scoring Index (T-VASI25/50/75/90) Score at Week 52**

End point title	Percentage of participants achieving a $\geq 25\%$ , $\geq 50\%$ , $75\%$ , and $\geq 90\%$ Improvement in the Total Body Vitiligo Area Scoring Index (T-VASI25/50/75/90) Score at Week 52
-----------------	--

End point description:

A T-VASI25/50/75/90 responder achieved  $\geq 25/50/75/90\%$  improvement from Baseline in T-VASI, calculated with contributions from 6 sites. The percentage of vitiligo involvement was estimated in hand units (percentage of BSA estimated to nearest 0.1%) by the Investigator using the Palmar Method. The Investigator used his/her hand to mimic the participant's hand size to evaluate percent BSA vitiligo involvement. The degree of depigmentation for each site was estimated to the nearest percentage: 0% (no depigmentation present), 10% (only specks of depigmentation present), 25% (pigmented area exceeded depigmented area), 50% (depigmented and pigmented area was equal), 75% (depigmented area exceeded pigmented area), 90% (specks of pigment present), 100% (no pigment present). T-VASI was then derived by multiplying the values assessed for the vitiligo involvement by the percentage of affected skin for each site and summing the values (range: 0-100; lower scores indicate increased improvement).

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 52

<b>End point values</b>	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	173	82		
Units: percentage of participants				
number (not applicable)				
T-VASI25	77.5	56.1		
T-VASI50	53.2	31.7		
T-VASI75	20.2	9.8		
T-VASI90	3.5	2.4		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of participants in each category of VNS during the treatment period (Double-Blind and Treatment-Extension Periods)

End point title	Percentage of participants in each category of VNS during the treatment period (Double-Blind and Treatment-Extension Periods)
-----------------	---

End point description:

The VNS is a patient-reported measure of vitiligo treatment success that is rated on a 5-point scale. The Baseline facial photograph was shown to the participants for reference, and a mirror was provided for the participants to assess the vitiligo on their face. The participant was asked to respond to the following query: Compared with before treatment, how noticeable is the vitiligo now? Responses: (1) more noticeable, (2) as noticeable, (3) slightly less noticeable, (4) a lot less noticeable, and (5) no longer noticeable.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 24 and Week 52

End point values	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID
Subject group type	Reporting group	Reporting group	Subject analysis set	Subject analysis set
Number of subjects analysed	195 <sup>[10]</sup>	90 <sup>[11]</sup>	173 <sup>[12]</sup>	82 <sup>[13]</sup>
Units: percentage of participants				
number (not applicable)				
Week 24, more noticeable	6.2	14.4	9999	9999
Week 24, as noticeable	17.4	46.7	9999	9999
Week 24, slightly less noticeable	51.3	35.6	9999	9999
Week 24, a lot less noticeable	24.1	3.3	9999	9999
Week 24, no longer noticeable	1.0	0.0	9999	9999
Week 52, more noticeable	9999	9999	4.0	4.9
Week 52, as noticeable	9999	9999	9.2	15.9
Week 52, slightly less noticeable	9999	9999	46.8	59.8
Week 52, a lot less noticeable	9999	9999	39.3	19.5
Week 52, no longer noticeable	9999	9999	0.6	0.0

Notes:

[10] - 9999=participants in this treatment group weren't analyzed at this time point.

[11] - 9999=participants in this treatment group weren't analyzed at this time point.

[12] - 9999=participants in this treatment group weren't analyzed at this time point.

[13] - 9999=participants in this treatment group weren't analyzed at this time point.

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in Dermatology Life Quality Index (DLQI) at Week 24

End point title	Change from Baseline in Dermatology Life Quality Index (DLQI)
-----------------	---

## End point description:

The DLQI is a 10-question validated questionnaire for use in participants aged 16 years and over to measure how much the skin problem has affected the participant over the previous 7 days. Each question is scored as: very much = 3; a lot = 2; a little = 1; not at all = 0; not relevant = 0. For Question 7, "Prevented work or studying" = 3. The DLQI was calculated by summing the score of each question, resulting in a maximum of 30 and a minimum of 0. The higher the score, the more quality of life is impaired.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline; Week 24

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	178	87		
Units: scores on a scale				
least squares mean (standard error)	-1.17 (± 0.27)	-0.85 (± 0.39)		

**Statistical analyses**

<b>Statistical analysis title</b>	mixed-effect model; repeated measurement
Comparison groups	Double-Blind Period: Ruxolitinib cream 1.5% BID v Double-Blind Period: Vehicle cream BID
Number of subjects included in analysis	265
Analysis specification	Pre-specified
Analysis type	
P-value	= 0.497
Method	mixed-effect model; repeated measurement
Parameter estimate	least squares mean difference
Point estimate	-0.32
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.26
upper limit	0.62
Variability estimate	Standard error of the mean
Dispersion value	0.48

**Secondary: Change from Baseline in DLQI at Week 52**

End point title	Change from Baseline in DLQI at Week 52
-----------------	---

## End point description:

The DLQI is a 10-question validated questionnaire for use in participants aged 16 years and over to measure how much the skin problem has affected the participant over the previous 7 days. Each

question is scored as: very much = 3; a lot = 2; a little = 1; not at all = 0; not relevant = 0. For Question 7, "Prevented work or studying" = 3. The DLQI was calculated by summing the score of each question, resulting in a maximum of 30 and a minimum of 0. The higher the score, the more quality of life is impaired.

End point type	Secondary
End point timeframe:	
Baseline; Week 52	

End point values	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID
Subject group type	Reporting group	Reporting group	Subject analysis set	Subject analysis set
Number of subjects analysed	204 <sup>[14]</sup>	105 <sup>[15]</sup>	204 <sup>[16]</sup>	105 <sup>[17]</sup>
Units: scores on a scale				
arithmetic mean (standard deviation)				
Baseline, n=204, 105	4.63 (± 4.446)	4.59 (± 4.871)	9999 (± 9999)	9999 (± 9999)
Week 52, n=157, 79	9999 (± 9999)	9999 (± 9999)	-1.40 (± 4.087)	-1.37 (± 3.617)

Notes:

[14] - 9999=participants in this treatment group weren't analyzed at this time point.

[15] - 9999=participants in this treatment group weren't analyzed at this time point.

[16] - 9999=participants in this treatment group weren't analyzed at this time point.

[17] - 9999=participants in this treatment group weren't analyzed at this time point.

### Statistical analyses

No statistical analyses for this end point

### Secondary: Change from Baseline in Children's Dermatology Life Quality Index (CDLQI) during the treatment period (Double-Blind and Treatment-Extension Periods)

End point title	Change from Baseline in Children's Dermatology Life Quality Index (CDLQI) during the treatment period (Double-Blind and Treatment-Extension Periods)
-----------------	--

End point description:

The DLQI is a 10-question validated questionnaire for use in participants aged 16 years and over to measure how much the skin problem has affected the participant over the previous 7 days. The CDLQI is the youth/children's version of the DLQI and was completed by adolescents aged ≥ 12 years to < 16 years. Each question is scored as: very much = 3; quite a lot = 2; only a little = 1; not at all = 0; question unanswered = 0. For Question 7: "Prevented school" = 3. The CDLQI was calculated by summing the score of each question, resulting in a maximum of 30 and a minimum of 0. The higher the score, the more quality of life is impaired.

End point type	Secondary
End point timeframe:	
Baseline; Week 24 and Week 52	

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Double-Blind Period: Vehicle cream BID	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID
Subject group type	Reporting group	Reporting group	Subject analysis set	Subject analysis set
Number of subjects analysed	17 <sup>[18]</sup>	4 <sup>[19]</sup>	17 <sup>[20]</sup>	4 <sup>[21]</sup>
Units: scores on a scale				
arithmetic mean (standard deviation)				
Baseline, n=16, 4	2.50 (± 2.805)	1.25 (± 1.893)	9999 (± 9999)	9999 (± 9999)
Week 24, n=16, 3	-0.25 (± 2.113)	0.00 (± 0.000)	9999 (± 9999)	9999 (± 9999)
Week 52, n=15, 3	9999 (± 9999)	9999 (± 9999)	-1.00 (± 2.507)	0.00 (± 1.000)

Notes:

[18] - 9999=participants in this treatment group weren't analyzed at this time point.

[19] - 9999=participants in this treatment group weren't analyzed at this time point.

[20] - 9999=participants in this treatment group weren't analyzed at this time point.

[21] - 9999=participants in this treatment group weren't analyzed at this time point.

## Statistical analyses

No statistical analyses for this end point

## Secondary: Trough plasma concentrations of ruxolitinib at Weeks 4, 24, and 40

End point title	Trough plasma concentrations of ruxolitinib at Weeks 4, 24, and 40 <sup>[22]</sup>
-----------------	--

End point description:

Trough plasma concentration was defined as the measurement of the plasma concentration of ruxolitinib before drug application.

End point type	Secondary
----------------	-----------

End point timeframe:

pre-dose at Weeks 4, 24, and 40

Notes:

[22] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Pharmacokinetics was assessed for ruxolitinib only; thus, data are not reported for the following arm: Double-Blind Period: Vehicle cream BID. Furthermore, no statistical analysis was conducted for this endpoint.

<b>End point values</b>	Double-Blind Period: Ruxolitinib cream 1.5% BID	Treatment-Extension (TE) Period: Ruxolitinib cream 1.5% BID	TE Period: Vehicle cream to Ruxolitinib cream 1.5% BID	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	206 <sup>[23]</sup>	173 <sup>[24]</sup>	80 <sup>[25]</sup>	
Units: nanomoles				
arithmetic mean (standard deviation)				
Week 4, n=206, 0, 0	57.1 (± 61.4)	9999 (± 9999)	9999 (± 9999)	
Week 24, n=191, 0, 0	56.3 (± 69.4)	9999 (± 9999)	9999 (± 9999)	
Week 40, n=0, 173, 80	9999 (± 9999)	55.5 (± 63.6)	50.1 (± 55.8)	

---

Notes:

[23] - 9999=participants in this treatment group weren't analyzed at this time point.

[24] - 9999=participants in this treatment group weren't analyzed at this time point.

[25] - 9999=participants in this treatment group weren't analyzed at this time point.

---

## **Statistical analyses**

No statistical analyses for this end point

## Adverse events

### Adverse events information<sup>[1]</sup>

Timeframe for reporting adverse events:

from the time of Informed Consent Form signing until at least 30 days after the last application of study drug (up to Week 52 + 30 days)

Adverse event reporting additional description:

Treatment-emergent adverse events (TEAEs): AEs reported for the first time or the worsening of a pre-existing event after the first application of study drug. For the Double-Blind Period, TEAEs are reported for members of the Safety Population. For the Treatment-Extension (TE) Period, TEAEs are reported for the TE Evaluable Population.

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	22
--------------------	----

### Reporting groups

Reporting group title	Vehicle cream BID
-----------------------	-------------------

Reporting group description:

Participants applied matching vehicle cream twice a day (BID) for 24 weeks in the Double-Blind Period.

Reporting group title	Ruxolitinib cream 1.5% BID
-----------------------	----------------------------

Reporting group description:

Participants applied ruxolitinib cream during the Double-Blind Treatment Period and the Treatment-Extension Period. Participants applied ruxolitinib 1.5% cream BID for 24 weeks. Participants who completed the Week 24 assessments with no safety concerns could continue into the 28-week Treatment-Extension Period. Participants who applied ruxolitinib cream 1.5% BID during the Double-Blind Period continued to apply ruxolitinib cream 1.5% BID for an additional 28 weeks in the Treatment-Extension Period. Participants who applied vehicle cream BID during the Double-Blind Period applied ruxolitinib cream 1.5% BID for 28 weeks in the Treatment-Extension Period.

Notes:

[1] - There are no non-serious adverse events recorded for these results. It is expected that there will be at least one non-serious adverse event reported.

Justification: No non-serious adverse events occurred in at least 5% of participants in either treatment arm.

<b>Serious adverse events</b>	Vehicle cream BID	Ruxolitinib cream 1.5% BID	
Total subjects affected by serious adverse events			
subjects affected / exposed	1 / 109 (0.92%)	8 / 311 (2.57%)	
number of deaths (all causes)	0	0	
number of deaths resulting from adverse events	0	0	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Prostate cancer			
subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Injury, poisoning and procedural complications			
Concussion			

subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Kidney contusion			
subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Tibia fracture			
subjects affected / exposed	1 / 109 (0.92%)	0 / 311 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Cardiac disorders			
Myocarditis			
subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Nervous system disorders			
Subacute combined cord degeneration			
subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Immune system disorders			
Hypersensitivity			
subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Gastrointestinal disorders			
Anal fistula			
subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
Appendicitis			

subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Hepatitis infectious mononucleosis</b>			
subjects affected / exposed	0 / 109 (0.00%)	1 / 311 (0.32%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	Vehicle cream BID	Ruxolitinib cream 1.5% BID	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	0 / 109 (0.00%)	0 / 311 (0.00%)	

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
12 December 2019	The primary purpose of this amendment was to incorporate revisions requested by the Voluntary Harmonisation Procedure (VHP). <ul style="list-style-type: none"><li>- Added language to exclude participant who had current and/or history of tuberculosis</li><li>- Added language to exclude participants who lived with anyone participating in any current Incyte-sponsored ruxolitinib cream study</li><li>- Added language to instruct that German participants whose hemoglobin was between 10 grams per deciliter (g/dL) and 10.5 g/dL during the screening visit should have been further evaluated per local guidelines before enrolling into the study</li></ul>
21 February 2020	The primary purpose of this amendment was to incorporate revisions requested by the German and French Ethics Committees (ECs) and FDA. <ul style="list-style-type: none"><li>- Reordered and revised the key secondary endpoints and updated the analysis plan</li><li>- Added 1 key exclusion criteria (exclude other forms of vitiligo) to the Population section, added information about the exit interview in the Study Design section, and added a Data and Safety Monitoring Board (DSMB) section to indicate that a DSMB was not required in this study</li><li>- Added language that targeted physical examination should have been conducted as indicated by symptoms reported by the participant, adverse events (AEs), or other findings. Abnormalities that were considered clinically significant in the judgment of the investigator were to be reported as AEs.</li><li>- Added language to instruct that adolescent participants screened in Germany whose hemoglobin was between 10 g/dL and 12 g/dL during the screening visit should have been further evaluated per local guidelines before being enrolled into the study</li></ul>

Notes:

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported