



Clinical trial results:

A phase III, randomized, controlled, observer-blind study to demonstrate effectiveness, immunogenicity and safety of GSK's meningococcal Group B and combined ABCWY vaccines when administered to healthy adolescents and young adults.

Summary

EudraCT number	2019-001666-15
Trial protocol	FI CZ EE Outside EU/EEA
Global end of trial date	13 September 2022

Results information

Result version number	v4 (current)
This version publication date	30 June 2024
First version publication date	27 March 2023
Version creation reason	

Trial information

Trial identification

Sponsor protocol code	205416
-----------------------	--------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	-
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	GlaxoSmithKline
Sponsor organisation address	Rue de l'Institut, 89, Rixensart, Belgium, 1330
Public contact	GSK Response Center, GlaxoSmithKline, 044 8664357343, GSKClinicalSupportHD@gsk.com
Scientific contact	GSK Response Center, GlaxoSmithKline, 044 8664357343, GSKClinicalSupportHD@gsk.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	Yes

Notes:

Results analysis stage

Analysis stage	Final
Date of interim/final analysis	24 October 2022
Is this the analysis of the primary completion data?	Yes
Primary completion date	13 September 2022
Global end of trial reached?	Yes
Global end of trial date	13 September 2022
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

•Effectiveness of rMenB+OMV NZ and MenABCWY vaccines -against a panel of N.meningitidis serogroup B strains at 1 month(M) after the 3 and 2-dose rMenB+OMV NZ series and last MenABCWY dose when compared to 1 M after MenACWY dose -As the percentages of subjects whose sera kill $\geq 70\%$ of strains tested using enc-hSBA at 1 M after the 3 and 2-dose rMenB+OMV NZ series and 1 M after last MenABCWY dose •Lot-lot consistency of immune responses of 3 lots of MenACWY component of MenABCWY vaccine,as measured by hSBA GMTs at 1 M after last dose •Immunological non-inferiority: MenABCWY versus MenACWY as measured by percentages of subjects achieving a 4-fold rise in hSBA titers at 1 M after last MenABCWY dose and 1 M after MenACWY dose •Effectiveness non-inferiority: MenABCWY versus rMenB+OMV NZ in terms of percentage of samples with bactericidal serum activity at 1 M after last ABCWY dose and 1 M after 3 or 2 dose rMenB+OMV series •Safety and reactogenicity of MenB,MenABCWY and MenACWY vaccines

Protection of trial subjects:

Vaccine administration is to be preceded by a review of the participants medical history (including previous vaccination and possible occurrence of undesirable events) and a general physical examination at the first visit and symptom-directed physical examination before subsequent vaccinations. Protocol procedures including blood sampling were to be done by a qualified healthcare professional. Vaccines/products will be administered only to eligible participants who had no contraindications to any components of the vaccines/products. Participants will be followed-up for 6 months after third vaccination/product administration. The participants will be observed closely for at least 30 minutes following the administration of the vaccine(s)/product(s), with appropriate medical treatment readily available in case of anaphylaxis and/or syncope.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	14 August 2020
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	No

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Australia: 295
Country: Number of subjects enrolled	Canada: 229
Country: Number of subjects enrolled	Czechia: 750
Country: Number of subjects enrolled	Estonia: 127
Country: Number of subjects enrolled	Finland: 819
Country: Number of subjects enrolled	Türkiye: 333
Country: Number of subjects enrolled	United States: 1085
Worldwide total number of subjects	3638
EEA total number of subjects	1696

Notes:

Subjects enrolled per age group	
In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	702
Adolescents (12-17 years)	1459
Adults (18-64 years)	1477
From 65 to 84 years	0
85 years and over	0

Subject disposition

Recruitment

Recruitment details:

As pre-specified in protocol: -Participant flow, Baseline characteristics, Effectiveness, and immunogenicity data are presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Lot-to-lot consistency analysis data are presented for individual ABCWY lot groups (ABCWY-1, ABCWY-2 and ABCWY-3).

Pre-assignment

Screening details:

Out of 3657 participants enrolled, 19 participants did not receive vaccination as they did not meet the eligibility criteria, therefore only 3638 participants were included in the Exposed Set and started the study.

Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind ^[1]
Roles blinded	Monitor, Data analyst

Blinding implementation details:

Observer-blinded study. Recipients & study evaluators will be unaware of vaccine administered.

Arms

Are arms mutually exclusive?	Yes
Arm title	MenB_0_2_6 Group

Arm description:

Participants received 3 doses of rMenB+OMV NZ vaccine on Day 1, Day 61 and Day 181. Participants received 1 dose of MenACWY vaccine at Day 211 as a standard of care.

Arm type	Experimental
Investigational medicinal product name	Meningococcal Groups A, C, W and Y Conjugate Vaccine (MenACWY)
Investigational medicinal product code	
Other name	Menveo
Pharmaceutical forms	Powder and solution for solution for injection
Routes of administration	Intramuscular use

Dosage and administration details:

1 dose of MenACWY vaccine at Day 211

Investigational medicinal product name	Meningococcal Group B Vaccine (rMenB+OMV NZ)
Investigational medicinal product code	
Other name	Bexsero
Pharmaceutical forms	Suspension for injection
Routes of administration	Intramuscular use

Dosage and administration details:

3 doses of rMenB+OMV NZ vaccine at Day 1, Day 61 and Day 181

Arm title	MenB_0_6 Group
------------------	----------------

Arm description:

Participants received 2 doses of rMenB+OMV NZ vaccine on Day 1, and Day 181, 1 dose of MenACWY vaccine on Day 61. Participants received 1 dose of Placebo on Day 211 to maintain blinding.

Arm type	Experimental
----------	--------------

Investigational medicinal product name	Meningococcal Group B Vaccine (rMenB+OMV NZ)
Investigational medicinal product code	
Other name	Bexsero
Pharmaceutical forms	Suspension for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
2 doses of rMenB+OMV NZ vaccine at Day 1 and Day 181	
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
1 dose of Placebo at Day 211	
Investigational medicinal product name	Meningococcal Groups A, C, W and Y Conjugate Vaccine (MenACWY)
Investigational medicinal product code	
Other name	Menveo
Pharmaceutical forms	Powder and solution for solution for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
1 dose of MenACWY vaccine at Day 61	
Arm title	ABCWY_Pooled
Arm description:	
Participants received 2 doses of either MenABCWY Lot 1, Lot 2, or Lot 3 vaccine on Day 1 and Day 181 and 1 dose of placebo on Day 61. Participants received 1 dose of placebo on Day 211 to maintain blinding. To evaluate the effectiveness of 2 doses of the MenABCWY vaccines against rMenB+OMV and MenACWY vaccines, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group.	
Arm type	Experimental
Investigational medicinal product name	Combined Meningococcal Groups A, B, C, W and Y vaccine (MenABCWY)
Investigational medicinal product code	MenABCWY
Other name	
Pharmaceutical forms	Powder and suspension for suspension for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
2 doses of MenABCWY vaccine at Day 1 and Day 181	
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
2 dose of Placebo at Day 61 and Day 211	
Arm title	ACWY Group
Arm description:	
Participants received 1 dose of MenACWY vaccine at Day 1, 1 dose of placebo at Day 61 and 1 dose of rMenB+OMV NZ vaccine on Day 181. Participants received 1 dose of rMenB+OMV NZ vaccine on Day 211 as standard of care.	
Arm type	Active comparator

Investigational medicinal product name	Meningococcal Groups A, C, W and Y Conjugate Vaccine (MenACWY)
Investigational medicinal product code	
Other name	Menveo
Pharmaceutical forms	Powder and solution for solution for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
1 dose of MenACWY vaccine at Day 1	
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
1 dose of Placebo at Day 61	
Investigational medicinal product name	Meningococcal Group B Vaccine (rMenB+OMV NZ)
Investigational medicinal product code	
Other name	Bexsero
Pharmaceutical forms	Suspension for injection
Routes of administration	Intramuscular use
Dosage and administration details:	
2 doses of rMenB+OMV NZ vaccine at Day 181 and Day 211	
Notes:	
[1] - The roles blinded appear to be inconsistent with a double blind trial.	
Justification: This is an Observer-blinded study. Recipients & study evaluators were unaware of the vaccine administered.	

Number of subjects in period 1	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled
Started	897	906	1657
ABCWY-1 Group	0	0	549
ABCWY-2 Group	0	0	554
ABCWY-3 Group	0	0	554
Completed	797	811	1497
Not completed	100	95	160
Consent withdrawn by subject	46	38	54
Adverse event, non-fatal	7	6	11
Not specified	1	-	1
MIGRATED / MOVED FROM THE STUDY AREA	6	7	9
Lost to follow-up	32	36	69
Protocol deviation	8	8	16

Number of subjects in period 1	ACWY Group
Started	178
ABCWY-1 Group	0
ABCWY-2 Group	0
ABCWY-3 Group	0

Completed	163
Not completed	15
Consent withdrawn by subject	7
Adverse event, non-fatal	1
Not specified	1
MIGRATED / MOVED FROM THE STUDY AREA	1
Lost to follow-up	4
Protocol deviation	1

Baseline characteristics

Reporting groups

Reporting group title	MenB_0_2_6 Group
Reporting group description:	
Participants received 3 doses of rMenB+OMV NZ vaccine on Day 1, Day 61 and Day 181. Participants received 1 dose of MenACWY vaccine at Day 211 as a standard of care.	
Reporting group title	MenB_0_6 Group
Reporting group description:	
Participants received 2 doses of rMenB+OMV NZ vaccine on Day 1, and Day 181, 1 dose of MenACWY vaccine on Day 61. Participants received 1 dose of Placebo on Day 211 to maintain blinding.	
Reporting group title	ABCWY_Pooled
Reporting group description:	
Participants received 2 doses of either MenABCWY Lot 1, Lot 2, or Lot 3 vaccine on Day 1 and Day 181 and 1 dose of placebo on Day 61. Participants received 1 dose of placebo on Day 211 to maintain blinding. To evaluate the effectiveness of 2 doses of the MenABCWY vaccines against rMenB+OMV and MenACWY vaccines, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group.	
Reporting group title	ACWY Group
Reporting group description:	
Participants received 1 dose of MenACWY vaccine at Day 1, 1 dose of placebo at Day 61 and 1 dose of rMenB+OMV NZ vaccine on Day 181. Participants received 1 dose of rMenB+OMV NZ vaccine on Day 211 as standard of care.	

Reporting group values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled
Number of subjects	897	906	1657
Age categorical			
Units: Subjects			
In utero	0	0	0
Preterm newborn infants (gestational age < 37 wks)	0	0	0
Newborns (0-27 days)	0	0	0
Infants and toddlers (28 days-23 months)	0	0	0
Children (2-11 years)	183	172	320
Adolescents (12-17 years)	349	368	666
Adults (18-64 years)	365	366	671
From 65-84 years	0	0	0
85 years and over	0	0	0
Age continuous			
Units: years			
arithmetic mean	16.5	16.5	16.5
standard deviation	± 4.7	± 4.7	± 4.7
Sex: Female, Male			
Units: Participants			
Female	464	446	933
Male	433	460	724
Race/Ethnicity, Customized			
Units: Subjects			
American Indian or Alaska Native	5	5	3
Asian	43	60	71
Black or African American	33	29	59

Native Hawaiian or Other Pacific Islander	3	1	3
Other, Unspecified	17	20	29
White	796	791	1492

Reporting group values	ACWY Group	Total	
Number of subjects	178	3638	
Age categorical Units: Subjects			
In utero	0	0	
Preterm newborn infants (gestational age < 37 wks)	0	0	
Newborns (0-27 days)	0	0	
Infants and toddlers (28 days-23 months)	0	0	
Children (2-11 years)	27	702	
Adolescents (12-17 years)	76	1459	
Adults (18-64 years)	75	1477	
From 65-84 years	0	0	
85 years and over	0	0	
Age continuous Units: years			
arithmetic mean	16.9		
standard deviation	± 4.6	-	
Sex: Female, Male Units: Participants			
Female	100	1943	
Male	78	1695	
Race/Ethnicity, Customized Units: Subjects			
American Indian or Alaska Native	0	13	
Asian	9	183	
Black or African American	6	127	
Native Hawaiian or Other Pacific Islander	0	7	
Other, Unspecified	1	67	
White	162	3241	

End points

End points reporting groups

Reporting group title	MenB_0_2_6 Group
Reporting group description: Participants received 3 doses of rMenB+OMV NZ vaccine on Day 1, Day 61 and Day 181. Participants received 1 dose of MenACWY vaccine at Day 211 as a standard of care.	
Reporting group title	MenB_0_6 Group
Reporting group description: Participants received 2 doses of rMenB+OMV NZ vaccine on Day 1, and Day 181, 1 dose of MenACWY vaccine on Day 61. Participants received 1 dose of Placebo on Day 211 to maintain blinding.	
Reporting group title	ABCWY_Pooled
Reporting group description: Participants received 2 doses of either MenABCWY Lot 1, Lot 2, or Lot 3 vaccine on Day 1 and Day 181 and 1 dose of placebo on Day 61. Participants received 1 dose of placebo on Day 211 to maintain blinding. To evaluate the effectiveness of 2 doses of the MenABCWY vaccines against rMenB+OMV and MenACWY vaccines, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group.	
Reporting group title	ACWY Group
Reporting group description: Participants received 1 dose of MenACWY vaccine at Day 1, 1 dose of placebo at Day 61 and 1 dose of rMenB+OMV NZ vaccine on Day 181. Participants received 1 dose of rMenB+OMV NZ vaccine on Day 211 as standard of care.	
Subject analysis set title	ABCWY-1 Group
Subject analysis set type	Per protocol
Subject analysis set description: Participants received 2 doses of MenABCWY lot 1 vaccine at Day 1 and Day 181 and 2 doses of placebo at Day 61 and Day 211.	
Subject analysis set title	ABCWY-2 Group
Subject analysis set type	Per protocol
Subject analysis set description: Participants received 2 doses of MenABCWY lot 2 vaccine at Day 1 and Day 181 and 2 doses of placebo at Day 61 and Day 211.	
Subject analysis set title	ABCWY-3 Group
Subject analysis set type	Per protocol
Subject analysis set description: Participants received 2 doses of MenABCWY lot 3 vaccine at Day 1 and Day 181 and 2 doses of placebo at Day 61 and Day 211.	

Primary: Percentage of blood samples without bactericidal serum activity against each of the endemic US N. meningitidis serogroup B strains at 1 month after the 3-dose (0,2,6-M), 2-dose(0,6-M) vaccination schedule of rMenB+OMV and 1 dose of MenACWY

End point title	Percentage of blood samples without bactericidal serum activity against each of the endemic US N. meningitidis serogroup B strains at 1 month after the 3-dose (0,2,6-M), 2-dose(0,6-M) vaccination schedule of rMenB+OMV and 1 dose of MenACWY ^[1]
-----------------	--

End point description:

The effectiveness (test-based) of rMenB+OMV vaccine at 1 month after the 3 doses in MenB_0_2_6 group and 1 month after the 2 dose schedule in MenB_0_6 group when compared to one dose of MenACWY vaccination in ACWY group, against a panel of N. meningitidis serogroup B strains was measured in terms of percentage of samples without bactericidal activity using endogenous complement human Serum Bactericidal Assay (enc-hSBA), which provides a qualitative assessment (yes/no) of the presence of sufficient bactericidal antibodies in human sera to kill a meningococcal strain at a specific dilution of 1:4. Participants were randomly selected for testing against each strain therefore only a subset of participants were tested for each of the strains. Number of Participants analyzed = Total number of participants included in PPS.

Analysis was performed on blood samples collected from Per Protocol Set (PPS).

End point type	Primary			
End point timeframe:				
At 1 month after vaccination schedule (i.e., Day 211 for MenB_0_2_6 group [3-dose schedule] and MenB_0_6 group, and Day 31 for ACWY group)				
Notes:				
[1] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: As specified in the Protocol, the analysis assesses the effectiveness of the MenABCWY vaccine compared to the rMenB+OMV vaccine in terms of the percentage of samples with bactericidal serum activity.				
End point values	MenB_0_2_6 Group	MenB_0_6 Group	ACWY Group	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	740	751	147	
Units: Percentage of blood samples				
number (not applicable)				
Number of Blood samples (N=25596,26142,4374)	13.3	14.4	79	

Statistical analyses

Statistical analysis title	Statistical analysis 2
Statistical analysis description:	
To demonstrate the effectiveness of the rMenB+OMV NZ vaccine against a randomly selected panel of endemic US N. meningitidis serogroup B invasive disease strains as measured by bactericidal activity using enc-hSBA at 1 month after the 2-dose (0,6-M) schedule in MenB_0_6 group when compared to 1 month after the MenACWY dose in the ACWY group.	
Comparison groups	MenB_0_6 Group v ACWY Group
Number of subjects included in analysis	898
Analysis specification	Pre-specified
Analysis type	other ^[2]
Parameter estimate	VE
Point estimate	81.8
Confidence interval	
level	Other: 97.5 %
sides	2-sided
lower limit	80.4
upper limit	83.1

Notes:

[2] - Effectiveness of rMenB+OMV NZ vaccine is demonstrated if the LL of the 2-sided 97.5% CI for VE against the selected strain panel between the MenB_0_6 and the ACWY groups is above 65%. VE is defined as $1 - RR = (1 - \text{percentage of samples without bactericidal serum activity at 1:4 dilution in MenB group} / \text{percentage of samples without bactericidal serum activity at 1:4 dilution in the ACWY group}) \times 100$ percentage.

Statistical analysis title	Statistical analysis 1
Statistical analysis description:	
To demonstrate the effectiveness of the rMenB+OMV NZ vaccine against a randomly selected panel of endemic US N. meningitidis serogroup B invasive disease strains as measured by bactericidal activity using enc-hSBA at 1 month after the 3-dose (0,2,6-months) schedule in MenB_0_2_6 group when compared to 1 month after the MenACWY dose in the ACWY group.	
Comparison groups	ACWY Group v MenB_0_2_6 Group

Number of subjects included in analysis	887
Analysis specification	Pre-specified
Analysis type	other ^[3]
Parameter estimate	VE (Vaccine Effectiveness)
Point estimate	83.2
Confidence interval	
level	Other: 97.5 %
sides	2-sided
lower limit	81.9
upper limit	84.4

Notes:

[3] - Effectiveness of rMenB+OMV NZ vaccine is demonstrated if the LL of the 2-sided 97.5% CI for Vaccine Effectiveness (VE) against the selected strain panel between the MenB_0_2_6 and the ACWY groups is above 65%. VE is defined as $1 - \text{Risk Ratio (RR)} = (1 - \text{percentage of samples without bactericidal serum activity at 1:4 dilution in MenB group} / \text{percentage of samples without bactericidal serum activity at 1:4 dilution in the ACWY group}) \times 100 \text{ percentage}$.

Primary: Percentage of blood samples without bactericidal serum activity against each of the endemic US N. meningitidis serogroup B strains at 1 month after the 2-dose (0,2-M) vaccination schedule of rMenB+OMV and 1 dose of MenACWY

End point title	Percentage of blood samples without bactericidal serum activity against each of the endemic US N. meningitidis serogroup B strains at 1 month after the 2-dose (0,2-M) vaccination schedule of rMenB+OMV and 1 dose of MenACWY ^[4]
-----------------	---

End point description:

The effectiveness (test-based) of rMenB+OMV vaccine at 1 month after the 2 doses in MenB_0_2_6 group when compared to one dose of MenACWY vaccination in ACWY group, against a panel of N. meningitidis serogroup B strains was measured in terms of percentage of samples without bactericidal activity using endogenous complement human Serum Bactericidal Assay (enc-hSBA), which provides a qualitative assessment (yes/no) of the presence of sufficient bactericidal antibodies in human sera to kill a meningococcal strain at a specific dilution of 1:4. Participants were randomly selected for testing against each strain therefore only a subset of participants were tested for each of the strains. Number of Participants analyzed = Total number of participants included in PPS.
Analysis was performed on blood samples collected from Per Protocol Set (PPS).

End point type	Primary
----------------	---------

End point timeframe:

At 1 month after vaccination schedule (i.e., Day 91 for the MenB_0_2_6 group [2-dose schedule] and Day 31 for ACWY group)

Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: As specified in the Protocol, the analysis assesses the effectiveness of the MenABCWY vaccine compared to the rMenB+OMV vaccine in terms of the percentage of samples with bactericidal serum activity.

End point values	MenB_0_2_6 Group	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	740	147		
Units: Percentage of blood samples				
number (not applicable)				
Number of Blood samples (N=27569,4374)	16.8	79		

Statistical analyses

Statistical analysis title	Statistical analysis 1
Statistical analysis description:	
To demonstrate the effectiveness of the rMenB+OMV NZ vaccine against a randomly selected panel of endemic US N. meningitidis serogroup B invasive disease strains as measured by bactericidal activity using enc-hSBA at 1 month after the 2-dose (0,2-M) schedule in MenB_0_2_6 group when compared to 1 month after the MenACWY dose in the ACWY group.	
Comparison groups	MenB_0_2_6 Group v ACWY Group
Number of subjects included in analysis	887
Analysis specification	Pre-specified
Analysis type	other ^[5]
Parameter estimate	VE
Point estimate	78.7
Confidence interval	
level	Other: 97.5 %
sides	2-sided
lower limit	77.2
upper limit	80.1

Notes:

[5] - Effectiveness of rMenB+OMV NZ vaccine is demonstrated if the LL of the 2-sided 97.5% CI for VE against the selected strain panel between the MenB_0_2_6 and the ACWY groups is above 65%. VE is defined as $1 - \text{RR} = (1 - \text{percentage of samples without bactericidal serum activity at 1:4 dilution in MenB group} / \text{percentage of samples without bactericidal serum activity at 1:4 dilution in the ACWY group}) \times 100 \text{ percentage}$.

Primary: Geometric mean titers (GMTs) against serogroups A, C, W and Y for each lot (ABCWY-1 Group, ABCWY-2 Group and ABCWY-3 Group) at 1 month after the last vaccination of MenABCWY

End point title	Geometric mean titers (GMTs) against serogroups A, C, W and Y for each lot (ABCWY-1 Group, ABCWY-2 Group and ABCWY-3 Group) at 1 month after the last vaccination of MenABCWY
-----------------	---

End point description:

Immune response was measured in terms of hSBA GMTs directed against serogroups A, C, W and Y. As pre- specified in the protocol, the data reported in this outcome measures data were presented for individual lots to demonstrate the consistency of the immune response of 3 lots (ABCWY- 1 Group, ABCWY-2 Group, and ABCWY-3 Group) of the ACWY component of the MenABCWY vaccine. Analysis was performed on PPS.

End point type	Primary
----------------	---------

End point timeframe:

At 1 month after the last vaccination of MenABCWY (Day 211)

End point values	ABCWY-1 Group	ABCWY-2 Group	ABCWY-3 Group	
Subject group type	Subject analysis set	Subject analysis set	Subject analysis set	
Number of subjects analysed	452	449	458	
Units: Titers				
geometric mean (confidence interval 95%)				
Men A (N=448,443,454)	336.4 (299.3 to 378.0)	349.9 (311.5 to 393.0)	390.4 (347.4 to 438.8)	
Men C (N= 448,449,456)	1036.7 (877.6 to 1224.5)	1130.2 (958.1 to 1333.4)	888.4 (752.1 to 1049.2)	
Men W (N= 452,449,458)	564.5 (497.9 to 639.9)	635.5 (561.0 to 719.9)	640.1 (564.6 to 725.6)	
Men Y (N=451,449,457)	536.7 (464.5 to 620.2)	623.9 (540.4 to 720.2)	644.3 (557.6 to 744.6)	

Statistical analyses

Statistical analysis title	Statistical analysis 1
Statistical analysis description: To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-2 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup A at 1 month after last vaccination (Day 211).	
Comparison groups	ABCWY-1 Group v ABCWY-2 Group
Number of subjects included in analysis	901
Analysis specification	Pre-specified
Analysis type	equivalence ^[6]
Parameter estimate	GMT ratio
Point estimate	0.96
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.84
upper limit	1.1

Notes:

[6] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y are within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 2
Statistical analysis description: To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup A at 1 month after last vaccination (Day 211).	
Comparison groups	ABCWY-1 Group v ABCWY-3 Group
Number of subjects included in analysis	910
Analysis specification	Pre-specified
Analysis type	equivalence ^[7]
Parameter estimate	GMT ratio
Point estimate	0.86
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.75
upper limit	0.98

Notes:

[7] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y is within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 5
Statistical analysis description: To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup C at 1 month after last vaccination (Day 211).	

Comparison groups	ABCWY-1 Group v ABCWY-3 Group
Number of subjects included in analysis	910
Analysis specification	Pre-specified
Analysis type	equivalence ^[8]
Parameter estimate	GMT ratio
Point estimate	1.17
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.97
upper limit	1.41

Notes:

[8] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y is within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 4
-----------------------------------	------------------------

Statistical analysis description:

To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-2 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup C at 1 month after last vaccination (Day 211).

Comparison groups	ABCWY-1 Group v ABCWY-2 Group
Number of subjects included in analysis	901
Analysis specification	Pre-specified
Analysis type	equivalence ^[9]
Parameter estimate	GMT ratio
Point estimate	0.92
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.76
upper limit	1.11

Notes:

[9] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y are within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 3
-----------------------------------	------------------------

Statistical analysis description:

To demonstrate lot-to-lot consistency of the immune responses of ABCWY-2 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup A at 1 month after last vaccination (Day 211).

Comparison groups	ABCWY-2 Group v ABCWY-3 Group
Number of subjects included in analysis	907
Analysis specification	Pre-specified
Analysis type	equivalence ^[10]
Parameter estimate	GMT ratio
Point estimate	0.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.78
upper limit	1.02

Notes:

[10] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y is within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 6
Statistical analysis description: To demonstrate lot-to-lot consistency of the immune responses of ABCWY-2 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup C at 1 month after last vaccination (Day 211).	
Comparison groups	ABCWY-2 Group v ABCWY-3 Group
Number of subjects included in analysis	907
Analysis specification	Pre-specified
Analysis type	equivalence ^[11]
Parameter estimate	GMT ratio
Point estimate	1.27
Confidence interval	
level	95 %
sides	2-sided
lower limit	1.05
upper limit	1.54

Notes:

[11] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y is within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 10
Statistical analysis description: To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-2 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup Y at 1 month after last vaccination (Day 211).	
Comparison groups	ABCWY-1 Group v ABCWY-2 Group
Number of subjects included in analysis	901
Analysis specification	Pre-specified
Analysis type	equivalence ^[12]
Parameter estimate	GMT ratio
Point estimate	0.86
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.73
upper limit	1.01

Notes:

[12] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y are within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 7
Statistical analysis description: To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-2 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup W at 1 month after last vaccination (Day 211).	
Comparison groups	ABCWY-1 Group v ABCWY-2 Group

Number of subjects included in analysis	901
Analysis specification	Pre-specified
Analysis type	equivalence ^[13]
Parameter estimate	GMT ratio
Point estimate	0.89
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.77
upper limit	1.02

Notes:

[13] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y are within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 8
-----------------------------------	------------------------

Statistical analysis description:

To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup W at 1 month after last vaccination (Day 211).

Comparison groups	ABCWY-1 Group v ABCWY-3 Group
Number of subjects included in analysis	910
Analysis specification	Pre-specified
Analysis type	
Parameter estimate	GMT ratio
Point estimate	0.88
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.77
upper limit	1.02

Statistical analysis title	Statistical analysis 9
-----------------------------------	------------------------

Statistical analysis description:

To demonstrate lot-to-lot consistency of the immune responses of ABCWY-2 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup W at 1 month after last vaccination (Day 211).

Comparison groups	ABCWY-2 Group v ABCWY-3 Group
Number of subjects included in analysis	907
Analysis specification	Pre-specified
Analysis type	equivalence ^[14]
Parameter estimate	GMT ratio
Point estimate	0.99
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.86
upper limit	1.14

Notes:

[14] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y is within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 11
Statistical analysis description:	
To demonstrate lot-to-lot consistency of the immune responses of ABCWY-1 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup Y at 1 month after last vaccination (Day 211).	
Comparison groups	ABCWY-1 Group v ABCWY-3 Group
Number of subjects included in analysis	910
Analysis specification	Pre-specified
Analysis type	equivalence ^[15]
Parameter estimate	GMT ratio
Point estimate	0.83
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.71
upper limit	0.98

Notes:

[15] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y is within the [0.5;2.0] equivalence interval for each pair of lots.

Statistical analysis title	Statistical analysis 12
Statistical analysis description:	
To demonstrate lot-to-lot consistency of the immune responses of ABCWY-2 and ABCWY-3 lots of the MenACWY component of the MenABCWY vaccine, as measured by hSBA GMTs directed against serogroup Y at 1 month after last vaccination (Day 211).	
Comparison groups	ABCWY-2 Group v ABCWY-3 Group
Number of subjects included in analysis	907
Analysis specification	Pre-specified
Analysis type	equivalence ^[16]
Parameter estimate	GMT ratio
Point estimate	0.97
Confidence interval	
level	95 %
sides	2-sided
lower limit	0.82
upper limit	1.14

Notes:

[16] - Lot-to-lot consistency is claimed if the 2-sided 95% CIs for the ratio of hSBA GMTs of antibodies against each of the serogroups A, C, W and Y is within the [0.5;2.0] equivalence interval for each pair of lots.

Primary: Percentage of participants whose sera kill $\geq 70\%$ of the strains tested using enc-hSBA at 1 month after the 2-dose (0,2-M) schedule of rMenB+OMV

End point title	Percentage of participants whose sera kill $\geq 70\%$ of the strains tested using enc-hSBA at 1 month after the 2-dose (0,2-M) schedule of rMenB+OMV ^{[17][18]}
-----------------	---

End point description:

The effectiveness (responder-based) of the rMenB+OMV NZ vaccine was measured in terms of percentage of participants whose sera kill $\geq 70\%$ of the strains tested using enc-hSBA, calculated based on Clopper Pearson method. Effectiveness is demonstrated if Lower Limit (LL) of the two-sided 97.5% CI for the percentages of subjects whose sera kill $\geq 70\%$ of strains is above 65%. Analysis was performed on the FAS.

End point type	Primary			
End point timeframe:				
At 1 month after vaccination schedule (i.e., Day 91 for the MenB_0_2_6 group [2-dose schedule])				
Notes:				
[17] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.				
Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.				
[18] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.				
Justification: As specified in the Protocol, the analysis assesses the effectiveness of the rMenB+OMV vaccine by assessing the percentages of subjects whose sera kill $\geq 70\%$ of strains tested.				
End point values	MenB_0_2_6 Group			
Subject group type	Reporting group			
Number of subjects analysed	831			
Units: Percentage of participants				
number (confidence interval 97.5%)	84.8 (81.8 to 87.5)			

Statistical analyses

No statistical analyses for this end point

Primary: Percentage of participants whose sera kill Greater Than or Equal to (\geq) 70% of the strains tested using enc-hSBA at 1 month after the 3-dose (0,2,6-M) schedule of rMenB+OMV and 2-dose(0,6-M) schedule of rMenB+OMV

End point title	Percentage of participants whose sera kill Greater Than or Equal to (\geq) 70% of the strains tested using enc-hSBA at 1 month after the 3-dose (0,2,6-M) schedule of rMenB+OMV and 2-dose(0,6-M) schedule of rMenB+OMV ^{[19][20]}
-----------------	---

End point description:

The effectiveness (responder-based) of the rMenB+OMV NZ vaccine was measured in terms of percentage of participants whose sera kill $\geq 70\%$ of the strains tested using enc-hSBA, calculated based on Clopper Pearson method. Effectiveness is demonstrated if Lower Limit (LL) of the two-sided 97.5% CI for the percentages of subjects whose sera kill $\geq 70\%$ of strains is above 65%. Analysis was performed on the Full Analysis Set (FAS).

End point type	Primary
----------------	---------

End point timeframe:

At 1 month after vaccination schedule (i.e., Day 211 for MenB_0_2_6 group [3-dose schedule] and MenB_0_6 group)

Notes:

[19] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

[20] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the effectiveness of the rMenB+OMV vaccine by assessing the percentages of subjects whose sera kill $\geq 70\%$ of strains tested.

End point values	MenB_0_2_6 Group	MenB_0_6 Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	790	813		
Units: Percentage of participants				
number (confidence interval 97.5%)	93.4 (91.2 to 95.2)	89.8 (87.2 to 92)		

Statistical analyses

No statistical analyses for this end point

Primary: Percentage of participants with 4-fold rise in hSBA titers against N. meningitidis serogroups A, C, W and Y at 1 month after last MenABCWY vaccination (pooled lots) and MenACWY vaccination (for the ACWY Group), relative to baseline

End point title	Percentage of participants with 4-fold rise in hSBA titers against N. meningitidis serogroups A, C, W and Y at 1 month after last MenABCWY vaccination (pooled lots) and MenACWY vaccination (for the ACWY Group), relative to baseline ^[21]
-----------------	---

End point description:

Four-fold rise is defined as: If the pre-vaccination hSBA titer is < 4, then post-vaccination hSBA titer should be ≥ 16 . If the pre-vaccination hSBA titer is \geq limit of detection (LOD) but < LL of quantification (LLOQ), then post-vaccination hSBA titer should be ≥ 4 times the LLOQ. If the pre-vaccination hSBA titer is \geq LLOQ, then post-vaccination hSBA titer should be ≥ 4 times the pre-vaccination hSBA titer. As pre-specified in the protocol, data reported in this outcome measure were presented for ACWY group, ABCWY pooled group to evaluate the immunological non-inferiority of 2 doses of the MenABCWY vaccines against MenACWY vaccines, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on PPS.

End point type	Primary
----------------	---------

End point timeframe:

At 1 month after vaccination schedule (i.e., Day 211 for the ABCWY_Pooled group and Day 31 for the ACWY Group) compared to Day 1 (baseline)

Notes:

[21] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the effectiveness of the rMenB+OMV vaccine compared to one dose of MenACWY vaccination in the ACWY group

End point values	ABCWY_Pooled	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1196	119		
Units: Percentage of participants				
number (confidence interval 95%)				
Men A (N=1170,112)	97.0 (95.9 to 97.9)	85.7 (77.8 to 91.6)		
Men C (N=1189,114)	97.2 (96.1 to 98.1)	50.0 (40.5 to 59.5)		
Men W (N=1185,115)	97.0 (95.9 to 97.9)	61.7 (52.2 to 70.6)		
Men Y (N=1196,119)	96.7 (95.6 to 97.7)	69.7 (60.7 to 77.8)		

Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description:	
To demonstrate the immunological non- inferiority of the MenABCWY vaccine compared to the MenACWY vaccine in participants without a previous MenACWY vaccination (unprimed) as measured by the percentages of participants, achieving a 4- fold rise in hSBA titers against N. meningitidis serogroup A at 1 month after the last MenABCWY vaccination (Day 211) and 1 month after the MenACWY vaccination.	
Comparison groups	ABCWY_Pooled v ACWY Group
Number of subjects included in analysis	1315
Analysis specification	Pre-specified
Analysis type	non-inferiority ^[22]
Parameter estimate	Difference in percentage of participants
Point estimate	11.29
Confidence interval	
level	95 %
sides	2-sided
lower limit	5.88
upper limit	19.01

Notes:

[22] - Non-inferiority of MenABCWY vaccine is demonstrated if the LL of the 2-sided 95% CI for the difference in percentage of participants with 4-fold rise between the 2 groups is above -10%.

Statistical analysis title	Statistical Analysis 2
Statistical analysis description:	
To demonstrate the immunological non- inferiority of the MenABCWY vaccine compared to the MenACWY vaccine in participants without a previous MenACWY vaccination (unprimed) as measured by the percentages of participants, achieving a 4- fold rise in hSBA titers against N. meningitidis serogroup C at 1 month after the last MenABCWY vaccination (Day 211) and 1 month after the MenACWY vaccination.	
Comparison groups	ABCWY_Pooled v ACWY Group
Number of subjects included in analysis	1315
Analysis specification	Pre-specified
Analysis type	non-inferiority ^[23]
Parameter estimate	Difference in percentage of participants
Point estimate	47.22
Confidence interval	
level	95 %
sides	2-sided
lower limit	38.14
upper limit	56.3

Notes:

[23] - Non-inferiority of MenABCWY vaccine is demonstrated if the LL of the 2-sided 95% CI for the difference in percentage of participants with 4-fold rise between the 2 groups is above -10%.

Statistical analysis title	Statistical Analysis 3
Statistical analysis description:	
To demonstrate the immunological non- inferiority of the MenABCWY vaccine compared to the MenACWY vaccine in participants without a previous MenACWY vaccination (unprimed) as measured by the percentages of participants, achieving a 4- fold rise in hSBA titers against N. meningitidis serogroup W at 1 month after the last MenABCWY vaccination (Day 211) and 1 month after the MenACWY vaccination.	
Comparison groups	ABCWY_Pooled v ACWY Group

Number of subjects included in analysis	1315
Analysis specification	Pre-specified
Analysis type	non-inferiority ^[24]
Parameter estimate	Difference in percentage of participants
Point estimate	35.31
Confidence interval	
level	95 %
sides	2-sided
lower limit	26.88
upper limit	44.49

Notes:

[24] - Non-inferiority of MenABCWY vaccine is demonstrated if the LL of the 2-sided 95% CI for the difference in percentage of participants with 4-fold rise between the 2 groups is above -10%.

Statistical analysis title	Statistical Analysis 4
-----------------------------------	------------------------

Statistical analysis description:

To demonstrate the immunological non- inferiority of the MenABCWY vaccine compared to the MenACWY vaccine in participants without a previous MenACWY vaccination (unprimed) as measured by the percentages of participants, achieving a 4- fold rise in hSBA titers against N. meningitidis serogroup Y at 1 month after the last MenABCWY vaccination (Day 211) and 1 month after the MenACWY vaccination.

Comparison groups	ABCWY_Pooled v ACWY Group
Number of subjects included in analysis	1315
Analysis specification	Pre-specified
Analysis type	non-inferiority ^[25]
Parameter estimate	Difference in percentage of participants
Point estimate	26.99
Confidence interval	
level	95 %
sides	2-sided
lower limit	19.38
upper limit	35.81

Notes:

[25] - Non-inferiority of MenABCWY vaccine is demonstrated if the LL of the 2-sided 95% CI for the difference in percentage of participants with 4-fold rise between the 2 groups is above -10%.

Primary: Percentage of blood samples with bactericidal serum activity against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after the last MenABCWY dose (pooled lots) and 2-dose(0,2-M) schedule of rMenB+OMV

End point title	Percentage of blood samples with bactericidal serum activity against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after the last MenABCWY dose (pooled lots) and 2-dose(0,2-M) schedule of rMenB+OMV ^[26]
-----------------	--

End point description:

The effectiveness was measured in terms of percentage of samples with bactericidal activity using enc-hSBA, which provides a qualitative assessment (yes/no) of the presence of sufficient bactericidal antibodies in human sera to kill a meningococcal strain at a specific dilution of 1:4. As pre-specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 Group and ABCWY pooled group to evaluate the effectiveness of 2 doses of the MenABCWY vaccines against MenACWY vaccines, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on blood samples collected from PPS, which included participants who received at least 1 dose of the study treatment to which they were randomized and have post-vaccination data for the specified analysis at specified timepoints and did not have any protocol deviations that lead to exclusion from the PPS.

End point type	Primary
----------------	---------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 211 for the ABCWY_Pooled group and Day 91 for the MenB_0_2_6 Group [2-dose schedule])

Notes:

[26] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the effectiveness of the MenABCWY vaccine compared to the rMenB+OMV vaccine in terms of the percentage of samples with bactericidal serum activity.

End point values	MenB_0_2_6 Group	ABCWY_Pooled		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	740	1356		
Units: Percentage of blood samples				
number (not applicable)				
Number of Blood samples Analyzed (27569, 25715)	83.1	82.5		

Statistical analyses

Statistical analysis title	Statistical analysis 1
----------------------------	------------------------

Statistical analysis description:

To demonstrate the non-inferiority of the effectiveness of the MenABCWY vaccine (0,6-months schedule) compared to the rMenB+OMV NZ vaccine (0,2-months) in terms of percentage of samples with bactericidal serum activity using enc-hSBA against a randomly selected panel of endemic US N. meningitidis serogroup B invasive disease strains.

Comparison groups	MenB_0_2_6 Group v ABCWY_Pooled
Number of subjects included in analysis	2096
Analysis specification	Pre-specified
Analysis type	non-inferiority ^[27]
Parameter estimate	Difference in percentage of participants
Point estimate	-0.61
Confidence interval	
level	95 %
sides	2-sided
lower limit	-1.25
upper limit	0.03

Notes:

[27] - Non-inferiority of MenABCWY to rMenB+OMV NZ is demonstrated if LL of the 2-sided 95% CI for the difference in percentages of samples with bactericidal serum activity at 1:4 dilution is above -5%.

Primary: Percentage of blood samples without bactericidal serum activity against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after last MenABCWY vaccination (pooled lots) and MenACWY vaccination (for the ACWY Group)

End point title	Percentage of blood samples without bactericidal serum activity against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after last MenABCWY vaccination (pooled lots) and MenACWY vaccination (for the ACWY Group) ^[28]
-----------------	---

End point description:

The effectiveness (test-based) of 2 doses of MenABCWY vaccine when compared to 1 dose of MenACWY vaccine, against a panel of N. meningitidis serogroup B strains was measured in terms of percentage of samples without bactericidal activity using enc-hSBA, which provides a qualitative assessment (yes/no) of the presence of sufficient bactericidal antibodies in human sera to kill a meningococcal strain at a specific dilution of 1:4. As pre-specified in the protocol, data reported in this outcome measure were presented for ACWY group and ABCWY pooled group to evaluate the effectiveness of 2 doses of the MenABCWY vaccines against MenACWY vaccines, participants from the ABCWY-1, ABCWY-2, and

ABCWY- 3 groups were pooled into a single group. Analysis was performed on blood samples collected from PPS.

End point type	Primary
----------------	---------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 211 for the ABCWY_Pooled group and Day 31 for the ACWY group)

Notes:

[28] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the effectiveness of the rMenB+OMV vaccine compared to one dose of MenACWY vaccination in the ACWY group

End point values	ABCWY_Pooled	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1356	147		
Units: Percentage of blood samples				
number (not applicable)				
Number of Blood samples analyzed (N=25715,4374)	17.4	79		

Statistical analyses

Statistical analysis title	Statistical analysis 1
----------------------------	------------------------

Statistical analysis description:

To demonstrate the effectiveness of the MenABCWY vaccine against a randomly selected panel of endemic US N. meningitidis serogroup B invasive disease strains as measured by enc-hSBA at 1 month after the last MenABCWY vaccination (Day 211) when compared to 1 month after the MenACWY vaccination.

Comparison groups	ABCWY_Pooled v ACWY Group
Number of subjects included in analysis	1503
Analysis specification	Pre-specified
Analysis type	other ^[29]
Parameter estimate	VE
Point estimate	77.9
Confidence interval	
level	95 %
sides	2-sided
lower limit	76.6
upper limit	79.2

Notes:

[29] - Effectiveness of MenABCWY vaccine is demonstrated if the LL of the 2-sided 95% CI for VE against the selected strain panel between the ABCWY and the ACWY groups is above 65%. VE is defined as $1 - RR = (1 - \text{percentage of samples without bactericidal serum activity at 1:4 dilution in ABCWY_Pooled group} / \text{percentage of samples without bactericidal serum activity at 1:4 dilution in the ACWY group}) \times 100$ percentage.

Primary: Percentage of participants whose sera kill $\geq 70\%$ of the strains tested using enc-hSBA at 1 month after the last vaccination in the ABCWY Group (pooled lots)

End point title	Percentage of participants whose sera kill $\geq 70\%$ of the strains tested using enc-hSBA at 1 month after the last vaccination in the ABCWY Group (pooled lots) ^{[30][31]}
-----------------	--

End point description:

The effectiveness (responder-based) of the MenABCWY vaccine is measured in terms of percentage of participants whose sera kill $\geq 70\%$ of the strains tested using enc-hSBA, being calculated based on Clopper Pearson method. Effectiveness is demonstrated Lower Limit (LL) of the two-sided 97.5% CI for the percentages of subjects whose sera kill $\geq 70\%$ of strains tested for MenABCWY is above 65%. Analysis was performed on the FAS.

End point type	Primary
----------------	---------

End point timeframe:

At 1 month after the last vaccination of MenABCWY (Day 211)

Notes:

[30] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

[31] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the effectiveness of the MenABCWY vaccine by assessing the percentages of subjects whose sera kill $\geq 70\%$ of strains tested.

End point values	ABCWY_Pooled			
Subject group type	Reporting group			
Number of subjects analysed	817			
Units: Percentage of participants				
number (confidence interval 95%)	84.1 (81.4 to 86.5)			

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any solicited local adverse events (AEs) after the first study intervention administration

End point title	Number of participants with any solicited local adverse events (AEs) after the first study intervention administration ^[32]
-----------------	--

End point description:

Assessed solicited local adverse events were injection or administration site pain, erythema, swelling, induration. Any = occurrence of the adverse event regardless of intensity grade. Any erythema and swelling = adverse event reported with a surface diameter greater than 0 millimeters. No solicited AEs were collected after vaccination on Day 211 because these vaccines were administered as part of standard of care and to maintain blinding. As pre- specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Solicited Safety Set (SSS).

End point type	Primary
----------------	---------

End point timeframe:

During 7 days after the first study intervention administration occurring at Day 1

Notes:

[32] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	885	894	1638	178
Units: participants				
Pain	807	819	1503	67
Erythema	90	86	216	11
Swelling	87	89	217	11
Induration	60	64	150	7

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any solicited local adverse events (AEs) after the second study intervention administration

End point title	Number of participants with any solicited local adverse events (AEs) after the second study intervention administration ^[33]
-----------------	---

End point description:

Assessed solicited local adverse events were injection or administration site pain, erythema, swelling, induration. Any = occurrence of the adverse event regardless of intensity grade. Any erythema and swelling = adverse event reported with a surface diameter greater than 0 millimeters. No solicited AEs were collected after vaccination on Day 211 because these vaccines were administered as part of standard of care and to maintain blinding. As pre- specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Solicited Safety Set (SSS).

End point type	Primary
----------------	---------

End point timeframe:

During 7 days after the second study intervention administration occurring at Day 61

Notes:

[33] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	823	813	1511	161
Units: participants				
Pain	714	224	205	30
Erythema	89	26	5	1
Swelling	99	22	5	1
Induration	67	19	6	0

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any solicited local adverse events (AEs) after

the third study intervention administration

End point title	Number of participants with any solicited local adverse events (AEs) after the third study intervention administration ^[34]
-----------------	--

End point description:

Assessed solicited local adverse events were injection site pain, erythema, swelling, induration. Any = occurrence of the adverse event regardless of intensity grade. Any erythema and swelling = adverse event reported with a surface diameter greater than 0 millimeters. No solicited AEs were collected after vaccination on Day 211 because these vaccines were administered as part of standard of care and to maintain blinding. As pre- specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Solicited Safety Set (SSS).

End point type	Primary
----------------	---------

End point timeframe:

During 7 days after the third study intervention administration occurring at Day 181

Notes:

[34] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	765	759	1428	148
Units: participants				
Pain	677	676	1258	126
Erythema	118	87	168	11
Swelling	107	85	176	13
Induration	52	57	114	12

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any solicited systemic AEs after the first study intervention administration

End point title	Number of participants with any solicited systemic AEs after the first study intervention administration ^[35]
-----------------	--

End point description:

Assessed solicited systemic AEs were fatigue, nausea, myalgia, arthralgia, headache and fever [temperature $\geq 38.0^{\circ}\text{C}$]. Any = occurrence of the adverse event regardless of intensity grade or relation to study vaccination. No solicited AEs were collected after vaccination on Day 211 because these vaccines were administered as part of standard of care and to maintain blinding. As pre- specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Solicited Safety Set (SSS).

End point type	Primary
----------------	---------

End point timeframe:

During 7 days after the first study intervention administration occurring at Day 1

Notes:

[35] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	885	894	1638	178
Units: participants				
Fatigue	423	414	828	78
Nausea	112	111	242	27
Myalgia	92	106	242	13
Arthralgia	56	70	133	17
Headache	358	330	681	69
Fever (C)	19	17	55	3

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any solicited systemic AEs after the second study intervention administration

End point title	Number of participants with any solicited systemic AEs after the second study intervention administration ^[36]
-----------------	---

End point description:

Assessed solicited systemic AEs were fatigue, nausea, myalgia, arthralgia, headache and fever [temperature $\geq 38.0^{\circ}\text{C}$]. Any = occurrence of the adverse event regardless of intensity grade or relation to study vaccination. No solicited AEs were collected after vaccination on Day 211 because these vaccines were administered as part of standard of care and to maintain blinding. As pre- specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Solicited Safety Set (SSS).

End point type	Primary
----------------	---------

End point timeframe:

During 7 days after the second study intervention administration occurring at Day 61

Notes:

[36] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	823	813	1511	161
Units: participants				
Fatigue	372	228	345	36
Nausea	104	56	85	18
Myalgia	110	46	55	3
Arthralgia	72	332	35	6
Headache	301	223	332	31
Fever (C)	22	12	18	1

Statistical analyses

Primary: Number of participants with any solicited systemic AEs after the third study intervention administration

End point title	Number of participants with any solicited systemic AEs after the third study intervention administration ^[37]
-----------------	--

End point description:

Assessed solicited systemic AEs were fatigue, nausea, myalgia, arthralgia, headache and fever [temperature $\geq 38.0^{\circ}\text{C}$]. Any = occurrence of the adverse event regardless of intensity grade or relation to study vaccination. No solicited AEs were collected after vaccination on Day 211 because these vaccines were administered as part of standard of care and to maintain blinding. As pre-specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Solicited Safety Set (SSS).

End point type	Primary
----------------	---------

End point timeframe:

During 7 days after the third study intervention administration occurring at Day 181

Notes:

[37] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	765	759	1428	148
Units: participants				
Fatigue	374	341	602	56
Nausea	94	84	147	14
Myalgia	106	109	168	17
Arthralgia	71	53	104	7
Headache	302	284	509	39
Fever (C)	21	23	27	2

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any unsolicited AEs after the first study intervention administration

End point title	Number of participants with any unsolicited AEs after the first study intervention administration ^[38]
-----------------	---

End point description:

Unsolicited AEs are defined as any AE reported in addition to those solicited during the clinical study. Also, any 'solicited' symptom with onset outside the specified period of follow-up for solicited symptoms were reported as an unsolicited AE. Any Unsolicited AEs (including Serious adverse events (SAEs), AEs leading to withdrawal, Adverse event of special interest (AESIs) and medically attended AEs were collected during 30 days after vaccination 1-3. For vaccination on Day 211 (Vaccination 4), conducted as part of standard of care and to maintain blinding, no eDiary data or all unsolicited AEs were collected for the 30 days following vaccination. However, SAEs, AEs leading to withdrawal, AESIs and medically attended AEs were collected throughout study duration. As pre-specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Unsolicited Safety Set (USS).

End point type	Primary
----------------	---------

End point timeframe:

During the 30 days after the first study intervention administration occurring at Day 1

Notes:

[38] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	893	900	1648	178
Units: participants	90	124	217	29

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any unsolicited AEs after the second study intervention administration

End point title	Number of participants with any unsolicited AEs after the second study intervention administration ^[39]
-----------------	--

End point description:

Unsolicited AEs are defined as any AE reported in addition to those solicited during the clinical study. Also, any 'solicited' symptom with onset outside the specified period of follow-up for solicited symptoms were reported as an unsolicited AE. Any Unsolicited AEs (including Serious adverse events (SAEs), AEs leading to withdrawal, Adverse event of special interest (AESIs) and medically attended AEs were collected during 30 days after vaccination 1-3. For vaccination on Day 211 (Vaccination 4), conducted as part of standard of care and to maintain blinding, no eDiary data or all unsolicited AEs were collected for the 30 days following vaccination. However, SAEs, AEs leading to withdrawal, AESIs and medically attended AEs were collected throughout study duration. As pre-specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Unsolicited Safety Set (USS).

End point type	Primary
----------------	---------

End point timeframe:

During the 30 days after the second study intervention administration occurring at Day 61

Notes:

[39] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	851	855	1579	170
Units: participants	106	88	160	15

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with any unsolicited AEs after the third study intervention administration

End point title	Number of participants with any unsolicited AEs after the third study intervention administration ^[40]
-----------------	---

End point description:

Unsolicited AEs are defined as any AE reported in addition to those solicited during the clinical study. Also, any 'solicited' symptom with onset outside the specified period of follow-up for solicited symptoms were reported as an unsolicited AE. Any Unsolicited AEs (including Serious adverse events (SAEs), AEs leading to withdrawal, Adverse event of special interest (AESIs) and medically attended AEs were collected during 30 days after vaccination 1-3. For vaccination on Day 211 (Vaccination 4), conducted as part of standard of care and to maintain blinding, no eDiary data or all unsolicited AEs were collected for the 30 days following vaccination. However, SAEs, AEs leading to withdrawal, AESIs and medically attended AEs were collected throughout study duration. As pre-specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Unsolicited Safety Set (USS).

End point type	Primary
----------------	---------

End point timeframe:

During the 30 days after the third study intervention administration occurring at Day 181

Notes:

[40] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	815	823	1521	166
Units: participants	96	94	183	19

Statistical analyses

No statistical analyses for this end point

Primary: Number of participants with SAEs, AEs leading to withdrawal, AESIs and medically attended AEs

End point title	Number of participants with SAEs, AEs leading to withdrawal, AESIs and medically attended AEs ^[41]
-----------------	---

End point description:

A SAEs is any untoward medical occurrence that results in death, is life-threatening, requires hospitalization or prolongation of existing hospitalization, results in disability/incapacity and is a congenital anomaly/birth defect in the offspring of a study subject. AESIs are predefined (serious or non-serious) AEs of scientific and medical concern specific to the product or program, for which ongoing monitoring and rapid communication by the investigator to the sponsor can be appropriate, because such an event might warrant further investigation to characterize and understand it. Medically attended AEs are symptoms or illnesses requiring hospitalization, or emergency room visit, or visit to/by a health care provider. As pre-specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 group, MenB_0_6 group, ACWY group, ABCWY pooled group. Analysis was performed on the Unsolicited Safety Set (USS).

End point type	Primary
----------------	---------

End point timeframe:

Throughout the study period (Day 1 to Day 361)

Notes:

[41] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: The analysis of this primary endpoint was descriptive i.e. no statistical hypothesis test was performed.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	893	900	1648	178
Units: participants				
SAEs	20	22	25	5
AEs leading to withdrawal	6	4	4	1
AESIs	1	1	6	0
medically attended AEs	238	288	479	44

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants with 4-fold rise in hSBA titers against N.meningitidis group B strains at 1 month after last MenABCWY dose (ABCWY group-pooled lots) and 1 month after 2-dose(0,2-M) schedule of rMenB+OMV NZ relative to baseline

End point title	Percentage of participants with 4-fold rise in hSBA titers against N.meningitidis group B strains at 1 month after last MenABCWY dose (ABCWY group-pooled lots) and 1 month after 2-dose(0,2-M) schedule of rMenB+OMV NZ relative to baseline ^[42]
-----------------	---

End point description:

The immunogenicity is measured as percentage of participants achieving a 4-fold rise in hSBA titers against N. meningitidis serogroup B indicator strains (M14459, 96217, M13520 and NZ98/254 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively). 4-fold rise per each indicator strain was defined as a post-vaccination hSBA titre ≥ 16 for participants with a pre-vaccination hSBA titre < 4 ; a post-vaccination hSBA titre ≥ 4 times the LLOQ for participants with a pre-vaccination hSBA titre $\geq \text{LOD}$ and $< \text{LLOQ}$; a post-vaccination hSBA titre ≥ 4 times the pre-vaccination hSBA titre for participants with a pre-vaccination hSBA titre $\geq \text{LLOQ}$. As pre-specified in the protocol, data reported in this outcome measure were presented for MenB_0_2_6 Group and ABCWY pooled group to evaluate the effectiveness of 2 doses of the MenABCWY vaccines against MenACWY vaccines, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. The analysis was performed on the PPS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 211 for the ABCWY_Pooled group and Day 91 for the MenB_0_2_6 Group [2-dose schedule]) compared to Day 1 (Baseline)

Notes:

[42] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immunological noninferiority of the MenABCWY vaccine compared to the rMenB+OMV vaccine.

End point values	MenB_0_2_6 Group	ABCWY_Pooled		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	719	678		
Units: Percentage of participants				
number (confidence interval 95%)				
fHbp (M14459) Ab (N=719,675)	74.7 (71.3 to 77.8)	79.7 (76.5 to 82.7)		
NadA (96217) Ab (N=717,671)	96.4 (94.7 to 97.6)	92.7 (90.5 to 94.5)		
NHBA (M13520) Ab (N=718,678)	58.6 (54.9 to 62.3)	61.9 (58.2 to 65.6)		

PorA (NZ98/254) Ab (N=704,642)	53.3 (49.5 to 57.0)	42.2 (38.4 to 46.1)		
--------------------------------	---------------------	---------------------	--	--

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of blood samples without bactericidal serum activity against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after the last MenACWY dose and 3-dose (0,2,6-M), 2-dose(0,6-M) schedule of rMenB+OMV and 1 dose of MenACWY

End point title	Percentage of blood samples without bactericidal serum activity against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after the last MenACWY dose and 3-dose (0,2,6-M), 2-dose(0,6-M) schedule of rMenB+OMV and 1 dose of MenACWY
-----------------	--

End point description:

The effectiveness of the 3 dose (0,2,6-M) and 2 dose (0,6-M) schedule of rMenB+OMV NZ vaccine and 2 doses of MenACWY vaccine when compared to 1 month after the MenACWY vaccination (Day 31), against a panel of N. meningitidis serogroup B strains is measured in terms of percentage of samples without bactericidal activity using enc-hSBA, which provides a qualitative assessment (yes/no) of the presence of sufficient bactericidal antibodies in human sera to kill a meningococcal strain at a specific dilution of 1:4. Participants were randomly selected for testing against each strain therefore only a subset of participants were tested for each of the strains. Number of Participants analyzed = Total number of participants included in FAS. As pre-specified in the protocol, data reported in this outcome measure were presented for the MenB_0_2_6 group, MenB_0_6 group, ACWY group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 211 for the MenB_0_2_6 group [3 dose schedule], MenB_0_6 group, ABCWY_Pooled group and Day 31 for the MenACWY group)

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	ACWY Group
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	790	813	817	136
Units: Percentage of blood samples				
number (not applicable)				
Meningitis B M10713 Ab (N= 252,247,251,45)	0.4	0.4	1.2	15.6
Meningitis B M08641 Ab (N= 229,259,237,29)	6.6	6.2	8	96.6
Meningitis B M12898 Ab (N= 242,265,249,41)	8.4	11.4	12.1	68.1
Meningitis B M09150 Ab (N= 249,222,273,36)	5.2	8.5	12.6	73
Meningitis B M09401 Ab (N= 243,275,288,49)	50.2	46.5	57	98
Meningitis B M07463 Ab (N= 259,274,256,49)	1.5	1.8	3.5	49
Meningitis B M10496 Ab (N= 263,261,257,47)	54.4	47.9	68.9	100

Meningitis B M14530 Ab (N=241,257,284,37)	2.1	1.6	3.2	100
Meningitis B M15668 Ab (N=237,270,262,49)	0.8	2.2	0.8	87.8
Meningitis B M14028 Ab (N=246,282,283,37)	2.8	5.7	4.9	100
Meningitis B M09909 Ab (N=244,265,261,42)	79.1	83.4	86.2	100
Meningitis B M14385 Ab (N=235,237,259,43)	0.9	0.8	0	16.3
Meningitis B M07992 Ab (N=237,252,274,36)	0.4	0	0.4	0
Meningitis B M09155 Ab (N=234,255,230,45)	1.7	2	1.3	97.8
Meningitis B M13085 Ab (N=256,253,255,44)	13.5	27.5	26.6	69.3
Meningitis B M18303 Ab (N=241,252,264,40)	2.9	4	9.5	100
Meningitis B M18711 Ab (N=252,237,262,33)	5.2	4.2	9.5	75.8
Meningitis B M15009 Ab (N=262,254,242,37)	11.8	20.5	31.8	86.5
Meningitis B M07773 Ab (N=269,262,251,35)	0.7	0.4	0.4	74.3
Meningitis B M09662 Ab (N=267,266,255,48)	61.4	50.8	66.3	95.8
Meningitis B M18483 Ab (N=228,233,222,37)	3	5.1	3.7	72.2
Meningitis B M11906 Ab (N=264,271,260,39)	23.7	30.3	39.9	84.7
Meningitis B M14987 Ab (N=250,274,270,35)	1.2	8.8	12.7	68.6
Meningitis B M12014 Ab (N=249,270,254,46)	0.4	0.7	1.2	67.4
Meningitis B M18200 Ab (N=228,247,242,47)	6.3	14.8	14.9	33.7
Meningitis B M08912 Ab (N=225,247,237,40)	0.4	0	0	0
Meningitis B M16748 Ab (N=262,252,260,45)	0.8	0	0	0
Meningitis B M08152 Ab (N=208,215,223,35)	22.4	25.3	32.8	66.8
Meningitis B M09973 Ab (N=252,270,247,48)	0.8	1.1	3	83.3
Meningitis B M15352 Ab (N=237,249,245,33)	8.9	8.4	17.8	97
Meningitis B M15165 Ab (N=249,250,240,42)	0	1.2	2.1	92.9
Meningitis B M08127 Ab (N=269,250,287,51)	0.7	0.4	0.7	84.3
Meningitis B M18347 Ab (N=234,262,239,44)	45.4	50.9	51.8	82.1
Meningitis B M12500 Ab (N=223,253,241,43)	0.9	2.4	2.5	95.3
Meningitis B M07499 Ab (N=242,251,256,51)	70.7	75.7	80.1	100
Meningitis B M09960 Ab (N=243,247,251,33)	1.2	0	0	3
Meningitis B M18045 Ab (N=236,251,272,41)	0	0.4	0	92.7
Meningitis B M10548 Ab (N=234,243,261,58)	8.1	11.9	12.5	74.1

Meningitis B M09354 Ab (N=246,240,241,40)	1.2	1.3	0	80
Meningitis B M11051 Ab (N=246,282,247,46)	61	64.2	66.7	97.8
Meningitis B M10104 Ab (N=247,241,239,41)	58.7	52.3	63.5	97.6
Meningitis B M13361 Ab (N=258,268,250,34)	0.8	0.4	0.4	85.3
Meningitis B M11042 Ab (N=217,249,254,43)	19.8	25.5	33.5	85.7
Meningitis B M18467 Ab (N=255,223,250,47)	1.2	0.4	0.8	78.7
Meningitis B M11113 Ab (N=255,253,269,51)	30.1	39.5	49.5	75.2
Meningitis B M07253 Ab (N=260,245,255,40)	34.7	33.8	50.2	86.4
Meningitis B M07356 Ab (N=235,235,266,29)	0.4	0	1.1	41.4
Meningitis B M10710 Ab (N=254,253,270,40)	1.6	2	1.4	92.5
Meningitis B M17147 Ab (N=261,279,214,41)	2.3	5.4	2.3	100
Meningitis B M14401 Ab (N=241,266,253,43)	1.7	0.4	1.2	83.7
Meningitis B M14293 Ab (N=251,247,224,47)	45.8	25.1	45.2	95.7
Meningitis B M08540 Ab (N=258,261,222,34)	1.6	0.8	0	38.2
Meningitis B M07960 Ab (N=277,241,241,39)	3.6	4.1	1.8	94.9
Meningitis B M16135 Ab (N=254,230,257,41)	0	1.7	0.8	95.1
Meningitis B M14548 Ab (N=267,266,274,38)	2.6	3.4	3.2	94.7
Meningitis B M09181 Ab (N=228,276,240,43)	0	0.4	0.4	72.1
Meningitis B M14224 Ab (N=248,279,277,40)	0.4	0.4	0	82.5
Meningitis B M07452 Ab (N=260,260,245,47)	2.7	8.1	17.1	85.1
Meningitis B M13520 Ab (N=217,222,242,30)	3.2	0.9	0	66.7
Meningitis B M09385 Ab (N=228,252,265,32)	0.4	1.6	0	46.9
Meningitis B M14881 Ab (N=240,278,277,40)	4.2	5.8	10	95
Meningitis B M13252 Ab (N=279,241,262,50)	0.7	1.2	0.4	98
Meningitis B M07818 Ab (N=232,244,228,43)	0.4	0.8	1.3	90.7
Meningitis B M09914 Ab (N=281,257,258,49)	85.4	86.8	88.3	98
Meningitis B M15083 Ab (N=231,262,241,40)	51.4	56.7	61	84.5
Meningitis B M11290 Ab (N=264,266,221,34)	61.4	61.7	67.4	100
Meningitis B M14988 Ab (N=241,269,246,40)	0.4	0	1.2	60
Meningitis B M10536 Ab (N=228,223,260,36)	19.7	14.3	14.3	91.7
Meningitis B M08959 Ab (N=252,241,258,47)	0.8	0.4	1.9	85.1

Meningitis B M08785 Ab (N=251,249,260,39)	0.8	0.4	0.4	53.8
Meningitis B M07245 Ab (N=238,248,230,43)	0	0	0.9	23.3
Meningitis B M19315 Ab (N=262,254,255,34)	3.8	3.1	9.5	79.4
Meningitis B M14376 Ab (N=234,220,222,41)	0	1.4	0	92.7
Meningitis B M08994 Ab (N=226,253,233,31)	2.5	7.6	6.9	62.7
Meningitis B M11646 Ab (N=241,224,250,36)	0	1.3	2	83.3
Meningitis B M13362 Ab (N=259,228,231,49)	0	0.4	0.4	81.6
Meningitis B M08080 Ab (N=236,247,242,45)	27.4	41	51.6	85.7
Meningitis B M08370 Ab (N=265,262,251,43)	2.3	1.5	3.7	97.7
Meningitis B M08129 Ab (N=245,254,257,28)	4.1	4.7	7	71.4
Meningitis B M07111 Ab (N=231,254,270,33)	0.4	1.2	0.7	90.9
Meningitis B M07537 Ab (N=241,242,263,47)	95.9	95.9	97.7	100
Meningitis B M13438 Ab (N=255,262,282,50)	1.2	0.8	0	16
Meningitis B M10661 Ab (N=246,275,273,33)	2	2.9	2.9	97
Meningitis B M10920 Ab (N=247,252,275,34)	29.1	27.8	37.8	91.2
Meningitis B M15564 Ab (N=240,273,263,40)	0.4	0.7	0.8	77.5
Meningitis B M10934 Ab (N=273,241,256,36)	0.4	0.8	0.8	100
Meningitis B M09400 Ab (N=254,281,279,39)	0.8	1.8	0.4	97.4
Meningitis B M08781 Ab (N=221,258,229,28)	71.9	74.4	78.2	100
Meningitis B M09173 Ab (N=238,249,247,42)	0.4	0.4	0.8	95.2
Meningitis B M14113 Ab (N=228,255,264,42)	15.4	21.2	22	100
Meningitis B M08389 Ab (N=254,259,250,45)	10.7	7	11.7	87.2
Meningitis B M16822 Ab (N=239,233,264,34)	67.8	76.4	82.2	100
Meningitis B M10995 Ab (N=268,258,239,47)	5.2	17.8	30.6	85.1
Meningitis B M08780 Ab (N=236,259,260,40)	1.3	0.8	1.5	92.5
Meningitis B M09910 Ab (N=247,246,257,43)	1.6	1.2	1.2	93
Meningitis B M08320 Ab (N=262,238,253,45)	33.6	39.7	43.3	87
Meningitis B M14879 Ab (N=239,242,247,47)	2.1	2.1	6.1	21.3
Meningitis B M09345 Ab (N=251,257,282,48)	19.3	20.1	33.4	81.2
Meningitis B M14594 Ab (N=245,241,238,44)	20.8	27.4	41.2	97.7
Meningitis B M07621 Ab (N=251,260,263,40)	1.2	1.2	1.1	77.5

Meningitis B M13568 Ab (N=261,264,253,40)	5.4	3.8	3.9	95
Meningitis B M18017 Ab (N=246,247,258,31)	0.4	0	1.6	96.8
Meningitis B M08420 Ab (N=243,255,256,40)	0.8	0.4	1.6	95
Meningitis B M07959 Ab (N=236,237,254,35)	1.7	2.5	1.7	97.1
Meningitis B M06970 Ab (N=236,261,253,32)	19.6	17.6	30.2	85.7
Meningitis B M10491 Ab (N=239,262,227,39)	5.4	8.4	17	82.1
Meningitis B M13569 Ab (N=221,243,247,31)	0.9	2.9	1.2	96.8
Meningitis B M10182 Ab (N=229,234,256,44)	0.4	0	0	0
Meningitis B M13547 Ab (N=251,258,238,44)	2.4	7	4.7	47.7
Meningitis B M15276 Ab (N=231,231,259,41)	0.4	0	1.5	87.8

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of blood samples without bactericidal serum activity using enc-hSBA against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after the 2-dose(0,6-M) schedule of rMenB+OMV and 1 dose of MenACWY

End point title	Percentage of blood samples without bactericidal serum activity using enc-hSBA against each of the endemic U.S N. meningitidis serogroup B strains at 1 month after the 2-dose(0,6-M) schedule of rMenB+OMV and 1 dose of
-----------------	---

End point description:

The effectiveness of the 2 dose (0,2-M) schedule of rMenB+OMV NZ vaccine when compared to 1 month after the MenACWY vaccination (Day 31), against a panel of N. meningitidis serogroup B strains is measured in terms of percentage of samples without bactericidal activity using enc-hSBA, which provides a qualitative assessment (yes/no) of the presence of sufficient bactericidal antibodies in human sera to kill a meningococcal strain at a specific dilution of 1:4.

Participants were randomly selected for testing against each strain therefore only a subset of participants were tested for each of the strains. Number of Participants analyzed = Total number of participants included in FAS.

Analysis was performed on blood samples collected from FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 91 for the MenB_0_2_6 group [2 dose schedule] and Day 31 for the MenACWY group)

Notes:

[43] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the effectiveness of the rMenB+OMV vaccine compared to one dose of MenACWY vaccination in the ACWY group.

End point values	MenB_0_2_6 Group	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	893	173		
Units: Percentage of blood samples				
number (not applicable)				
Meningitis B M10713 Ab (N=266,45)	1.9	15.6		
Meningitis B M08641 Ab (N=250,29)	11.2	96.6		
Meningitis B M12898 Ab (N=260,41)	12.4	69		
Meningitis B M09150 Ab (N=250,36)	9.3	68.6		
Meningitis B M09401 Ab (N=253,49)	47	98		
Meningitis B M07463 Ab (N=265,49)	2.6	49		
Meningitis B M10496 Ab (N=277,47)	50.5	100		
Meningitis B M14530 Ab (N=254,37)	3.9	100		
Meningitis B M15668 Ab (N=252,49)	0.4	87.8		
Meningitis B M14028 Ab (N=257,37)	5.4	100		
Meningitis B M09909 Ab (N=266,42)	82	100		
Meningitis B M14385 Ab (N=227,43)	0	16.3		
Meningitis B M07992 Ab (N=249,36)	0	0		
Meningitis B M09155 Ab (N=243,45)	2.5	97.8		
Meningitis B M13085 Ab (N=271,44)	21	66.4		
Meningitis B M18303 Ab (N=252,40)	9.1	100		
Meningitis B M18711 Ab (N=261,33)	7.7	75.8		
Meningitis B M15009 Ab (N=243,37)	22.2	86.5		
Meningitis B M07773 Ab (N=281,35)	1.1	74.3		
Meningitis B M09662 Ab (N=284,48)	58.1	95.8		
Meningitis B M18483 Ab (N=234,37)	4.3	77.3		
Meningitis B M11906 Ab (N=277,39)	38.8	85.3		
Meningitis B M14987 Ab (N=264,35)	10.7	62.3		
Meningitis B M12014 Ab (N=267,46)	0.7	63.2		
Meningitis B M18200 Ab (N=241,47)	12.1	34.4		
Meningitis B M08912 Ab (N=246,40)	0	0		
Meningitis B M16748 Ab (N=281,45)	0	0		
Meningitis B M08152 Ab (N=224,35)	35.6	67.2		
Meningitis B M09973 Ab (N=271,48)	1.5	83.3		
Meningitis B M15352 Ab (N=219,33)	11.9	97		
Meningitis B M15165 Ab (N=265,42)	0.8	92.9		
Meningitis B M08127 Ab (N=285,51)	0.4	84.3		
Meningitis B M18347 Ab (N=249,44)	60.5	80.4		
Meningitis B M12500 Ab (N=236,43)	3.4	95.3		
Meningitis B M07499 Ab (N=259,51)	81.1	100		
Meningitis B M09960 Ab (N=251,33)	4.4	3		
Meningitis B M18045 Ab (N=247,41)	0	92.7		
Meningitis B M10548 Ab (N=244,58)	11.9	74.1		
Meningitis B M09354 Ab (N=258,40)	0.4	80		
Meningitis B M11051 Ab (N=270,46)	68.9	97.8		
Meningitis B M10104 Ab (N=249,41)	61.8	97.6		
Meningitis B M13361 Ab (N=269,34)	0.4	85.3		
Meningitis B M11042 Ab (N=210,43)	30.1	84.2		
Meningitis B M18467 Ab (N=272,47)	0.4	78.7		
Meningitis B M11113 Ab (N=273,51)	43.1	75.6		
Meningitis B M07253 Ab (N=274,40)	43.3	85.3		

Meningitis B M07356 Ab (N=247,29)	0	41.4		
Meningitis B M10710 Ab (N=269,40)	2.2	92.5		
Meningitis B M17147 Ab (N=274,41)	5.1	100		
Meningitis B M14401 Ab (N=251,43)	0.8	83.7		
Meningitis B M14293 Ab (N=264,47)	36.4	95.7		
Meningitis B M08540 Ab (N=271,34)	0.4	38.2		
Meningitis B M07960 Ab (N=292,39)	4.1	94.9		
Meningitis B M16135 Ab (N=263,41)	0.8	95.1		
Meningitis B M14548 Ab (N=286,38)	2.8	94.7		
Meningitis B M09181 Ab (N=252,43)	0	72.1		
Meningitis B M14224 Ab (N=268,40)	0.7	82.5		
Meningitis B M07452 Ab (N=267,47)	13.5	85.1		
Meningitis B M13520 Ab (N=234,30)	0.9	66.7		
Meningitis B M09385 Ab (N=230,32)	2.2	46.9		
Meningitis B M14881 Ab (N=258,40)	14.7	95		
Meningitis B M13252 Ab (N=286,50)	3.1	98		
Meningitis B M07818 Ab (N=249,43)	0.4	90.7		
Meningitis B M09914 Ab (N=293,49)	87.7	98		
Meningitis B M15083 Ab (N=260,40)	63.7	78		
Meningitis B M11290 Ab (N=278,34)	70.5	100		
Meningitis B M14988 Ab (N=251,40)	0.4	60		
Meningitis B M10536 Ab (N=241,36)	23.2	91.7		
Meningitis B M08959 Ab (N=267,47)	0.4	85.1		
Meningitis B M08785 Ab (N=262,39)	0.4	53.8		
Meningitis B M07245 Ab (N=254,43)	0	23.3		
Meningitis B M19315 Ab (N=270,34)	6.3	79.4		
Meningitis B M14376 Ab (N=269,41)	0	92.7		
Meningitis B M08994 Ab (N=233,31)	8.9	55.7		
Meningitis B M11646 Ab (N=252,36)	1.6	83.3		
Meningitis B M13362 Ab (N=272,49)	1.5	81.6		
Meningitis B M08080 Ab (N=244,45)	46.8	87.1		
Meningitis B M08370 Ab (N=280,43)	3.9	97.7		
Meningitis B M08129 Ab (N=242,28)	7.4	71.4		
Meningitis B M07111 Ab (N=244,33)	0.4	90.9		
Meningitis B M07537 Ab (N=253,47)	96.4	100		
Meningitis B M13438 Ab (N=263,50)	0.4	16		
Meningitis B M10661 Ab (N=259,33)	2.7	97		
Meningitis B M10920 Ab (N=259,34)	50.2	91.2		
Meningitis B M15564 Ab (N=258,40)	0.8	77.5		
Meningitis B M10934 Ab (N=286,36)	0	100		
Meningitis B M09400 Ab (N=261,39)	1.1	97.4		
Meningitis B M08781 Ab (N=223,28)	75.8	100		
Meningitis B M09173 Ab (N=244,42)	0	95.2		
Meningitis B M14113 Ab (N=244,42)	28.3	100		
Meningitis B M08389 Ab (N=258,45)	1.9	86.7		
Meningitis B M16822 Ab (N=240,34)	85	100		
Meningitis B M10995 Ab (N=282,47)	16.3	85.1		
Meningitis B M08780 Ab (N=241,40)	0.8	92.5		
Meningitis B M09910 Ab (N=256,43)	1.2	93		
Meningitis B M08320 Ab (N=277,45)	45.1	86.8		
Meningitis B M14879 Ab (N=252,47)	3.3	23.2		
Meningitis B M09345 Ab (N=263,48)	32.3	74.9		

Meningitis B M14594 Ab (N=254,44)	35.8	97.7		
Meningitis B M07621 Ab (N=259,40)	0.4	77.5		
Meningitis B M13568 Ab (N=268,40)	9.7	95		
Meningitis B M18017 Ab (N=262,31)	0	96.8		
Meningitis B M08420 Ab (N=260,40)	0.8	95		
Meningitis B M07959 Ab (N=243,35)	1.2	97.1		
Meningitis B M06970 Ab (N=243,32)	28.8	90.6		
Meningitis B M10491 Ab (N=254,39)	16.1	82.1		
Meningitis B M13569 Ab (N=233,31)	0.4	96.8		
Meningitis B M10182 Ab (N=251,44)	0	0		
Meningitis B M13547 Ab (N=260,44)	5	47.7		
Meningitis B M15276 Ab (N=238,41)	0.4	87.8		

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants classified by percentage of serogroup B invasive disease strains killed using enc-hSBA in each subject at 1 month after vaccination schedule for the MenB_0_2_6 group [3 dose], MenB_0_6 group and last MenABCWY dose (pooled lots)

End point title	Percentage of participants classified by percentage of serogroup B invasive disease strains killed using enc-hSBA in each subject at 1 month after vaccination schedule for the MenB_0_2_6 group [3 dose], MenB_0_6 group and last MenABCWY dose (pooled lots) ^[44]
-----------------	--

End point description:

The percentage of participants classified by percentage of N.meningitidis serogroup B invasive strains killed using enc-hSBA and the corresponding 2- sided 95% CI based on Clopper-Pearson method is calculated for each vaccine group. As pre-specified in the protocol, data reported in this outcome measure were presented for the MenB_0_2_6 group, MenB_0_6 group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (Day 211 for MenB_0_2_6 group (3 dose schedule), MenB_0_6 group and ABCWY_Pooled group)

Notes:

[44] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the percentage of serogroup B invasive disease strains killed within a subject using enc-hSBA of the rMenB+OMV and MenABCWY vaccines.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	790	813	817	
Units: Percentage of participants				
number (confidence interval 95%)				
>=50% killed strains	98.7 (97.7 to 99.4)	98.5 (97.4 to 99.2)	98 (96.8 to 98.9)	
>=55% killed strains	98.4 (97.2 to 99.1)	97.4 (96.1 to 98.4)	96.8 (95.4 to 97.9)	

>=60% killed strains	97.8 (96.6 to 98.7)	96.8 (95.3 to 97.9)	95.2 (93.5 to 96.6)	
>=65% killed strains	96.5 (94.9 to 97.6)	93.6 (91.7 to 95.2)	90.1 (87.8 to 92)	
>=70% killed strains	93.4 (91.5 to 95)	89.8 (87.5 to 91.8)	84.1 (81.4 to 86.5)	
>=75% killed strains	86.8 (84.3 to 89.1)	82.2 (79.4 to 84.7)	74.7 (71.5 to 77.6)	
>=80% killed strains	79.2 (76.2 to 82)	75.5 (72.4 to 78.4)	66 (62.6 to 69.2)	
>=85% killed strains	62.8 (59.3 to 66.2)	60.4 (56.9 to 63.8)	50.1 (46.6 to 53.5)	
>=90% killed strains	43.7 (40.2 to 47.2)	41.3 (37.9 to 44.8)	32.1 (28.9 to 35.4)	
>=95% killed strains	22.5 (19.7 to 25.6)	21 (18.3 to 24)	13.7 (11.4 to 16.3)	
100% killed strains	10 (8 to 12.3)	8.4 (6.6 to 10.5)	6.1 (4.6 to 8)	

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants classified by percentage of serogroup B invasive disease strains killed using enc-hSBA in each subject at 1 month after vaccination with rMenB+OMV NZ (0,2-months)

End point title	Percentage of participants classified by percentage of serogroup B invasive disease strains killed using enc-hSBA in each subject at 1 month after vaccination with rMenB+OMV NZ (0,2-months) ^[45]
-----------------	---

End point description:

The percentage of participants are classified by percentage of N.meningitidis serogroup B invasive strains killed using enc-hSBA and the corresponding 2- sided 95% CI based on Clopper-Pearson method is calculated for each vaccine group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (Day 91 for MenB_0_2_6 group [2 dose schedule])

Notes:

[45] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the percentage of serogroup B invasive disease strains killed within a subject using enc-hSBA of the rMenB+OMV and MenABCWY vaccines.

End point values	MenB_0_2_6 Group			
Subject group type	Reporting group			
Number of subjects analysed	831			
Units: Percentage of participants				
number (confidence interval 95%)				
>=50% killed strains	98.6 (97.5 to 99.3)			
>=55% killed strains	97.7 (96.5 to 98.6)			
>=60% killed strains	96.5 (95 to 97.7)			

>=65% killed strains	92.2 (90.1 to 93.9)			
>=70% killed strains	84.8 (82.2 to 87.2)			
>=75% killed strains	75.7 (72.6 to 78.6)			
>=80% killed strains	66.7 (63.3 to 69.9)			
>=85% killed strains	49.7 (46.2 to 53.2)			
>=90% killed strains	33.8 (30.6 to 37.1)			
>=95% killed strains	16.2 (13.8 to 18.9)			
100% killed strains	7.7 (6 to 9.7)			

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants with hSBA titers >= LLOQ for each and all serogroup B indicator strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months and 0,6-months) and last dose of MenABCWY (ABCWY group-pooled lots)

End point title	Percentage of participants with hSBA titers >= LLOQ for each and all serogroup B indicator strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months and 0,6-months) and last dose of MenABCWY (ABCWY group-pooled lots) ^[46]
-----------------	--

End point description:

The immune response to rMenB+OMV NZ and MenABCWY vaccine is evaluated by measuring bactericidal activity against each (individual response) and all (composite response) N. meningitidis serogroup B indicator strains (M14459, 96217, M13520 and NZ98/254 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively). As pre-specified in the protocol, data reported in this outcome measure were presented for the MenB_0_2_6 group, MenB_0_6 group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1 (pre-vaccination) and 1 month after the vaccination schedule (i.e., Day 211 for MenB_0_2_6 group [3 dose schedule], MenB_0_6 group and ABCWY_Pooled group)

Notes:

[46] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV and MenABCWY vaccines against N. meningitidis serogroup B indicator strains.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	749	731	780	
Units: Percentage of participants				
number (confidence interval 95%)				
fHbp (M14459) Ab, Day 1 (N=749,730,762)	4.9 (3.5 to 6.7)	3.4 (2.2 to 5)	5.4 (3.9 to 7.2)	

fHbp (M14459) Ab, Day 211 (N=690,707,738)	97.4 (95.9 to 98.4)	94.6 (92.7 to 96.2)	95.9 (94.2 to 97.2)	
NadA (96217) Ab, Day 1 (N=744,731,780)	6.2 (4.6 to 8.2)	4.4 (3.0 to 6.1)	6.2 (4.6 to 8.1)	
NadA (96217) Ab, Day 211 (N=691,707,734)	100 (99.5 to 100)	98 (96.7 to 98.9)	96.2 (94.5 to 97.5)	
NHBA (M13520) Ab, Day 1 (N=749,731,764)	23.2 (20.3 to 26.4)	20.9 (18 to 24.1)	18.5 (15.8 to 21.4)	
NHBA (M13520) Ab, Day 211 (N=695,711,738)	97.0 (95.4 to 98.1)	97.5 (96.0 to 98.5)	95.3 (93.5 to 96.7)	
PorA (NZ98/254) Ab, Day 1 (N=738,716,751)	2.3 (1.3 to 3.7)	1.4 (0.7 to 2.6)	2.1 (1.2 to 3.4)	
PorA (NZ98/254) Ab, Day 211 (N=657,684,709)	85.8 (82.9 to 88.4)	82.6 (79.5 to 85.4)	75.3 (72.0 to 78.5)	
Composite Response, Day=1 (N=727,708,747)	1.1 (0.5 to 2.2)	0.6 (0.2 to 1.4)	1.1 (0.5 to 2.1)	
Composite Response, Day=211 (N=654,683,707)	83.3 (80.3 to 86.1)	80.7 (77.5 to 83.6)	71.4 (67.9 to 74.7)	

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants with hSBA titers \geq LLOQ for each and all serogroup B indicator strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2-months)

End point title	Percentage of participants with hSBA titers \geq LLOQ for each and all serogroup B indicator strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2-months) ^[47]
-----------------	---

End point description:

The immune response to rMenB+OMV NZ is evaluated by measuring bactericidal activity against each (individual response) and all (composite response) N. meningitidis serogroup B indicator strains (M14459, 96217, M13520 and NZ98/254 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively). Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1 (pre-vaccination) and 1 month after the vaccination schedule (i.e., Day 91 for MenB_0_2_6 group [2 dose schedule])

Notes:

[47] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV vaccines against N. meningitidis serogroup B indicator strains.

End point values	MenB_0_2_6 Group			
Subject group type	Reporting group			
Number of subjects analysed	753			
Units: Percentage of participants				
number (confidence interval 95%)				
fHbp (M14459) Ab, Day 1 (N=749)	4.9 (3.5 to 6.7)			
fHbp (M14459) Ab, Day 91 (N=750)	92.9 (90.9 to 94.7)			
NadA (96217) Ab, Day 1 (N=744)	6.2 (4.6 to 8.2)			

NadA (96217) Ab, Day 91 (N=753)	99.5 (98.6 to 99.9)			
NHBA (M13520) Ab, Day 1 (N=749)	23.2 (20.3 to 26.4)			
NHBA (M13520) Ab, Day 91 (N=750)	96.1 (94.5 to 97.4)			
PorA N (NZ98/254) Ab, Day 1 (N=738)	2.3 (1.3 to 3.7)			
PorA N (NZ98/254) Ab, Day 91 (N=745)	80 (76.9 to 82.8)			
Composite Response, Day=1 (N=727)	1.1 (0.5 to 2.2)			
Composite Response, Day=91 (N=744)	75.5 (72.3 to 78.6)			

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants with 4-fold rise in hSBA titers for each of the serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2 months)

End point title	Percentage of participants with 4-fold rise in hSBA titers for each of the serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2 months) ^[48]
-----------------	--

End point description:

The immune response to 2 dose (0,2-M) is evaluated by measuring bactericidal activity against each of the N. meningitidis serogroup B test strains- M14459, 96217, NZ98/254 and M13520 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively compared to baseline. Four-fold rise per each indicator strain was defined as a post-vaccination hSBA titre ≥ 16 for participants with a pre-vaccination hSBA titre < 4 a post-vaccination hSBA titre ≥ 4 times the LLOQ for subjects with a pre-vaccination hSBA titre $\geq \text{LOD}$ and $< \text{LLOQ}$ a post-vaccination hSBA titre ≥ 4 times the pre-vaccination hSBA titre for subjects with a pre-vaccination hSBA titre $\geq \text{LLOQ}$. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 91 for MenB_0_2_6 [2-dose schedule]) compared to Day 1 (baseline)

Notes:

[48] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV vaccines against N. meningitidis serogroup B indicator strains.

End point values	MenB_0_2_6 Group			
Subject group type	Reporting group			
Number of subjects analysed	739			
Units: Percentage of participants				
number (confidence interval 95%)				
fHbp (M14459) Ab (N=739)	74.6 (71.3 to 77.7)			
NadA (96217) Ab (N=738)	96.3 (94.7 to 97.6)			
NHBA (M13520) Ab (N=739)	58.5 (54.8 to 62.0)			
PorA (NZ98/254) Ab (N=724)	53.5 (49.7 to 57.1)			

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants with 4-fold rise in hSBA titers for each of the serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months and 0,6-months) and last dose of MenABCWY (ABCWY group-pooled lots)

End point title	Percentage of participants with 4-fold rise in hSBA titers for each of the serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months and 0,6-months) and last dose of MenABCWY (ABCWY group-pooled lots) ^[49]
-----------------	---

End point description:

The immune response to 3 dose (0,2,6-M), 2 dose (0,6-M) schedule of rMenB+OMV NZ and 2 doses of MenABCWY vaccine was evaluated by measuring bactericidal activity against each of the N. meningitidis serogroup B test strains- M14459, 96217, NZ98/254 and M13520 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively compared to baseline. Four-fold rise per each indicator strain was defined as a post-vaccination hSBA titre ≥ 16 for subjects with a pre-vaccination hSBA titre < 4 a post-vaccination hSBA titer ≥ 4 times the LLOQ for subjects with a pre-vaccination hSBA titre $\geq \text{LOD}$ and $< \text{LLOQ}$ a post-vaccination hSBA titre ≥ 4 times the pre-vaccination hSBA titre for participants with a pre-vaccination hSBA titre $\geq \text{LLOQ}$. As pre-specified in the protocol, data reported in this outcome measure were presented for the MenB_0_2_6 group, MenB_0_6 group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 211 for MenB_0_2_6 group [3 dose schedule], MenB_0_6 group and ABCWY_Pooled group) compared to Day 1 (baseline)

Notes:

[49] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV and MenABCWY vaccines against N. meningitidis serogroup B indicator strains.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	685	704	731 ^[50]	
Units: Percentage of participants				
number (confidence interval 95%)				
fHbp (M14459) Ab (N=679,699, 729)	86.7 (84.0 to 89.2)	82.4 (79.4 to 85.2)	78.9 (75.7 to 81.8)	
NadA (96217) Ab (N=679,700,725)	98.7 (97.5 to 99.4)	95.3 (93.4 to 96.7)	92.3 (90.1 to 94.1)	
NHBA (M13520) Ab (N=685,704,731)	66.9 (63.2 to 70.4)	69.5 (65.9 to 72.8)	61.1 (57.5 to 64.7)	
PorA (NZ98/254) Ab (N=637,664,693)	56.5 (52.6 to 60.4)	57.2 (53.4 to 61.0)	42.4 (38.7 to 46.2)	

Notes:

[50] - Analysis of final results for the ABCWY group is ongoing and will be updated subsequently.

Statistical analyses

No statistical analyses for this end point

Secondary: hSBA Geometric Mean Titres (GMTs) against each of the N. meningitidis serogroup B strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months, 0,6-months) and last dose of MenABCWY (ABCWY group-pooled lots)

End point title	hSBA Geometric Mean Titres (GMTs) against each of the N. meningitidis serogroup B strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months, 0,6-months) and last dose of MenABCWY (ABCWY group-pooled lots) ^[51]
-----------------	--

End point description:

The immune response to rMenB+OMV NZ and MenABCWY vaccine was evaluated by measuring bactericidal activity against N. meningitidis serogroup B test strains in terms of GMTs after vaccination compared to baseline (Day 1). For each N. meningitidis serogroup B test strain (M14459, M13520, 96217 and NZ98/254 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively), The GMTs (After vaccination/baseline) are calculated, with their associated 2-sided 95% CIs. As pre-specified in the protocol, data reported in this outcome measure were presented for the MenB_0_2_6 group, MenB_0_6 group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1 (pre-vaccination) and at 1 month after the vaccination schedule (i.e., Day 211 for MenB_0_2_6 group (3 dose schedule), MenB_0_6 group and ABCWY_Pooled group)

Notes:

[51] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV and MenABCWY vaccines against N. meningitidis serogroup B indicator strains

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	749	731	780	
Units: Titers				
geometric mean (confidence interval 95%)				
fHbp (M14459) Ab, Day 1 (N=749,730,762)	2.8 (2.7 to 2.8)	2.7 (2.6 to 2.8)	2.8 (2.7 to 2.9)	
fHbp (M14459) Ab, Day 211 (N=690,707,738)	30.8 (28.3 to 33.5)	28.1 (25.9 to 30.6)	25 (23 to 27.1)	
NadA (96217) Ab, Day 1 (N=744,731,780)	8.4 (8.1 to 8.6)	8.3 (8 to 8.6)	8.5 (8.2 to 8.8)	
NadA (96217) Ab, Day 211 (N=691,707,734)	267.2 (243.7 to 293)	215.1 (196.2 to 235.9)	150.6 (137.3 to 165.2)	
NHBA (M13520) Ab, Day 1 (N=749,731,764)	3.4 (3.1 to 3.7)	3.2 (3 to 3.5)	3.1 (2.8 to 3.4)	
NHBA (M13520) Ab, Day 211 (N=695,711,738)	30.6 (27.7 to 33.7)	33.2 (30.2 to 36.6)	25.2 (22.9 to 27.8)	
PorA (NZ98/254) Ab, Day 1 (N=738,716,751)	3.2 (3.1 to 3.2)	3.1 (3 to 3.2)	3.1 (3.1 to 3.2)	
PorA (NZ98/254) Ab, Day 211 (N=657,684,709)	18.1 (16.3 to 20.1)	17.7 (15.9 to 19.6)	12.9 (11.6 to 14.4)	

Statistical analyses

Secondary: hSBA GMTs against each of the N. meningitidis serogroup B strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2-months)

End point title	hSBA GMTs against each of the N. meningitidis serogroup B strains at Day 1 and at 1 month after vaccination with rMenB+OMV NZ (0,2-months) ^[52]
-----------------	--

End point description:

The immune response to rMenB+OMV NZ and MenABCWY vaccine was evaluated by measuring bactericidal activity against N. meningitidis serogroup B test strains in terms of GMTs after vaccination compared to baseline (Day 1). For each N. meningitidis serogroup B test strain (M14459, M13520, 96217 and NZ98/254 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively), The GMTs (After vaccination/baseline) were calculated, with their associated 2-sided 95% CIs. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1 (pre-vaccination) and at 1 month after the vaccination schedule (i.e., Day 91 for MenB_0_2_6 group [2-dose schedule])

Notes:

[52] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV vaccines against N. meningitidis serogroup B indicator strains.

End point values	MenB_0_2_6 Group			
Subject group type	Reporting group			
Number of subjects analysed	753			
Units: Titers				
geometric mean (confidence interval 95%)				
fHbp (M14459) Ab, Day 1(N=749)	2.8 (2.7 to 2.8)			
fHbp (M14459) Ab, Day 91(N=750)	20.9 (18.9 to 23.1)			
NadA (96217) Ab, Day 1(N=744)	8.4 (8.1 to 8.6)			
NadA (96217) Ab, Day 91(N=753)	178.5 (161.7 to 197.2)			
NHBA (M13520) Ab, Day 1(N=749)	3.4 (3.1 to 3.7)			
NHBA (M13520) Ab, Day 91(N=750)	27.2 (24.1 to 30.6)			
PorA (NZ98/254) Ab, Day 1(N=738)	3.2 (3.1 to 3.2)			
PorA (NZ98/254) Ab, Day 91(N=745)	17.1 (15.2 to 19.3)			

Statistical analyses

No statistical analyses for this end point

Secondary: hSBA Geometric Mean Ratios (GMRs) for each of the N. meningitidis serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months, 0,6-months) and last dose of MenABCWY (ABCWY group-pooled lots)

End point title	hSBA Geometric Mean Ratios (GMRs) for each of the N. meningitidis serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2,6-months, 0,6-months) and last
-----------------	--

End point description:

The immune response to rMenB+OMV NZ vaccine was evaluated by measuring bactericidal activity against *N. meningitidis* serogroup B test strains after vaccination compared to baseline (Day 1). For each *N. meningitidis* serogroup B test strain (M14459, M13520, 96217 and NZ98/254 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively), the GMRs (after vaccination/baseline) are calculated, with their associated 2-sided 95% CIs. As pre-specified in the protocol, data reported in this outcome measure were presented for the MenB_0_2_6 group, MenB_0_6 group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 211 for MenB_0_2_6 group (3 dose schedule), MenB_0_6 group and ABCWY_Pooled group) compared to Day 1 (baseline)

Notes:

[53] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV and MenABCWY vaccines against *N. meningitidis* serogroup B indicator strains.

End point values	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY_Pooled	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	685	704	731	
Units: Ratio				
geometric mean (confidence interval 95%)				
fHbp (M14459) Ab (N=679,699,729)	11.2 (10.3 to 12.2)	10.5 (9.6 to 11.4)	9 (8.2 to 9.8)	
NadA (96217) Ab (N=679,700,725)	32.1 (29.1 to 35.3)	25.8 (23.5 to 28.4)	17.7 (16 to 19.4)	
NHBA (M13520) Ab (N=685,704,731)	9.1 (8.2 to 10.1)	10.6 (9.5 to 11.7)	8.2 (7.4 to 9.1)	
PorA (NZ98/254) Ab (N=637,664,693)	5.8 (5.2 to 6.5)	5.8 (5.2 to 6.4)	4.1 (3.7 to 4.6)	

Statistical analyses

No statistical analyses for this end point

Secondary: hSBA GMRs for each of the *N. meningitidis* serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2-months)

End point title	hSBA GMRs for each of the <i>N. meningitidis</i> serogroup B strains at 1 month after vaccination with rMenB+OMV NZ (0,2-months) ^[54]
-----------------	--

End point description:

The immune response to rMenB+OMV NZ vaccine was evaluated by measuring bactericidal activity against *N. meningitidis* serogroup B test strains after vaccination compared to baseline (Day 1). For each *N. meningitidis* serogroup B test strain (M14459, M13520, 96217 and NZ98/254 for fHbp, NadA, NHBA and PorA P1.4 antigens, respectively), the GMRs (after vaccination/baseline) are calculated, with their associated 2-sided 95% CIs. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after the vaccination schedule (i.e., Day 91 for MenB_0_2_6 group [2-dose schedule]) compared to Day 1 (baseline)

Notes:

[54] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the rMenB+OMV vaccines against N. meningitidis serogroup B indicator strains.

End point values	MenB_0_2_6 Group			
Subject group type	Reporting group			
Number of subjects analysed	739			
Units: Ratio				
geometric mean (confidence interval 95%)				
fHbp (M14459) Ab (N=739)	7.7 (6.9 to 8.5)			
NadA (96217) Ab (N=738)	21.7 (19.5 to 24)			
NHBA (M13520) Ab (N=739)	8 (7.1 to 9)			
PorA (NZ98/254) Ab (N=724)	5.5 (4.9 to 6.2)			

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants with hSBA titers \geq LLOQ for each of the N. meningitidis serogroups A,C,W,Y at Day 1 and at 1 month after the first and the last MenABCWY vaccination for ABCWY_Pooled group and 1 month after the MenACWY vaccine for ACWY group

End point title	Percentage of participants with hSBA titers \geq LLOQ for each of the N. meningitidis serogroups A,C,W,Y at Day 1 and at 1 month after the first and the last MenABCWY vaccination for ABCWY_Pooled group and 1 month after the MenACWY vaccine for ACWY group ^[55]
-----------------	--

End point description:

The immune responses to MenABCWY and MenACWY vaccines were evaluated by measuring bactericidal activity against N. meningitidis serogroups A, C, W and Y after vaccination compared to baseline (Day 1) and expressed as the percentage of participants with hSBA titers \geq LLOQ for serogroups A, C, W and Y at baseline and 1 month after vaccination schedule of MenABCWY and MenACWY vaccines. As pre-specified in the protocol, data reported in this outcome measure were presented for the ACWY group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1, and 1 month after vaccination schedule (i.e, at Day 31 for ABCWY group [pooled lots -first dose] and for ACWY Group, and at Day 211 for ABCWY group [pooled lots – second dose])

Notes:

[55] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the MenABCWY and MenACWY vaccines against N. meningitidis serogroups A, C, W, and Y.

End point values	ABCWY_Pooled	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1489	141		
Units: Percentage of participants				
number (confidence interval 95%)				
Men A, Day 1 (N=1452, 137)	9.2 (7.7 to 10.8)	11.7 (6.8 to 18.3)		
Men A, Day 31 (N=132, 133)	79.5 (71.7 to 86.1)	90.2 (83.9 to 94.7)		
Men A, Day 211 (N=1446)	98.6 (97.9 to 99.2)	99999 (99999 to 99999)		
Men C, Day 1 (N=1487, 139)	29.8 (27.5 to 32.2)	28.8 (21.4 to 37.1)		
Men C, Day 31 (N=139,136)	74.8 (66.8 to 81.8)	64.0 (55.3 to 72.0)		
Men C, Day 211 (N=1457)	99.6 (99.1 to 99.8)	99999 (99999 to 99999)		
Men W, Day 1 (N=1473,140)	12.6 (10.9 to 14.4)	12.9 (7.8 to 19.6)		
Men W, Day 31 (N=142,137)	80.3 (72.8 to 86.5)	69.3 (60.9 to 76.9)		
Men W, Day 211 (N=1463)	99.2 (98.7 to 99.6)	99999 (99999 to 99999)		
Men Y, Day 1 (N=1489,141)	12.2 (10.6 to 14.0)	13.5 (8.3 to 20.2)		
Men Y, Day 91 (N=146,140)	82.2 (75.0 to 88.0)	80.0 (72.4 to 86.3)		
Men Y, Day 211 (N=1461)	99.2 (98.7 to 99.6)	99999 (99999 to 99999)		

Statistical analyses

No statistical analyses for this end point

Secondary: Percentage of participants with 4-fold rise in hSBA titers for each of the N. meningitidis serogroups A, C, W and Y at 1 month after the first MenABCWY dose for ABCWY_Pooled group compared to the MenACWY vaccine for ACWY group relative to baseline (Day 1)

End point title	Percentage of participants with 4-fold rise in hSBA titers for each of the N. meningitidis serogroups A, C, W and Y at 1 month after the first MenABCWY dose for ABCWY_Pooled group compared to the MenACWY vaccine for ACWY group relative to baseline (Day 1) ^[56]
-----------------	---

End point description:

The immune response to MenABCWY vaccine compared to MenACWY vaccine was evaluated by measuring bactericidal activity against each of the N. meningitidis serogroups A, C, W and Y at Day 31 compared to baseline (Day 1). Four-fold rise is defined as: - If the pre-vaccination hSBA titer is < 4, then post-vaccination hSBA titer should be ≥ 16. - If the pre-vaccination hSBA titer is ≥ LOD but < LLOQ, then post-vaccination hSBA titer should be ≥ 4 times the LLOQ. - If the pre-vaccination hSBA titer is ≥ LLOQ, then post-vaccination hSBA titer should be ≥ 4 times the pre-vaccination hSBA titer. The corresponding 2- sided 95% CI based on Clopper-Pearson method is calculated for each vaccine group. As pre-specified in the protocol, data reported in this outcome measure were presented for the ACWY group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At 1 month after vaccination schedule (i.e, at Day 31 for ABCWY group [pooled group] and for ACWY

Notes:

[56] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the effectiveness of the rMenB+OMV vaccine compared to one dose of MenACWY vaccination in the ACWY group.

End point values	ABCWY_Pooled	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	146	140		
Units: Percentage of participants				
number (confidence interval 95%)				
Men A (N=127,129)	74.0 (65.5 to 81.4)	86.0 (78.8 to 91.5)		
Men C (N=139,134)	66.9 (58.4 to 74.6)	56.7 (47.9 to 65.2)		
Men W (N=139,136)	74.1 (66.0 to 81.2)	66.2 (57.6 to 74.1)		
Men Y (N=146,140)	76.0 (68.3 to 82.7)	72.1 (63.9 to 79.4)		

Statistical analyses

No statistical analyses for this end point

Secondary: hSBA GMTs against each of the N. meningitidis serogroups A, C, W and Y at Day 1 and 1 month after the first and the last MenABCWY vaccination for the ABCWY_Pooled group and at 1 month after the MenACWY vaccination for ACWY Group

End point title	hSBA GMTs against each of the N. meningitidis serogroups A, C, W and Y at Day 1 and 1 month after the first and the last MenABCWY vaccination for the ABCWY_Pooled group and at 1 month after the MenACWY vaccination for ACWY Group ^[57]
-----------------	--

End point description:

The immune responses to MenABCWY and MenACWY vaccines are evaluated by measuring bactericidal activity against N. meningitidis serogroups A, C, W and Y in terms of GMTs after vaccination compared to baseline (Day 1). For each N. meningitidis serogroups A, C, W and Y, the GMTs (after vaccination/baseline) are calculated, with their associated 2-sided 95% CIs. As pre-specified in the protocol, data reported in this outcome measure were presented for the ACWY group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1, and 1 month after vaccination schedule (i.e, at Day 31 for ABCWY group [pooled lots -first dose] and for ACWY Group, and at Day 211 for ABCWY group [pooled lots – second dose])

Notes:

[57] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the MenABCWY and MenACWY vaccines against N. meningitidis serogroups A, C, W, and Y

End point values	ABCWY_Pooled	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1489	141		
Units: Titers				
geometric mean (confidence interval 95%)				
Men A, Day 1 (N=1452,137)	11.1 (10.3 to 11.9)	12.7 (10.9 to 14.8)		
Men A, Day 31 (N=132,133)	175.3 (121.2 to 253.3)	474.8 (331.3 to 680.3)		
Men A, Day 211 (N=1446)	352.0 (322.0 to 384.7)	99999 (99999 to 99999)		
Men C, Day 1 (N=1487,139)	12.0 (10.9 to 13.3)	11.4 (9.1 to 14.3)		
Men C, Day 31 (N=139,136)	674.8 (355.9 to 1279.4)	379.0 (204.4 to 703.0)		
Men C, Day 211 (N=1457)	1162.5 (1015.7 to 1330.5)	99999 (99999 to 99999)		
Men W, Day 1 (N=1473,140)	8.0 (7.4 to 8.7)	7.4 (6.2 to 8.9)		
Men W, Day 31 (N=142,137)	374.0 (243.4 to 574.8)	194.3 (128.3 to 294.2)		
Men W, Day 211 (N=1463)	666.5 (603.2 to 736.3)	99999 (99999 to 99999)		
Men Y, Day 1 (N=1489,141)	9.3 (8.7 to 10.0)	9.9 (8.6 to 11.5)		
Men Y, Day 31 (N=146,140)	375.4 (246.9 to 570.7)	320.9 (213.8 to 481.7)		
Men Y, Day 211 (N=1461)	655.9 (587.0 to 733.0)	99999 (99999 to 99999)		

Statistical analyses

No statistical analyses for this end point

Secondary: GMRs for each of the N. meningitidis serogroups A, C, W and Y at 1 month after the first and the last MenABCWY vaccination for the ABCWY_Pooled group and at 1 month after the MenACWY vaccination for ACWY Group

End point title	GMRs for each of the N. meningitidis serogroups A, C, W and Y at 1 month after the first and the last MenABCWY vaccination for the ABCWY_Pooled group and at 1 month after the MenACWY vaccination for ACWY Group ^[58]
-----------------	---

End point description:

The immune responses to MenABCWY and MenACWY vaccines are evaluated by measuring bactericidal activity against N. meningitidis serogroups A, C, W and Y. For each N. meningitidis serogroups A, C, W and Y, the GMRs (after vaccination/baseline) are calculated, with their associated 2-sided 95% CIs. As pre-specified in the protocol, data reported in this outcome measure were presented for the ACWY group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

1 month after vaccination schedule (i.e, at Day 31 for ABCWY group [pooled lots -first dose] and for ACWY Group, and at Day 211 for ABCWY group [pooled lots – second dose]) compared to baseline (Day 1)

Notes:

[58] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the MenABCWY and MenACWY against N. meningitidis serogroups A, C, W, and Y.

End point values	ABCWY_Pooled	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	1446	140		
Units: Ratio				
geometric mean (confidence interval 95%)				
Men A, Day 31 (N=127,129)	11.8 (7.9 to 17.7)	31.8 (21.4 to 47.1)		
Men A, Day 211 (N=1397)	31.2 (28.3 to 34.5)	99999 (99999 to 99999)		
Men C, Day 31 (N=139,134)	30.9 (16.9 to 56.2)	22.9 (12.8 to 41.1)		
Men C, Day 211 (N=1439)	96.9 (84.5 to 111.1)	99999 (99999 to 99999)		
Men W, Day 31 (N=139,136)	32.9 (21.7 to 50.0)	23.2 (15.5 to 34.7)		
Men W, Day 211 (N=1432)	83.8 (74.9 to 93.8)	99999 (99999 to 99999)		
Men Y, Day 31 (N=146,140)	28.1 (18.3 to 43.2)	25.6 (16.9 to 38.8)		
Men Y, Day 211 (N=1446)	70.2 (62.3 to 79.1)	99999 (99999 to 99999)		

Statistical analyses

No statistical analyses for this end point

Secondary: Total Immunoglobulin G (IgG) antibodies concentrations against N. meningitidis serogroups A, C, W and Y at Day 1 and 1 month after the first and the last MenABCWY vaccination for ABCWY_Pooled group and 1 month after the MenACWY vaccination for ACWY Group

End point title	Total Immunoglobulin G (IgG) antibodies concentrations against N. meningitidis serogroups A, C, W and Y at Day 1 and 1 month after the first and the last MenABCWY vaccination for ABCWY_Pooled group and 1 month after the MenACWY vaccination for ACWY Group ^[59]
-----------------	--

End point description:

The immune responses to MenABCWY and MenACWY vaccines were evaluated by measuring the total IgG in terms of electrochemiluminescence-based multiplex (EGL) geometric mean concentrations (GMCs) which was an alternative assay to Enzyme-Linked Immunosorbent Assay (ELISA). EGL (validated assay) was used because ELISA is not validated. As pre- specified in the protocol, data reported in this outcome measure were presented for the ACWY group and ABCWY pooled group, participants from the ABCWY-1, ABCWY-2, and ABCWY-3 groups were pooled into a single group. Analysis was performed on the FAS.

End point type	Secondary
----------------	-----------

End point timeframe:

At Day 1, and 1 month after vaccination schedule (i.e, at Day 31 for ABCWY group [pooled lots -first dose] and for ACWY Group, and at Day 211 for ABCWY group [pooled lots - second dose])

Notes:

[59] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: As specified in the Protocol, the analysis assesses the immune response of the MenABCWY and MenACWY vaccines against N. meningitidis serogroups A, C, W, and Y.

End point values	ABCWY_Pooled	ACWY Group		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	179	172		
Units: microgram per milliliter(µg/mL)				
geometric mean (confidence interval 95%)				
Men A, Day 1 (N=170, 179)	1.9 (1.6 to 2.2)	2.3 (2.0 to 2.7)		
Men A, Day 31 (N=168, 177)	18.1 (13.8 to 23.8)	53.7 (40.8 to 70.7)		
Men A, Day 211 (N=164)	30.2 (23.3 to 39.2)	99999 (99999 to 99999)		
Men C, Day 1 (N=170, 179)	0.7 (0.6 to 0.8)	0.8 (0.7 to 1.0)		
Men C, Day 31 (N=170, 175)	15.5 (11.6 to 20.7)	13.8 (10.4 to 18.4)		
Men C, Day 211 (N=163)	17.0 (13.1 to 22.0)	99999 (99999 to 99999)		
Men W, Day 1 (N=170, 179)	0.5 (0.4 to 0.6)	0.6 (0.5 to 0.7)		
Men W, Day 31 (N=172, 178)	9.1 (6.5 to 12.8)	8.4 (6.0 to 11.6)		
Men W, Day 211 (N=164)	21.7 (16.0 to 29.5)	99999 (99999 to 99999)		
Men Y, Day 1 (N=170, 179)	0.9 (0.7 to 1.0)	0.9 (0.8 to 1.0)		
Men Y, Day 31 (N=172, 178)	12.9 (9.2 to 18.0)	14.3 (10.2 to 19.8)		
Men Y, Day 211 (N=164)	26.3 (19.6 to 35.4)	99999 (99999 to 99999)		

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

SAEs and Non-serious AEs (Other AEs) were collected through the entire period of the study (from Day 1 up to study end [Day 361])

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	25.0
--------------------	------

Reporting groups

Reporting group title	MenB_0_2_6 Group
-----------------------	------------------

Reporting group description:

Participants received 3 doses of rMenB+OMV NZ vaccine on Day 1, Day 61 and Day 181. Participants received 1 dose of MenACWY vaccine at Day 211 as a standard of care.

Reporting group title	MenB_0_6 Group
-----------------------	----------------

Reporting group description:

Participants received 2 doses of rMenB+OMV NZ vaccine on Day 1, and Day 181, 1 dose of MenACWY vaccine on Day 61. Participants received 1 dose of Placebo on Day 211 to maintain blinding.

Reporting group title	ABCWY-3 Group
-----------------------	---------------

Reporting group description:

Participants received 2 doses of MenABCWY lot 3 vaccine at Day 1 and Day 181 and 2 doses of placebo at Day 61 and Day 211.

Reporting group title	ABCWY-1 Group
-----------------------	---------------

Reporting group description:

Participants received 2 doses of MenABCWY lot 1 vaccine at Day 1 and Day 181 and 2 doses of placebo at Day 61 and Day 211.

Reporting group title	ABCWY-2 Group
-----------------------	---------------

Reporting group description:

Participants received 2 doses of MenABCWY lot 2 vaccine at Day 1 and Day 181 and 2 doses of placebo at Day 61 and Day 211.

Reporting group title	ACWY Group
-----------------------	------------

Reporting group description:

Participants received 1 dose of MenACWY vaccine at Day 1, 1 dose of placebo at Day 61 and 1 dose of rMenB+OMV NZ vaccine on Day 181. Participants received 1 dose of rMenB+OMV NZ vaccine on Day 211 as standard of care.

Serious adverse events	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY-3 Group
Total subjects affected by serious adverse events			
subjects affected / exposed	20 / 897 (2.23%)	22 / 906 (2.43%)	8 / 554 (1.44%)
number of deaths (all causes)	1	1	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Testis cancer			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Brain stem glioma			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ovarian fibroma			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Haematoma			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
Complication of pregnancy			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Abortion spontaneous			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pre-eclampsia			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Placental insufficiency			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

General disorders and administration site conditions			
Chest pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyrexia			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			
Anaphylactic reaction			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders			
Ovarian cyst ruptured			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Dyspnoea			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Depression suicidal			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Depression			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anxiety			

subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Major depression			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Substance-induced psychotic disorder			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicide attempt			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Mental disorder			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anorexia nervosa			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicidal ideation			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychotic disorder			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Thinking abnormal			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Troponin T increased			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Concussion			
subjects affected / exposed	3 / 897 (0.33%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 3	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Alcohol poisoning			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 1	0 / 0	0 / 0
Intentional overdose			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Lower limb fracture			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Post procedural haemorrhage			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Poisoning			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 1	0 / 0	0 / 0

Overdose			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 1	0 / 0
Road traffic accident			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin laceration			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Subdural haemorrhage			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Traumatic liver injury			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Toxicity to various agents			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tendon rupture			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ulna fracture			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury			

subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Brain contusion			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Back injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hand fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Head injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neck injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skull fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tibia fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Congenital, familial and genetic disorders			
Urachal abnormality			

subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Aphasia			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Headache			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dysarthria			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dizziness			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorder			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Paresis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Petit mal epilepsy			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Speech disorder			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Subarachnoid haemorrhage			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neuromyelitis optica spectrum disorder			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Lymphadenitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
Hypermetropia			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nausea			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Colitis ulcerative			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	1 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Cholelithiasis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatitis toxic			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
Excessive granulation tissue			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vancomycin infusion reaction			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Nephrolithiasis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urethral stenosis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Rhabdomyolysis			

subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Intervertebral disc protrusion			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Appendicitis			
subjects affected / exposed	1 / 897 (0.11%)	3 / 906 (0.33%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 1	0 / 3	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cellulitis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Peritonsillar abscess			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tooth abscess			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tonsillitis			

subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infected dermal cyst			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pharyngeal abscess			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Post-acute COVID-19 syndrome			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Type 1 diabetes mellitus			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	ABCWY-1 Group	ABCWY-2 Group	ACWY Group
Total subjects affected by serious adverse events			
subjects affected / exposed	8 / 549 (1.46%)	9 / 554 (1.62%)	5 / 178 (2.81%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Testis cancer			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Brain stem glioma			

subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ovarian fibroma			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vascular disorders			
Haematoma			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
Complication of pregnancy			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Abortion spontaneous			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pre-eclampsia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Placental insufficiency			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
General disorders and administration site conditions			
Chest pain			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pyrexia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Immune system disorders			
Anaphylactic reaction			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Reproductive system and breast disorders			
Ovarian cyst ruptured			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Dyspnoea			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Depression suicidal			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Depression			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anxiety			

subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Major depression			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Substance-induced psychotic disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicide attempt			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Mental disorder			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Anorexia nervosa			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Suicidal ideation			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychotic disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Thinking abnormal			

subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Investigations			
Troponin T increased			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Concussion			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Alcohol poisoning			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Intentional overdose			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Lower limb fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Post procedural haemorrhage			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Poisoning			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Overdose			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Road traffic accident			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin laceration			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Subdural haemorrhage			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Traumatic liver injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Toxicity to various agents			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tendon rupture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Ulna fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Brain contusion			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Back injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hand fracture			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Head injury			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neck injury			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skull fracture			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tibia fracture			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Congenital, familial and genetic disorders			
Urachal abnormality			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorders			
Aphasia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Headache			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dysarthria			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Dizziness			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nervous system disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Paresis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Petit mal epilepsy			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Speech disorder			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Subarachnoid haemorrhage			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Neuromyelitis optica spectrum disorder			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Blood and lymphatic system disorders			
Lymphadenitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Eye disorders			
Hypermetropia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vomiting			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Nausea			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Colitis ulcerative			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Cholelithiasis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatitis toxic			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
Excessive granulation tissue			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Vancomycin infusion reaction			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Renal and urinary disorders			
Nephrolithiasis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Urethral stenosis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Musculoskeletal and connective tissue disorders			
Rhabdomyolysis			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Intervertebral disc protrusion			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Appendicitis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cellulitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastroenteritis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Peritonsillar abscess			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Sepsis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tooth abscess			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Tonsillitis			

subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infected dermal cyst			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pharyngeal abscess			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Post-acute COVID-19 syndrome			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Metabolism and nutrition disorders			
Type 1 diabetes mellitus			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Frequency threshold for reporting non-serious adverse events: 0 %

Non-serious adverse events	MenB_0_2_6 Group	MenB_0_6 Group	ABCWY-3 Group
Total subjects affected by non-serious adverse events			
subjects affected / exposed	876 / 897 (97.66%)	872 / 906 (96.25%)	544 / 554 (98.19%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Skin papilloma			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	1	2	0
Benign soft tissue neoplasm			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Basal cell carcinoma			

subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Melanocytic naevus subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Fibroadenoma of breast subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Lipoma subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Vascular disorders			
Pallor subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Varicose vein subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Haematoma subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Hot flush subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Hypertension subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	2 / 906 (0.22%) 2	0 / 554 (0.00%) 0
Peripheral venous disease subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Hyperaemia subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Hypotension subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1

Orthostatic hypotension subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
General disorders and administration site conditions			
Fatigue subjects affected / exposed occurrences (all)	604 / 897 (67.34%) 1183	579 / 906 (63.91%) 1005	358 / 554 (64.62%) 597
Feeling hot subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	2 / 554 (0.36%) 2
Administration site erythema subjects affected / exposed occurrences (all)	209 / 897 (23.30%) 300	157 / 906 (17.33%) 203	94 / 554 (16.97%) 120
Administration site induration subjects affected / exposed occurrences (all)	138 / 897 (15.38%) 180	113 / 906 (12.47%) 142	72 / 554 (13.00%) 88
Cyst subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Administration site swelling subjects affected / exposed occurrences (all)	202 / 897 (22.52%) 293	158 / 906 (17.44%) 200	95 / 554 (17.15%) 121
Asthenia subjects affected / exposed occurrences (all)	2 / 897 (0.22%) 2	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Chest pain subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	2 / 906 (0.22%) 2	1 / 554 (0.18%) 1
Chills subjects affected / exposed occurrences (all)	8 / 897 (0.89%) 8	3 / 906 (0.33%) 4	3 / 554 (0.54%) 3
Administration site pain subjects affected / exposed occurrences (all)	851 / 897 (94.87%) 2229	853 / 906 (94.15%) 1768	524 / 554 (94.58%) 1028
Induration			

subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Influenza like illness			
subjects affected / exposed	2 / 897 (0.22%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	2	3	0
Injection site bruising			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	1	0	1
Injection site haematoma			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Injection site hypoaesthesia			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Injection site warmth			
subjects affected / exposed	2 / 897 (0.22%)	1 / 906 (0.11%)	2 / 554 (0.36%)
occurrences (all)	3	1	2
Injection site mass			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	1	2	1
Injection site pain			
subjects affected / exposed	3 / 897 (0.33%)	8 / 906 (0.88%)	3 / 554 (0.54%)
occurrences (all)	3	8	3
Injection site pruritus			
subjects affected / exposed	2 / 897 (0.22%)	1 / 906 (0.11%)	3 / 554 (0.54%)
occurrences (all)	4	2	3
Injection site rash			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	2 / 554 (0.36%)
occurrences (all)	1	2	2
Injection site swelling			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Injection site induration			
subjects affected / exposed	8 / 897 (0.89%)	5 / 906 (0.55%)	5 / 554 (0.90%)
occurrences (all)	11	5	5
Malaise			

subjects affected / exposed	4 / 897 (0.45%)	3 / 906 (0.33%)	0 / 554 (0.00%)
occurrences (all)	5	3	0
Non-cardiac chest pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Pain			
subjects affected / exposed	4 / 897 (0.45%)	2 / 906 (0.22%)	3 / 554 (0.54%)
occurrences (all)	5	2	3
Peripheral swelling			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Pyrexia			
subjects affected / exposed	66 / 897 (7.36%)	59 / 906 (6.51%)	32 / 554 (5.78%)
occurrences (all)	69	62	32
Vaccination site warmth			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	1	0	1
Thirst			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Vaccination site pain			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Vaccination site pruritus			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Vaccination site urticaria			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Swelling			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	1	0	1
Vessel puncture site pain			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	0	2	0
Axillary pain			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Chest discomfort			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Discomfort			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Injection site papule			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Injection site discomfort			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Injection site eczema			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Injection site erythema			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Injection site haemorrhage			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Feeling cold			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Injection site paraesthesia			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Injection site reaction			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Medical device pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Oedema peripheral			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Swelling face			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Vaccination site erythema			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Vaccination site reaction			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Vaccination site bruising			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Immune system disorders			
Anaphylactoid reaction			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Drug hypersensitivity			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Food allergy			
subjects affected / exposed	2 / 897 (0.22%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	2	1	1
Hypersensitivity			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Multiple allergies			
subjects affected / exposed	2 / 897 (0.22%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	2	1	0
Seasonal allergy			
subjects affected / exposed	7 / 897 (0.78%)	7 / 906 (0.77%)	6 / 554 (1.08%)
occurrences (all)	7	7	6
Allergy to animal			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1

Allergy to arthropod sting subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Dust allergy subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Social circumstances Menarche subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Reproductive system and breast disorders			
Breast cyst subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Ovarian cyst subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Nipple enlargement subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Menstruation irregular subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Intermenstrual bleeding subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Heavy menstrual bleeding subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Penile rash subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Dysmenorrhoea subjects affected / exposed occurrences (all)	14 / 897 (1.56%) 19	10 / 906 (1.10%) 10	10 / 554 (1.81%) 14
Breast tenderness			

subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Breast mass			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Breast inflammation			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Endometriosis			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Vaginal haemorrhage			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	1
Testicular pain			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Premenstrual syndrome			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Abnormal uterine bleeding			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Amenorrhoea			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Bartholin's cyst			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Breast discharge			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Breast pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Menstruation delayed			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Premenstrual pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Uterine spasm			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Breast haematoma			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Respiratory, thoracic and mediastinal disorders			
Cough			
subjects affected / exposed	4 / 897 (0.45%)	3 / 906 (0.33%)	7 / 554 (1.26%)
occurrences (all)	4	4	7
Asthma			
subjects affected / exposed	4 / 897 (0.45%)	6 / 906 (0.66%)	0 / 554 (0.00%)
occurrences (all)	5	8	0
Dyspnoea			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Rhinitis allergic			
subjects affected / exposed	2 / 897 (0.22%)	3 / 906 (0.33%)	0 / 554 (0.00%)
occurrences (all)	2	3	0
Rhinalgia			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Respiratory tract congestion			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Oropharyngeal pain			
subjects affected / exposed	12 / 897 (1.34%)	11 / 906 (1.21%)	9 / 554 (1.62%)
occurrences (all)	12	12	9
Nasal congestion			

subjects affected / exposed	12 / 897 (1.34%)	6 / 906 (0.66%)	7 / 554 (1.26%)
occurrences (all)	13	6	7
Epistaxis			
subjects affected / exposed	1 / 897 (0.11%)	3 / 906 (0.33%)	3 / 554 (0.54%)
occurrences (all)	1	3	3
Rhinorrhoea			
subjects affected / exposed	4 / 897 (0.45%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	4	1	1
Wheezing			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Upper-airway cough syndrome			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Tonsillolith			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Sneezing			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	2	0	1
Bronchial hyperreactivity			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Hiccups			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Increased upper airway secretion			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Pleuritic pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Respiratory disorder			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Throat clearing			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Psychiatric disorders			
Adjustment disorder with depressed mood			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Adjustment disorder			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Aggression			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Anorexia nervosa			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	1
Anxiety			
subjects affected / exposed	6 / 897 (0.67%)	8 / 906 (0.88%)	3 / 554 (0.54%)
occurrences (all)	6	8	3
Anxiety disorder			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	2	0	1
Attention deficit hyperactivity disorder			
subjects affected / exposed	0 / 897 (0.00%)	4 / 906 (0.44%)	1 / 554 (0.18%)
occurrences (all)	0	4	1
Binge eating			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Confusional state			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Depression			
subjects affected / exposed	6 / 897 (0.67%)	4 / 906 (0.44%)	4 / 554 (0.72%)
occurrences (all)	6	4	4
Depression suicidal			

subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Disruptive mood dysregulation disorder			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Insomnia			
subjects affected / exposed	3 / 897 (0.33%)	3 / 906 (0.33%)	5 / 554 (0.90%)
occurrences (all)	3	3	5
Panic attack			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	3	0	1
Sleep disorder			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	1	1	1
Stress			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Suicidal ideation			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Tic			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Mixed anxiety and depressive disorder			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Acute stress disorder			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Anger			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Hallucination			

subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Irritability subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
School refusal subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Obsessive-compulsive disorder subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Investigations			
Serum ferritin decreased subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
SARS-CoV-2 test positive subjects affected / exposed occurrences (all)	6 / 897 (0.67%) 6	7 / 906 (0.77%) 7	3 / 554 (0.54%) 3
Liver function test increased subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Heart rate irregular subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Haemoglobin decreased subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Computerised tomogram abdomen abnormal subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Cardiac murmur subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Body temperature increased			

subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Thyroid hormones decreased subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Streptococcus test positive subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Red blood cell count increased subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Blood pressure increased subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Heart rate increased subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Weight decreased subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Injury, poisoning and procedural complications			
Alcohol poisoning subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Ankle fracture subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Arthropod bite subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	1 / 554 (0.18%) 1
Arthropod sting subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Clavicle fracture			

subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Fibula fracture			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	0	2	0
Contusion			
subjects affected / exposed	3 / 897 (0.33%)	2 / 906 (0.22%)	2 / 554 (0.36%)
occurrences (all)	3	2	5
Eye abrasion			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Eye injury			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Facial bones fracture			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	2	0	1
Fall			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	2 / 554 (0.36%)
occurrences (all)	1	2	2
Concussion			
subjects affected / exposed	3 / 897 (0.33%)	0 / 906 (0.00%)	2 / 554 (0.36%)
occurrences (all)	3	0	2
Humerus fracture			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Human bite			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Lower limb fracture			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Limb injury			
subjects affected / exposed	3 / 897 (0.33%)	5 / 906 (0.55%)	3 / 554 (0.54%)
occurrences (all)	3	5	3
Ligament sprain			

subjects affected / exposed	5 / 897 (0.56%)	16 / 906 (1.77%)	5 / 554 (0.90%)
occurrences (all)	7	17	5
Ligament rupture			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	1	1	1
Joint injury			
subjects affected / exposed	4 / 897 (0.45%)	3 / 906 (0.33%)	2 / 554 (0.36%)
occurrences (all)	4	3	2
Joint dislocation			
subjects affected / exposed	0 / 897 (0.00%)	4 / 906 (0.44%)	0 / 554 (0.00%)
occurrences (all)	0	4	0
Infusion related reaction			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Foot fracture			
subjects affected / exposed	2 / 897 (0.22%)	5 / 906 (0.55%)	1 / 554 (0.18%)
occurrences (all)	2	6	1
Hand fracture			
subjects affected / exposed	3 / 897 (0.33%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	3	1	2
Head injury			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	2 / 554 (0.36%)
occurrences (all)	1	0	2
Procedural nausea			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Procedural dizziness			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Procedural complication			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Post-traumatic neck syndrome			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Post procedural complication			

subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	1	2	0
Procedural pain			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	1
Musculoskeletal foreign body			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Muscle strain			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Muscle rupture			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	2	0	1
Meniscus injury			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Nail injury			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	0	2	0
Skin abrasion			
subjects affected / exposed	3 / 897 (0.33%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	3	3	1
Scratch			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Road traffic accident			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	0	2	1
Radius fracture			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	1	2	1
Procedural vomiting			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Skin laceration			

subjects affected / exposed	5 / 897 (0.56%)	5 / 906 (0.55%)	2 / 554 (0.36%)
occurrences (all)	5	5	2
Traumatic haematoma			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Torus fracture			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Tibia fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Thermal burn			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	1	0	1
Tendon injury			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Ulna fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	2 / 554 (0.36%)
occurrences (all)	0	0	2
Upper limb fracture			
subjects affected / exposed	1 / 897 (0.11%)	4 / 906 (0.44%)	0 / 554 (0.00%)
occurrences (all)	1	5	0
Wrist fracture			
subjects affected / exposed	2 / 897 (0.22%)	2 / 906 (0.22%)	2 / 554 (0.36%)
occurrences (all)	2	2	2
Vaccination complication			
subjects affected / exposed	2 / 897 (0.22%)	3 / 906 (0.33%)	0 / 554 (0.00%)
occurrences (all)	2	3	0
Anaesthetic complication			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Animal bite			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Animal scratch			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Bone contusion			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Bursa injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Compression fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Ear canal injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Foreign body in respiratory tract			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Ligament injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Limb crushing injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Limb fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Musculoskeletal injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Nasal injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Post procedural haemorrhage			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Soft tissue injury			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Sports injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Stress fracture			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Tooth injury			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Traumatic haemorrhage			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Wound			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Sunburn			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Congenital, familial and genetic disorders			
Familial mediterranean fever			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Multiple endocrine neoplasia Type 1			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Rathke's cleft cyst			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Type V hyperlipidaemia			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Dermoid cyst			

subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Pectus excavatum subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Cardiac disorders			
Supraventricular tachycardia subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Palpitations subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	1 / 554 (0.18%) 1
Tachycardia paroxysmal subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Tachycardia subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Nervous system disorders			
Myoclonus subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Migraine subjects affected / exposed occurrences (all)	3 / 897 (0.33%) 3	5 / 906 (0.55%) 5	2 / 554 (0.36%) 2
Lethargy subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	2 / 906 (0.22%) 2	0 / 554 (0.00%) 0
Hypoaesthesia subjects affected / exposed occurrences (all)	2 / 897 (0.22%) 2	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Headache subjects affected / exposed occurrences (all)	578 / 897 (64.44%) 999	525 / 906 (57.95%) 874	320 / 554 (57.76%) 541
Dyskinesia			

subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Dizziness			
subjects affected / exposed	6 / 897 (0.67%)	6 / 906 (0.66%)	3 / 554 (0.54%)
occurrences (all)	8	6	3
Taste disorder			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Syncope			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	7 / 554 (1.26%)
occurrences (all)	1	2	7
Speech disorder			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Seizure			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Psychomotor hyperactivity			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Presyncope			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	4 / 554 (0.72%)
occurrences (all)	1	1	4
Neuromuscular blockade			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Somnolence			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Carpal tunnel syndrome			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Neuralgia			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Paraesthesia			

subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Restless legs syndrome subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Sleep deficit subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Tension headache subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Blood and lymphatic system disorders			
Anaemia subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	2 / 554 (0.36%) 3
Lymphadenopathy subjects affected / exposed occurrences (all)	5 / 897 (0.56%) 6	7 / 906 (0.77%) 7	3 / 554 (0.54%) 3
Iron deficiency anaemia subjects affected / exposed occurrences (all)	4 / 897 (0.45%) 4	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Lymphadenitis subjects affected / exposed occurrences (all)	2 / 897 (0.22%) 2	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Coagulopathy subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Increased tendency to bruise subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Ear and labyrinth disorders			
Tinnitus subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Ear discomfort			

subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Ear pain			
subjects affected / exposed	3 / 897 (0.33%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	3	2	1
Eustachian tube dysfunction			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Hypoacusis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Cerumen impaction			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	2 / 554 (0.36%)
occurrences (all)	0	2	2
Tympanic membrane perforation			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Vertigo			
subjects affected / exposed	4 / 897 (0.45%)	3 / 906 (0.33%)	0 / 554 (0.00%)
occurrences (all)	4	3	0
Excessive cerumen production			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
External ear inflammation			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Motion sickness			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Tympanosclerosis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Eye disorders			
Astigmatism			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0

Eye swelling			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Blindness transient			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Conjunctivitis allergic			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Eye irritation			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Eye pain			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Blepharitis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Glaucoma			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Keratitis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Ocular hyperaemia			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Swelling of eyelid			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Vision blurred			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Conjunctival haemorrhage			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0

Chalazion			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Eye inflammation			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Eye pruritus			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Eyelid exfoliation			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Eyelid ptosis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	0
Myopia			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Gastrointestinal disorders			
Colitis			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	0	2	0
Abdominal distension			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Abdominal pain			
subjects affected / exposed	7 / 897 (0.78%)	11 / 906 (1.21%)	4 / 554 (0.72%)
occurrences (all)	7	12	7
Abdominal pain lower			
subjects affected / exposed	3 / 897 (0.33%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	3	0	0
Abdominal pain upper			
subjects affected / exposed	7 / 897 (0.78%)	9 / 906 (0.99%)	9 / 554 (1.62%)
occurrences (all)	7	10	10
Anal fissure			

subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Abdominal discomfort			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	1	2	1
Constipation			
subjects affected / exposed	3 / 897 (0.33%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	3	1	1
Dental caries			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Diarrhoea			
subjects affected / exposed	14 / 897 (1.56%)	16 / 906 (1.77%)	3 / 554 (0.54%)
occurrences (all)	15	19	3
Dyspepsia			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	2 / 554 (0.36%)
occurrences (all)	0	0	2
Haemorrhoids			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	0	2	0
Food poisoning			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	2 / 554 (0.36%)
occurrences (all)	0	2	2
Gastritis			
subjects affected / exposed	1 / 897 (0.11%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	1	2	1
Gastrooesophageal reflux disease			
subjects affected / exposed	2 / 897 (0.22%)	4 / 906 (0.44%)	0 / 554 (0.00%)
occurrences (all)	2	4	0
Haematemesis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Enteritis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Hyperchlorhydria			

subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Hypoaesthesia oral			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Irritable bowel syndrome			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	0	2	1
Lip swelling			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	1	1	1
Mouth ulceration			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Toothache			
subjects affected / exposed	1 / 897 (0.11%)	5 / 906 (0.55%)	2 / 554 (0.36%)
occurrences (all)	1	6	2
Oral pain			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Oral pruritus			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Pancreatitis relapsing			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Peptic ulcer			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Nausea			
subjects affected / exposed	245 / 897 (27.31%)	204 / 906 (22.52%)	139 / 554 (25.09%)
occurrences (all)	324	262	170
Vomiting			
subjects affected / exposed	6 / 897 (0.67%)	11 / 906 (1.21%)	4 / 554 (0.72%)
occurrences (all)	6	11	4
Palatal disorder			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Coeliac disease			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Embedded tooth			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Eruption			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Flatulence			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Gingival pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Aphthous ulcer			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Tongue discolouration			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Transient lingual papillitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Hepatobiliary disorders			
Hepatic steatosis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Skin and subcutaneous tissue disorders			
Dermatitis contact			
subjects affected / exposed	3 / 897 (0.33%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	3	2	2
Dermatitis atopic			

subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	2 / 554 (0.36%)
occurrences (all)	1	1	2
Dermatitis allergic			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Dermatitis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Cold urticaria			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Acne			
subjects affected / exposed	8 / 897 (0.89%)	7 / 906 (0.77%)	4 / 554 (0.72%)
occurrences (all)	8	7	5
Miliaria			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Pain of skin			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Pityriasis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Pruritus			
subjects affected / exposed	3 / 897 (0.33%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	3	0	0
Rash			
subjects affected / exposed	3 / 897 (0.33%)	2 / 906 (0.22%)	4 / 554 (0.72%)
occurrences (all)	3	2	4
Rash macular			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Ingrowing nail			
subjects affected / exposed	3 / 897 (0.33%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	3	1	1
Hyperhidrosis			

subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	0	2	1
Hand dermatitis			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	1	1	1
Erythema			
subjects affected / exposed	3 / 897 (0.33%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	4	0	0
Rash maculo-papular			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	2	0	1
Eczema			
subjects affected / exposed	4 / 897 (0.45%)	2 / 906 (0.22%)	2 / 554 (0.36%)
occurrences (all)	4	2	2
Rash pruritic			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Skin lesion			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Urticaria			
subjects affected / exposed	2 / 897 (0.22%)	3 / 906 (0.33%)	1 / 554 (0.18%)
occurrences (all)	2	3	1
Rash erythematous			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Alopecia			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Dermatitis acneiform			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Drug eruption			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Nail disorder			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Psoriasis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	2 / 554 (0.36%)
occurrences (all)	0	0	2
Seborrhoeic dermatitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Skin induration			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Urticaria chronic			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Renal and urinary disorders			
Haematuria			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Dysuria			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	3
Urinary tract inflammation			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Urinary retention			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Pollakiuria			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Nephrolithiasis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	2 / 554 (0.36%)
occurrences (all)	0	1	2
Nephritis			
subjects affected / exposed	3 / 897 (0.33%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	3	0	0

Hydronephrosis subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Endocrine disorders			
Hypothyroidism subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	1 / 554 (0.18%) 1
Thyroid stimulating hormone deficiency subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Growth hormone deficiency subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Hyperthyroidism subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Thyroid cyst subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Musculoskeletal and connective tissue disorders			
Arthralgia subjects affected / exposed occurrences (all)	154 / 897 (17.17%) 205	127 / 906 (14.02%) 163	76 / 554 (13.72%) 90
Arthritis subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Axillary mass subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Back pain subjects affected / exposed occurrences (all)	7 / 897 (0.78%) 7	15 / 906 (1.66%) 15	2 / 554 (0.36%) 2
Muscle swelling subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Knee deformity			

subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Joint warmth			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Joint swelling			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Joint hyperextension			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Muscular weakness			
subjects affected / exposed	0 / 897 (0.00%)	3 / 906 (0.33%)	0 / 554 (0.00%)
occurrences (all)	0	3	0
Growing pains			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Foot deformity			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Flank pain			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Epiphysiolysis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Muscle tightness			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	1
Intervertebral disc protrusion			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	1
Spinal pain			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Sever's disease			

subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Rotator cuff syndrome			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Pain in extremity			
subjects affected / exposed	5 / 897 (0.56%)	11 / 906 (1.21%)	5 / 554 (0.90%)
occurrences (all)	5	11	8
Osteochondrosis			
subjects affected / exposed	4 / 897 (0.45%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	4	1	0
Neck pain			
subjects affected / exposed	1 / 897 (0.11%)	3 / 906 (0.33%)	0 / 554 (0.00%)
occurrences (all)	1	3	0
Neck mass			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Myositis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Myalgia			
subjects affected / exposed	240 / 897 (26.76%)	209 / 906 (23.07%)	130 / 554 (23.47%)
occurrences (all)	325	276	169
Musculoskeletal chest pain			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	0	2	0
Temporomandibular joint syndrome			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Tendon pain			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Tenosynovitis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	1	0	1
Myokymia			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Bone pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Coccydynia			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Costochondritis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Joint effusion			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Pain in jaw			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Rhabdomyolysis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Sacral pain			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Scoliosis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Short stature			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Synovial cyst			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Tendonitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Torticollis			

subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Spinal flattening subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Infections and infestations			
Acarodermatitis subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	1 / 554 (0.18%) 1
Acute sinusitis subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	4 / 906 (0.44%) 4	1 / 554 (0.18%) 1
Adenovirus infection subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Asymptomatic COVID-19 subjects affected / exposed occurrences (all)	3 / 897 (0.33%) 3	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Bacterial vaginosis subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	2 / 906 (0.22%) 2	2 / 554 (0.36%) 2
Body tinea subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Bronchitis subjects affected / exposed occurrences (all)	2 / 897 (0.22%) 2	2 / 906 (0.22%) 2	2 / 554 (0.36%) 2
Bullous impetigo subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Cystitis subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	3 / 906 (0.33%) 3	0 / 554 (0.00%) 0
COVID-19 pneumonia subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0

Cellulitis			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	1 / 554 (0.18%)
occurrences (all)	0	2	1
Chlamydial infection			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Coronavirus infection			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	1	1	1
COVID-19			
subjects affected / exposed	93 / 897 (10.37%)	107 / 906 (11.81%)	61 / 554 (11.01%)
occurrences (all)	96	111	64
Ear infection			
subjects affected / exposed	3 / 897 (0.33%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	3	2	0
Endometritis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Enterobiasis			
subjects affected / exposed	0 / 897 (0.00%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	0	2	0
Enterovirus infection			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Erythema migrans			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Genital herpes			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Eyelid infection			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Folliculitis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	1	0	1

Gastroenteritis			
subjects affected / exposed	6 / 897 (0.67%)	4 / 906 (0.44%)	5 / 554 (0.90%)
occurrences (all)	6	5	5
Gastroenteritis viral			
subjects affected / exposed	3 / 897 (0.33%)	6 / 906 (0.66%)	2 / 554 (0.36%)
occurrences (all)	3	6	2
Eye infection			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Infectious mononucleosis			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Influenza			
subjects affected / exposed	10 / 897 (1.11%)	11 / 906 (1.21%)	7 / 554 (1.26%)
occurrences (all)	10	12	7
Localised infection			
subjects affected / exposed	1 / 897 (0.11%)	4 / 906 (0.44%)	2 / 554 (0.36%)
occurrences (all)	1	4	2
Lower respiratory tract infection			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Lyme disease			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	1	1	1
Infected bite			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Gingivitis			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Helicobacter gastritis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Herpes zoster			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	2 / 554 (0.36%)
occurrences (all)	1	0	2

Hordeolum			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Impetigo			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	2	0	1
Mycoplasma genitalium infection			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Nasopharyngitis			
subjects affected / exposed	29 / 897 (3.23%)	37 / 906 (4.08%)	24 / 554 (4.33%)
occurrences (all)	33	42	27
Onychomycosis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Oral herpes			
subjects affected / exposed	0 / 897 (0.00%)	5 / 906 (0.55%)	0 / 554 (0.00%)
occurrences (all)	0	6	0
Otitis externa			
subjects affected / exposed	4 / 897 (0.45%)	4 / 906 (0.44%)	1 / 554 (0.18%)
occurrences (all)	5	4	1
Otitis media			
subjects affected / exposed	6 / 897 (0.67%)	3 / 906 (0.33%)	1 / 554 (0.18%)
occurrences (all)	6	3	1
Pharyngitis			
subjects affected / exposed	8 / 897 (0.89%)	19 / 906 (2.10%)	9 / 554 (1.62%)
occurrences (all)	9	24	9
Otosalpingitis			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Paronychia			
subjects affected / exposed	2 / 897 (0.22%)	1 / 906 (0.11%)	3 / 554 (0.54%)
occurrences (all)	2	1	3
Parotitis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0

Pelvic inflammatory disease subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Otitis media acute subjects affected / exposed occurrences (all)	3 / 897 (0.33%) 4	1 / 906 (0.11%) 2	0 / 554 (0.00%) 0
Pharyngitis streptococcal subjects affected / exposed occurrences (all)	2 / 897 (0.22%) 2	4 / 906 (0.44%) 4	1 / 554 (0.18%) 1
Postoperative wound infection subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Pulpitis dental subjects affected / exposed occurrences (all)	2 / 897 (0.22%) 2	2 / 906 (0.22%) 2	1 / 554 (0.18%) 1
Respiratory tract infection subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Respiratory tract infection bacterial subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Skin infection subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Rhinitis subjects affected / exposed occurrences (all)	4 / 897 (0.45%) 4	8 / 906 (0.88%) 8	2 / 554 (0.36%) 2
Rhinovirus infection subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Sialoadenitis subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Sinusitis subjects affected / exposed occurrences (all)	4 / 897 (0.45%) 4	3 / 906 (0.33%) 3	2 / 554 (0.36%) 2

Respiratory tract infection viral subjects affected / exposed occurrences (all)	2 / 897 (0.22%) 3	7 / 906 (0.77%) 8	4 / 554 (0.72%) 6
Soft tissue infection subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Streptococcal infection subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Suspected COVID-19 subjects affected / exposed occurrences (all)	5 / 897 (0.56%) 5	6 / 906 (0.66%) 6	5 / 554 (0.90%) 6
Tinea versicolour subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Tonsillitis subjects affected / exposed occurrences (all)	10 / 897 (1.11%) 10	15 / 906 (1.66%) 17	14 / 554 (2.53%) 14
Upper respiratory tract infection subjects affected / exposed occurrences (all)	35 / 897 (3.90%) 38	42 / 906 (4.64%) 46	25 / 554 (4.51%) 26
Tooth abscess subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	3 / 906 (0.33%) 3	0 / 554 (0.00%) 0
Tooth infection subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Tracheitis subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Trichomoniasis subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Tonsillitis streptococcal subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	2 / 906 (0.22%) 2	0 / 554 (0.00%) 0

Upper respiratory tract infection bacterial			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	1
Urinary tract infection			
subjects affected / exposed	9 / 897 (1.00%)	9 / 906 (0.99%)	3 / 554 (0.54%)
occurrences (all)	11	10	3
Vaginal infection			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	1 / 554 (0.18%)
occurrences (all)	0	1	1
Varicella			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Viral infection			
subjects affected / exposed	4 / 897 (0.45%)	5 / 906 (0.55%)	1 / 554 (0.18%)
occurrences (all)	5	6	1
Yersinia infection			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Viral upper respiratory tract infection			
subjects affected / exposed	3 / 897 (0.33%)	4 / 906 (0.44%)	3 / 554 (0.54%)
occurrences (all)	3	4	3
Vulvovaginal candidiasis			
subjects affected / exposed	2 / 897 (0.22%)	2 / 906 (0.22%)	0 / 554 (0.00%)
occurrences (all)	2	2	0
Vulvovaginal mycotic infection			
subjects affected / exposed	1 / 897 (0.11%)	3 / 906 (0.33%)	2 / 554 (0.36%)
occurrences (all)	1	3	2
Wound infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Viral pharyngitis			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Conjunctivitis			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	4 / 554 (0.72%)
occurrences (all)	0	0	4
Conjunctivitis viral			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Ear lobe infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Fungal foot infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Fungal skin infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Gastrointestinal viral infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Oral candidiasis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Helminthic infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Herpes simplex			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Laryngitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Laryngotracheitis obstructive			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Molluscum contagiosum			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Hand-foot-and-mouth disease			

subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Subcutaneous abscess			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Tinea pedis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Vaccination site cellulitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Vaccination site pustule			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Vulvovaginitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Skin bacterial infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Parasitic gastroenteritis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Pericoronitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Pharyngotonsillitis			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Post-acute COVID-19 syndrome			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Respiratory syncytial virus infection			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	0	0	0
Metabolism and nutrition disorders			

Gluten sensitivity			
subjects affected / exposed	0 / 897 (0.00%)	0 / 906 (0.00%)	1 / 554 (0.18%)
occurrences (all)	0	0	1
Hypercholesterolaemia			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Hyperlipidaemia			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Hyponatraemia			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0
Insulin resistance			
subjects affected / exposed	0 / 897 (0.00%)	3 / 906 (0.33%)	0 / 554 (0.00%)
occurrences (all)	0	3	0
Iron deficiency			
subjects affected / exposed	1 / 897 (0.11%)	3 / 906 (0.33%)	3 / 554 (0.54%)
occurrences (all)	1	3	3
Folate deficiency			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Abnormal loss of weight			
subjects affected / exposed	1 / 897 (0.11%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	1	1	0
Glucose tolerance impaired			
subjects affected / exposed	2 / 897 (0.22%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	2	0	0
Lactose intolerance			
subjects affected / exposed	1 / 897 (0.11%)	0 / 906 (0.00%)	0 / 554 (0.00%)
occurrences (all)	1	0	0
Obesity			
subjects affected / exposed	2 / 897 (0.22%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	2	1	0
Type 2 diabetes mellitus			
subjects affected / exposed	0 / 897 (0.00%)	1 / 906 (0.11%)	0 / 554 (0.00%)
occurrences (all)	0	1	0

Vitamin B complex deficiency subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	1 / 554 (0.18%) 1
Vitamin B12 deficiency subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	1 / 554 (0.18%) 1
Vitamin D deficiency subjects affected / exposed occurrences (all)	1 / 897 (0.11%) 1	2 / 906 (0.22%) 2	0 / 554 (0.00%) 0
Zinc deficiency subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	1 / 906 (0.11%) 1	0 / 554 (0.00%) 0
Decreased appetite subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	3 / 554 (0.54%) 3
Haemochromatosis subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Hypertriglyceridaemia subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	1 / 554 (0.18%) 1
Hypoglycaemia subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0
Hypovitaminosis subjects affected / exposed occurrences (all)	0 / 897 (0.00%) 0	0 / 906 (0.00%) 0	0 / 554 (0.00%) 0

Non-serious adverse events	ABCWY-1 Group	ABCWY-2 Group	ACWY Group
Total subjects affected by non-serious adverse events			
subjects affected / exposed	529 / 549 (96.36%)	537 / 554 (96.93%)	173 / 178 (97.19%)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Skin papilloma			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Benign soft tissue neoplasm			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Basal cell carcinoma			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Melanocytic naevus			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Fibroadenoma of breast			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Lipoma			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vascular disorders			
Pallor			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Varicose vein			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Haematoma			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Hot flush			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Hypertension			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Peripheral venous disease			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Hyperaemia			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0

Hypotension subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Orthostatic hypotension subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
General disorders and administration site conditions			
Fatigue subjects affected / exposed occurrences (all)	352 / 549 (64.12%) 601	356 / 554 (64.26%) 609	105 / 178 (58.99%) 174
Feeling hot subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	1 / 178 (0.56%) 1
Administration site erythema subjects affected / exposed occurrences (all)	112 / 549 (20.40%) 134	115 / 554 (20.76%) 143	21 / 178 (11.80%) 24
Administration site induration subjects affected / exposed occurrences (all)	73 / 549 (13.30%) 93	78 / 554 (14.08%) 92	17 / 178 (9.55%) 19
Cyst subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Administration site swelling subjects affected / exposed occurrences (all)	101 / 549 (18.40%) 134	109 / 554 (19.68%) 146	22 / 178 (12.36%) 25
Asthenia subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Chest pain subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	1 / 554 (0.18%) 1	1 / 178 (0.56%) 1
Chills subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	3 / 554 (0.54%) 3	0 / 178 (0.00%) 0
Administration site pain			

subjects affected / exposed	515 / 549 (93.81%)	522 / 554 (94.22%)	149 / 178 (83.71%)
occurrences (all)	1003	1025	240
Induration			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Influenza like illness			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Injection site bruising			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Injection site haematoma			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Injection site hypoaesthesia			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Injection site warmth			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Injection site mass			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Injection site pain			
subjects affected / exposed	5 / 549 (0.91%)	5 / 554 (0.90%)	1 / 178 (0.56%)
occurrences (all)	5	5	1
Injection site pruritus			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	0	3	1
Injection site rash			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Injection site swelling			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Injection site induration			

subjects affected / exposed	3 / 549 (0.55%)	5 / 554 (0.90%)	2 / 178 (1.12%)
occurrences (all)	4	5	2
Malaise			
subjects affected / exposed	1 / 549 (0.18%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	1	2	0
Non-cardiac chest pain			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	1	0	1
Pain			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Peripheral swelling			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Pyrexia			
subjects affected / exposed	37 / 549 (6.74%)	46 / 554 (8.30%)	7 / 178 (3.93%)
occurrences (all)	41	49	7
Vaccination site warmth			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Thirst			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vaccination site pain			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vaccination site pruritus			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Vaccination site urticaria			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Swelling			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Vessel puncture site pain			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Axillary pain			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Chest discomfort			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Discomfort			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Injection site papule			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Injection site discomfort			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Injection site eczema			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Injection site erythema			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Injection site haemorrhage			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Feeling cold			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Injection site paraesthesia			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Injection site reaction			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Medical device pain			

subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Oedema peripheral			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Swelling face			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Vaccination site erythema			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vaccination site reaction			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Vaccination site bruising			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Immune system disorders			
Anaphylactoid reaction			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Drug hypersensitivity			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Food allergy			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Hypersensitivity			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Multiple allergies			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Seasonal allergy			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	2 / 178 (1.12%)
occurrences (all)	2	1	2

Allergy to animal subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Allergy to arthropod sting subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Dust allergy subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Social circumstances Menarche subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Reproductive system and breast disorders			
Breast cyst subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Ovarian cyst subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Nipple enlargement subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Menstruation irregular subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Intermenstrual bleeding subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Heavy menstrual bleeding subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Penile rash subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Dysmenorrhoea			

subjects affected / exposed	9 / 549 (1.64%)	6 / 554 (1.08%)	6 / 178 (3.37%)
occurrences (all)	11	6	6
Breast tenderness			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Breast mass			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Breast inflammation			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Endometriosis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Vaginal haemorrhage			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Testicular pain			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Premenstrual syndrome			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Abnormal uterine bleeding			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Amenorrhoea			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Bartholin's cyst			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Breast discharge			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Breast pain			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Menstruation delayed			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Premenstrual pain			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Uterine spasm			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Breast haematoma			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Respiratory, thoracic and mediastinal disorders			
Cough			
subjects affected / exposed	2 / 549 (0.36%)	7 / 554 (1.26%)	3 / 178 (1.69%)
occurrences (all)	2	7	3
Asthma			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Dyspnoea			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Rhinitis allergic			
subjects affected / exposed	1 / 549 (0.18%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	1	2	1
Rhinalgia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Respiratory tract congestion			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Oropharyngeal pain			

subjects affected / exposed	13 / 549 (2.37%)	7 / 554 (1.26%)	0 / 178 (0.00%)
occurrences (all)	13	7	0
Nasal congestion			
subjects affected / exposed	6 / 549 (1.09%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	8	3	0
Epistaxis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	0	1	1
Rhinorrhoea			
subjects affected / exposed	3 / 549 (0.55%)	4 / 554 (0.72%)	2 / 178 (1.12%)
occurrences (all)	3	4	2
Wheezing			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Upper-airway cough syndrome			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Tonsillolith			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Sneezing			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Bronchial hyperreactivity			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Hiccups			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Increased upper airway secretion			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Pleuritic pain			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Respiratory disorder			

subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Throat clearing			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Psychiatric disorders			
Adjustment disorder with depressed mood			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Adjustment disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Aggression			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Anorexia nervosa			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Anxiety			
subjects affected / exposed	6 / 549 (1.09%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	6	2	0
Anxiety disorder			
subjects affected / exposed	1 / 549 (0.18%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	1	3	0
Attention deficit hyperactivity disorder			
subjects affected / exposed	0 / 549 (0.00%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	0	3	0
Binge eating			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Confusional state			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Depression			

subjects affected / exposed	4 / 549 (0.73%)	6 / 554 (1.08%)	2 / 178 (1.12%)
occurrences (all)	4	6	2
Depression suicidal			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Disruptive mood dysregulation disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Insomnia			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	0	2	1
Panic attack			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Sleep disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Stress			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Suicidal ideation			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Tic			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Mixed anxiety and depressive disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Acute stress disorder			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Anger			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Hallucination			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Irritability			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
School refusal			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Obsessive-compulsive disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Investigations			
Serum ferritin decreased			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
SARS-CoV-2 test positive			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	2	1	1
Liver function test increased			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Heart rate irregular			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Haemoglobin decreased			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Computerised tomogram abdomen abnormal			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Cardiac murmur			

subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Body temperature increased subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Thyroid hormones decreased subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	1 / 178 (0.56%) 1
Streptococcus test positive subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Red blood cell count increased subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Blood pressure increased subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Heart rate increased subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Weight decreased subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Injury, poisoning and procedural complications			
Alcohol poisoning subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Ankle fracture subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	2 / 554 (0.36%) 2	0 / 178 (0.00%) 0
Arthropod bite subjects affected / exposed occurrences (all)	2 / 549 (0.36%) 2	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Arthropod sting			

subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Clavicle fracture			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Fibula fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Contusion			
subjects affected / exposed	2 / 549 (0.36%)	4 / 554 (0.72%)	0 / 178 (0.00%)
occurrences (all)	3	4	0
Eye abrasion			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Eye injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Facial bones fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Fall			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	2 / 178 (1.12%)
occurrences (all)	0	1	2
Concussion			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
Humerus fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Human bite			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Lower limb fracture			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Limb injury			

subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Ligament sprain			
subjects affected / exposed	5 / 549 (0.91%)	4 / 554 (0.72%)	1 / 178 (0.56%)
occurrences (all)	5	4	1
Ligament rupture			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	1	0	1
Joint injury			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Joint dislocation			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	1	1	1
Infusion related reaction			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Foot fracture			
subjects affected / exposed	2 / 549 (0.36%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	2	2	0
Hand fracture			
subjects affected / exposed	2 / 549 (0.36%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	2	3	0
Head injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Procedural nausea			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Procedural dizziness			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Procedural complication			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Post-traumatic neck syndrome			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Post procedural complication			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Procedural pain			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
Musculoskeletal foreign body			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Muscle strain			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
Muscle rupture			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Meniscus injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Nail injury			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Skin abrasion			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	2 / 178 (1.12%)
occurrences (all)	1	1	2
Scratch			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Road traffic accident			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Radius fracture			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	2	1	1
Procedural vomiting			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Skin laceration			
subjects affected / exposed	0 / 549 (0.00%)	3 / 554 (0.54%)	1 / 178 (0.56%)
occurrences (all)	0	3	1
Traumatic haematoma			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Torus fracture			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Tibia fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Thermal burn			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Tendon injury			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Ulna fracture			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	2	0	1
Upper limb fracture			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Wrist fracture			
subjects affected / exposed	4 / 549 (0.73%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	4	3	0
Vaccination complication			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Anaesthetic complication			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Animal bite			

subjects affected / exposed	1 / 549 (0.18%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	1	3	0
Animal scratch			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Bone contusion			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Bursa injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Compression fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Ear canal injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Foreign body in respiratory tract			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Ligament injury			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Limb crushing injury			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Limb fracture			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Musculoskeletal injury			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Nasal injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Post procedural haemorrhage			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Soft tissue injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Sports injury			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Stress fracture			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Tooth injury			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Traumatic haemorrhage			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Wound			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Sunburn			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Congenital, familial and genetic disorders			
Familial mediterranean fever			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Multiple endocrine neoplasia Type 1			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Rathke's cleft cyst			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Type V hyperlipidaemia			

subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Dermoid cyst subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Pectus excavatum subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Cardiac disorders			
Supraventricular tachycardia subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Palpitations subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	2 / 554 (0.36%) 2	0 / 178 (0.00%) 0
Tachycardia paroxysmal subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Tachycardia subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	2 / 554 (0.36%) 2	0 / 178 (0.00%) 0
Nervous system disorders			
Myoclonus subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Migraine subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	2 / 554 (0.36%) 2	0 / 178 (0.00%) 0
Lethargy subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Hypoaesthesia subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Headache			

subjects affected / exposed	300 / 549 (54.64%)	309 / 554 (55.78%)	97 / 178 (54.49%)
occurrences (all)	512	525	148
Dyskinesia			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Dizziness			
subjects affected / exposed	1 / 549 (0.18%)	3 / 554 (0.54%)	1 / 178 (0.56%)
occurrences (all)	1	3	1
Taste disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Syncope			
subjects affected / exposed	2 / 549 (0.36%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	2	2	1
Speech disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Seizure			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Psychomotor hyperactivity			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Presyncope			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Neuromuscular blockade			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Somnolence			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Carpal tunnel syndrome			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Neuralgia			

subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Paraesthesia			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Restless legs syndrome			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Sleep deficit			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Tension headache			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Blood and lymphatic system disorders			
Anaemia			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Lymphadenopathy			
subjects affected / exposed	4 / 549 (0.73%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	4	2	1
Iron deficiency anaemia			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Lymphadenitis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Coagulopathy			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Increased tendency to bruise			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Ear and labyrinth disorders			
Tinnitus			

subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Ear discomfort			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Ear pain			
subjects affected / exposed	2 / 549 (0.36%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	2	3	0
Eustachian tube dysfunction			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Hypoacusis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Cerumen impaction			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Tympanic membrane perforation			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vertigo			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Excessive cerumen production			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
External ear inflammation			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Motion sickness			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Tympanosclerosis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Eye disorders			

Astigmatism			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Eye swelling			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Blindness transient			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Conjunctivitis allergic			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Eye irritation			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Eye pain			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Blepharitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Glaucoma			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Keratitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Ocular hyperaemia			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Swelling of eyelid			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vision blurred			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0

Conjunctival haemorrhage subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Chalazion subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 2	0 / 178 (0.00%) 0
Eye inflammation subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Eye pruritus subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Eyelid exfoliation subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Eyelid ptosis subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Myopia subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Gastrointestinal disorders			
Colitis subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Abdominal distension subjects affected / exposed occurrences (all)	2 / 549 (0.36%) 2	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Abdominal pain subjects affected / exposed occurrences (all)	6 / 549 (1.09%) 7	3 / 554 (0.54%) 3	1 / 178 (0.56%) 1
Abdominal pain lower subjects affected / exposed occurrences (all)	2 / 549 (0.36%) 2	0 / 554 (0.00%) 0	1 / 178 (0.56%) 1
Abdominal pain upper			

subjects affected / exposed	6 / 549 (1.09%)	3 / 554 (0.54%)	2 / 178 (1.12%)
occurrences (all)	6	4	2
Anal fissure			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Abdominal discomfort			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Constipation			
subjects affected / exposed	4 / 549 (0.73%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	4	0	0
Dental caries			
subjects affected / exposed	3 / 549 (0.55%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	4	0	0
Diarrhoea			
subjects affected / exposed	6 / 549 (1.09%)	4 / 554 (0.72%)	5 / 178 (2.81%)
occurrences (all)	6	4	5
Dyspepsia			
subjects affected / exposed	3 / 549 (0.55%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	3	0	1
Haemorrhoids			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Food poisoning			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	2	1	1
Gastritis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Gastrooesophageal reflux disease			
subjects affected / exposed	2 / 549 (0.36%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	2	3	0
Haematemesis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Enteritis			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Hyperchlorhydria			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Hypoaesthesia oral			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Irritable bowel syndrome			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Lip swelling			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Mouth ulceration			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Toothache			
subjects affected / exposed	3 / 549 (0.55%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	3	2	1
Oral pain			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Oral pruritus			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Pancreatitis relapsing			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Peptic ulcer			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Nausea			
subjects affected / exposed	121 / 549 (22.04%)	138 / 554 (24.91%)	45 / 178 (25.28%)
occurrences (all)	156	174	59
Vomiting			

subjects affected / exposed	2 / 549 (0.36%)	6 / 554 (1.08%)	1 / 178 (0.56%)
occurrences (all)	2	6	1
Palatal disorder			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Coeliac disease			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Embedded tooth			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Eructation			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Flatulence			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Gingival pain			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Aphthous ulcer			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Tongue discolouration			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Transient lingual papillitis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Hepatobiliary disorders			
Hepatic steatosis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Skin and subcutaneous tissue disorders			
Dermatitis contact			

subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
Dermatitis atopic			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0
Dermatitis allergic			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Dermatitis			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Cold urticaria			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Acne			
subjects affected / exposed	6 / 549 (1.09%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	7	2	1
Miliaria			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Pain of skin			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Pityriasis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Pruritus			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Rash			
subjects affected / exposed	7 / 549 (1.28%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	7	2	0
Rash macular			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Ingrowing nail			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Hyperhidrosis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Hand dermatitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Erythema			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Rash maculo-papular			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Eczema			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	3	1	0
Rash pruritic			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Skin lesion			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Urticaria			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	2	1	1
Rash erythematous			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Alopecia			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Dermatitis acneiform			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Drug eruption			

subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Nail disorder			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Psoriasis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Seborrhoeic dermatitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Skin induration			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Urticaria chronic			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Renal and urinary disorders			
Haematuria			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Dysuria			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Urinary tract inflammation			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Urinary retention			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Pollakiuria			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Nephrolithiasis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0

Nephritis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Hydronephrosis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Endocrine disorders			
Hypothyroidism			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Thyroid stimulating hormone deficiency			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Growth hormone deficiency			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Hyperthyroidism			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Thyroid cyst			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Musculoskeletal and connective tissue disorders			
Arthralgia			
subjects affected / exposed	84 / 549 (15.30%)	76 / 554 (13.72%)	25 / 178 (14.04%)
occurrences (all)	105	95	30
Arthritis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Axillary mass			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Back pain			
subjects affected / exposed	5 / 549 (0.91%)	8 / 554 (1.44%)	2 / 178 (1.12%)
occurrences (all)	5	8	2
Muscle swelling			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Knee deformity			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Joint warmth			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Joint swelling			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Joint hyperextension			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Muscular weakness			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Growing pains			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Foot deformity			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Flank pain			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Epiphysiolysis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Muscle tightness			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Intervertebral disc protrusion			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Spinal pain			

subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Sever's disease			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Rotator cuff syndrome			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Pain in extremity			
subjects affected / exposed	2 / 549 (0.36%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	2	2	1
Osteochondrosis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Neck pain			
subjects affected / exposed	3 / 549 (0.55%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	3	0	0
Neck mass			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Myositis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Myalgia			
subjects affected / exposed	122 / 549 (22.22%)	132 / 554 (23.83%)	30 / 178 (16.85%)
occurrences (all)	153	178	35
Musculoskeletal chest pain			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Temporomandibular joint syndrome			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Tendon pain			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Tenosynovitis			

subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Myokymia			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Bone pain			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Coccydynia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Costochondritis			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Joint effusion			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Pain in jaw			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Rhabdomyolysis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Sacral pain			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Scoliosis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Short stature			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Synovial cyst			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Tendonitis			

subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Torticollis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Spinal flattening			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Infections and infestations			
Acarodermatitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Acute sinusitis			
subjects affected / exposed	1 / 549 (0.18%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	1	2	0
Adenovirus infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Asymptomatic COVID-19			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
Bacterial vaginosis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Body tinea			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0
Bronchitis			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Bullous impetigo			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Cystitis			
subjects affected / exposed	1 / 549 (0.18%)	4 / 554 (0.72%)	1 / 178 (0.56%)
occurrences (all)	1	4	1

COVID-19 pneumonia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Cellulitis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Chlamydial infection			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Coronavirus infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
COVID-19			
subjects affected / exposed	63 / 549 (11.48%)	71 / 554 (12.82%)	26 / 178 (14.61%)
occurrences (all)	63	71	26
Ear infection			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	1 / 178 (0.56%)
occurrences (all)	0	2	1
Endometritis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Enterobiasis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Enterovirus infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Erythema migrans			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Genital herpes			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Eyelid infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0

Folliculitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Gastroenteritis			
subjects affected / exposed	5 / 549 (0.91%)	6 / 554 (1.08%)	0 / 178 (0.00%)
occurrences (all)	5	6	0
Gastroenteritis viral			
subjects affected / exposed	2 / 549 (0.36%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	2	3	0
Eye infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Infectious mononucleosis			
subjects affected / exposed	1 / 549 (0.18%)	2 / 554 (0.36%)	2 / 178 (1.12%)
occurrences (all)	1	2	2
Influenza			
subjects affected / exposed	10 / 549 (1.82%)	5 / 554 (0.90%)	1 / 178 (0.56%)
occurrences (all)	10	5	1
Localised infection			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0
Lower respiratory tract infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Lyme disease			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Infected bite			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Gingivitis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Helicobacter gastritis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0

Herpes zoster			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	1	0	1
Hordeolum			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Impetigo			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Mycoplasma genitalium infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Nasopharyngitis			
subjects affected / exposed	16 / 549 (2.91%)	20 / 554 (3.61%)	11 / 178 (6.18%)
occurrences (all)	17	21	11
Onychomycosis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Oral herpes			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Otitis externa			
subjects affected / exposed	4 / 549 (0.73%)	5 / 554 (0.90%)	0 / 178 (0.00%)
occurrences (all)	4	5	0
Otitis media			
subjects affected / exposed	4 / 549 (0.73%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	4	1	0
Pharyngitis			
subjects affected / exposed	10 / 549 (1.82%)	9 / 554 (1.62%)	2 / 178 (1.12%)
occurrences (all)	12	9	2
Otosalpingitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Paronychia			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0

Parotitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Pelvic inflammatory disease			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Otitis media acute			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	2	1	1
Pharyngitis streptococcal			
subjects affected / exposed	1 / 549 (0.18%)	4 / 554 (0.72%)	1 / 178 (0.56%)
occurrences (all)	1	4	1
Postoperative wound infection			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Pulpitis dental			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Respiratory tract infection			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
Respiratory tract infection bacterial			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Skin infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Rhinitis			
subjects affected / exposed	2 / 549 (0.36%)	2 / 554 (0.36%)	2 / 178 (1.12%)
occurrences (all)	2	2	2
Rhinovirus infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Sialoadenitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0

Sinusitis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	2 / 178 (1.12%)
occurrences (all)	1	0	2
Respiratory tract infection viral			
subjects affected / exposed	3 / 549 (0.55%)	4 / 554 (0.72%)	2 / 178 (1.12%)
occurrences (all)	4	4	2
Soft tissue infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Streptococcal infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Suspected COVID-19			
subjects affected / exposed	2 / 549 (0.36%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	2	3	0
Tinea versicolour			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Tonsillitis			
subjects affected / exposed	4 / 549 (0.73%)	2 / 554 (0.36%)	5 / 178 (2.81%)
occurrences (all)	5	3	6
Upper respiratory tract infection			
subjects affected / exposed	24 / 549 (4.37%)	22 / 554 (3.97%)	6 / 178 (3.37%)
occurrences (all)	25	23	6
Tooth abscess			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0
Tooth infection			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	1 / 178 (0.56%)
occurrences (all)	1	1	1
Tracheitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Trichomoniasis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1

Tonsillitis streptococcal			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Upper respiratory tract infection bacterial			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Urinary tract infection			
subjects affected / exposed	6 / 549 (1.09%)	4 / 554 (0.72%)	3 / 178 (1.69%)
occurrences (all)	10	4	3
Vaginal infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	2
Varicella			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Viral infection			
subjects affected / exposed	2 / 549 (0.36%)	4 / 554 (0.72%)	0 / 178 (0.00%)
occurrences (all)	2	5	0
Yersinia infection			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Viral upper respiratory tract infection			
subjects affected / exposed	6 / 549 (1.09%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	6	0	0
Vulvovaginal candidiasis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vulvovaginal mycotic infection			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Wound infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Viral pharyngitis			

subjects affected / exposed	3 / 549 (0.55%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	3	0	0
Conjunctivitis			
subjects affected / exposed	2 / 549 (0.36%)	3 / 554 (0.54%)	0 / 178 (0.00%)
occurrences (all)	2	3	0
Conjunctivitis viral			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Ear lobe infection			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Fungal foot infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Fungal skin infection			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Gastrointestinal viral infection			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Oral candidiasis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Helminthic infection			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Herpes simplex			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Laryngitis			
subjects affected / exposed	1 / 549 (0.18%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	1	1	0
Laryngotracheitis obstructive			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Molluscum contagiosum			

subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Hand-foot-and-mouth disease			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Subcutaneous abscess			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Tinea pedis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Vaccination site cellulitis			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vaccination site pustule			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Vulvovaginitis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Skin bacterial infection			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0
Parasitic gastroenteritis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Pericoronitis			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Pharyngotonsillitis			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Post-acute COVID-19 syndrome			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0
Respiratory syncytial virus infection			

subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Metabolism and nutrition disorders			
Gluten sensitivity			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	1 / 178 (0.56%)
occurrences (all)	0	0	1
Hypercholesterolaemia			
subjects affected / exposed	1 / 549 (0.18%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	1	0	0
Hyperlipidaemia			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Hyponatraemia			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Insulin resistance			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Iron deficiency			
subjects affected / exposed	2 / 549 (0.36%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	2	1	0
Folate deficiency			
subjects affected / exposed	2 / 549 (0.36%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	2	0	0
Abnormal loss of weight			
subjects affected / exposed	0 / 549 (0.00%)	1 / 554 (0.18%)	0 / 178 (0.00%)
occurrences (all)	0	1	0
Glucose tolerance impaired			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Lactose intolerance			
subjects affected / exposed	0 / 549 (0.00%)	0 / 554 (0.00%)	0 / 178 (0.00%)
occurrences (all)	0	0	0
Obesity			
subjects affected / exposed	0 / 549 (0.00%)	2 / 554 (0.36%)	0 / 178 (0.00%)
occurrences (all)	0	2	0

Type 2 diabetes mellitus subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Vitamin B complex deficiency subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Vitamin B12 deficiency subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Vitamin D deficiency subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Zinc deficiency subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Decreased appetite subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Haemochromatosis subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Hypertriglyceridaemia subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0
Hypoglycaemia subjects affected / exposed occurrences (all)	0 / 549 (0.00%) 0	1 / 554 (0.18%) 1	0 / 178 (0.00%) 0
Hypovitaminosis subjects affected / exposed occurrences (all)	1 / 549 (0.18%) 1	0 / 554 (0.00%) 0	0 / 178 (0.00%) 0

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
23 May 2019	As per the recommendation from CBER, the scope of the study has been extended to include the 3-dose (0,2,6-M) schedule and an additional 2-dose schedule (0,6-M) along with the 2-dose (0,2-M) schedule planned originally. The study will assess the immunogenicity of the 2-dose and 3-doses vaccination with rMenB+OMV NZ vaccine along with effectiveness and safety.
18 March 2020	The scope of this post-marketing commitment study has been extended to demonstrate the effectiveness, immunogenicity and safety of GSK's investigational combined meningococcal ABCWY vaccine (from a phase III MenABCWY study) along with the rMenB+OMV NZ vaccine.
23 September 2020	This protocol is amended primarily as a consequence of feedback from regulatory authorities of participating countries following their review of Protocol Amendment 2. Additional changes have been made to improve the clarity of the text.
09 May 2021	The protocol is being amended to document the increase in blood volumes drawn at certain visits (Visit 2 and Visit 6). The allowed windows for study visits during special circumstances have also been widened to maintain subject visit compliance during the COVID-19 pandemic. Additionally, considering that some of the study interventions are combination products constituted of a device and biologic product (pre-filled syringes), the amended protocol provides instructions for collection of safety information related to the use of medical devices. The reporting period for pregnancies has also been updated in line with the current guidelines

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported