

Secondary Efficacy Endpoint: Percent Change in Lipid Parameter Lp(a) from Baseline at Week 24/LOCF (Full Analysis Set)

	Observed Value	Change from Baseline	Percent Change from Baseline	Fisher Combined p-value^a
Lp(a) (mg/dL)				0.0021
n	15	14	14	
Mean (SD)	32.959 (31.2902)	-10.553 (25.8881)	0.877 (46.9887)	
Median	30.00	0.000	0.000	
Range	2.00 to 116.00	-81.00 to 19.00	-67.77 to 118.75	
(95% CI) ^b			(-26.2535, 28.0075)	
Lp(a) (nmol/L)				
n	28	27	27	
Mean (SD)	82.493 (75.5080)	-47.383 (59.2811)	-26.757 (32.5675)	
Median	54.300	-30.000	-37.947	
Range	7.00 to 294.20	-222.00 to 29.00	-71.01 to 56.75	
(95% CI) ^b			(-39.6398, -13.8733)	

Abbreviations: CI=confidence interval; LOCF=last observation carried forward; Lp(a)=lipoprotein(a); N=number of subjects in the analysis set; n=number of subjects taking a concomitant medication; SD=standard deviation.

^a Individual p-values were 0.9454 and 0.0002 for data reported in mg/dL and nmol/L, respectively.

^b 95% CI for mean percentage change.