



Clinical trial results:

A Phase IIb, randomized, partially blind, active controlled, dose-range finding study of GSK3640254 compared to a reference arm of dolutegravir, each in combination with nucleoside reverse transcriptase inhibitors, in HIV-1 infected antiretroviral treatment-naïve adults

Summary

EudraCT number	2019-004435-23
Trial protocol	PT FR DE IT
Global end of trial date	29 May 2023

Results information

Result version number	v1
This version publication date	21 September 2023
First version publication date	21 September 2023

Trial information

Trial identification

Sponsor protocol code	208379
-----------------------	--------

Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT04493216
WHO universal trial number (UTN)	-

Notes:

Sponsors

Sponsor organisation name	ViiV Healthcare
Sponsor organisation address	980 GreatWest Road, Brentford, Middlesex, United Kingdom, TW8 9GS
Public contact	GSK Response Center, ViiV Healthcare, 1 8664357343, GSKClinicalSupportHD@gsk.com
Scientific contact	GSK Response Center, ViiV Healthcare, 1 8664357343, GSKClinicalSupportHD@gsk.com

Notes:

Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

Results analysis stage

Analysis stage	Interim
Date of interim/final analysis	21 October 2022
Is this the analysis of the primary completion data?	Yes
Primary completion date	05 September 2022
Global end of trial reached?	Yes
Global end of trial date	29 May 2023
Was the trial ended prematurely?	No

Notes:

General information about the trial

Main objective of the trial:

To evaluate antiviral efficacy of GSK3640254 relative to DTG, each given in combination with 2 NRTIs, enabling the selection of an optimal dose for GSK3640254

Protection of trial subjects:

Not Applicable

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	18 November 2020
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

Population of trial subjects

Subjects enrolled per country

Country: Number of subjects enrolled	Argentina: 36
Country: Number of subjects enrolled	Canada: 4
Country: Number of subjects enrolled	France: 2
Country: Number of subjects enrolled	Germany: 6
Country: Number of subjects enrolled	Italy: 11
Country: Number of subjects enrolled	Portugal: 8
Country: Number of subjects enrolled	Russian Federation: 20
Country: Number of subjects enrolled	South Africa: 16
Country: Number of subjects enrolled	Spain: 41
Country: Number of subjects enrolled	United States: 17
Worldwide total number of subjects	161
EEA total number of subjects	68

Notes:

Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23	0

months)	
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	160
From 65 to 84 years	1
85 years and over	0

Subject disposition

Recruitment

Recruitment details: -

Pre-assignment

Screening details:

The results presented are based on the data up to a maximum of Week 48 (Partially Blind Randomised Phase). Data collection is still ongoing and additional results will be provided after study completion.

Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator

Arms

Are arms mutually exclusive?	Yes
Arm title	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF

Arm description:

Participants with human immunodeficiency virus type 1 (HIV-1), orally received one 100 mg tablet per day of GSK3640254 and two tablets per day of matching placebo in a blinded setting. Open label dual nucleoside reverse transcriptase inhibitors (NRTIs) background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.

Arm type	Experimental
Investigational medicinal product name	2 NRTIs
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Open label dual NRTI background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.

Investigational medicinal product name	100 mg GSK3640254
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants with human immunodeficiency virus type 1 (HIV-1) infected antiretroviral treatment-naïve adults orally received one 100 mg tablet per day of GSK3640254 and two tablets per day of matching placebo as blinded.

Arm title	GSK3640254 150 mg + ABC/3TC or FTC/TAF
------------------	--

Arm description:

Participants with HIV-1, orally received 150 mg (one 100 mg tablet + two 25 mg tablets per day) of GSK3640254 in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Arm type	Experimental
----------	--------------

Investigational medicinal product name	2 NRTIs
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Open label dual NRTI background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.

Investigational medicinal product name	150 mg GSK3640254
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants with HIV-1 infected antiretroviral treatment-naïve adults orally received 150 mg (one 100 mg tablet + two 25 mg tablets per day) of GSK3640254 as blinded.

Arm title	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
------------------	--

Arm description:

Participants with HIV-1, orally received two 100 mg tablets (200 mg) per day of GSK3640254 and one tablet per day of matching placebo in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Arm type	Experimental
Investigational medicinal product name	2 NRTIs
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Open label dual NRTI background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.

Investigational medicinal product name	200 mg GSK3640254
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants with HIV-1 infected antiretroviral treatment-naïve adults orally received two 100 mg tablets (200 mg) per day of GSK3640254 and one tablet per day of matching placebo as blinded.

Arm title	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
------------------	---

Arm description:

Participants with HIV-1, orally received one 50 mg tablet per day of DTG in an open label setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Arm type	Active comparator
Investigational medicinal product name	2 NRTIs
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Open label dual NRTI background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.

Investigational medicinal product name	50 mg DTG
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Tablet
Routes of administration	Oral use

Dosage and administration details:

Participants with HIV-1 infected antiretroviral treatment-naïve adults orally received one 50 mg tablet per day of DTG in an open label setting.

Number of subjects in period 1	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
Started	40	43	42
Completed	0	0	0
Not completed	40	43	42
Consent withdrawn by subject	1	1	1
Physician decision	-	-	1
Adverse event, non-fatal	1	4	5
Protocol Deviation	1	2	-
Ongoing	18	18	16
Study Terminated by Sponsor	14	15	11
Lost to follow-up	1	1	-
Subject Reached Protocol-Defined Stopping Criteria	4	2	8

Number of subjects in period 1	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Started	36
Completed	0
Not completed	36
Consent withdrawn by subject	2
Physician decision	-
Adverse event, non-fatal	2
Protocol Deviation	2
Ongoing	16
Study Terminated by Sponsor	11
Lost to follow-up	1
Subject Reached Protocol-Defined Stopping Criteria	2

Baseline characteristics

Reporting groups

Reporting group title	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF
-----------------------	---

Reporting group description:

Participants with human immunodeficiency virus type 1 (HIV-1), orally received one 100 mg tablet per day of GSK3640254 and two tablets per day of matching placebo in a blinded setting. Open label dual nucleoside reverse transcriptase inhibitors (NRTIs) background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.

Reporting group title	GSK3640254 150 mg + ABC/3TC or FTC/TAF
-----------------------	--

Reporting group description:

Participants with HIV-1, orally received 150 mg (one 100 mg tablet + two 25 mg tablets per day) of GSK3640254 in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Reporting group title	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
-----------------------	--

Reporting group description:

Participants with HIV-1, orally received two 100 mg tablets (200 mg) per day of GSK3640254 and one tablet per day of matching placebo in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Reporting group title	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
-----------------------	---

Reporting group description:

Participants with HIV-1, orally received one 50 mg tablet per day of DTG in an open label setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Reporting group values	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
Number of subjects	40	43	42
Age categorical			
Units: Subjects			
19-64 years	40	42	42
>=65 years	0	1	0
Sex: Female, Male			
Units: Participants			
Female	7	9	12
Male	33	34	30
Race/Ethnicity, Customized			
Units: Subjects			
AMERICAN INDIAN OR ALASKA NATIVE	1	2	1
ASIAN	0	2	2
BLACK OR AFRICAN AMERICAN	6	8	6
WHITE	32	31	32
MIXED RACE	1	0	0
MISSING	0	0	1
Age, Continuous			
Units: YEARS			
arithmetic mean	32.8	38.1	33.7
standard deviation	± 6.20	± 12.54	± 10.59

Reporting group values	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF	Total	
Number of subjects	36	161	
Age categorical Units: Subjects			
19-64 years	36	160	
>=65 years	0	1	
Sex: Female, Male Units: Participants			
Female	10	38	
Male	26	123	
Race/Ethnicity, Customized Units: Subjects			
AMERICAN INDIAN OR ALASKA NATIVE	0	4	
ASIAN	1	5	
BLACK OR AFRICAN AMERICAN	6	26	
WHITE	29	124	
MIXED RACE	0	1	
MISSING	0	1	
Age, Continuous Units: YEARS			
arithmetic mean	35.3		
standard deviation	± 9.85	-	

End points

End points reporting groups

Reporting group title	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF
Reporting group description: Participants with human immunodeficiency virus type 1 (HIV-1), orally received one 100 mg tablet per day of GSK3640254 and two tablets per day of matching placebo in a blinded setting. Open label dual nucleoside reverse transcriptase inhibitors (NRTIs) background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.	
Reporting group title	GSK3640254 150 mg + ABC/3TC or FTC/TAF
Reporting group description: Participants with HIV-1, orally received 150 mg (one 100 mg tablet + two 25 mg tablets per day) of GSK3640254 in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.	
Reporting group title	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
Reporting group description: Participants with HIV-1, orally received two 100 mg tablets (200 mg) per day of GSK3640254 and one tablet per day of matching placebo in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.	
Reporting group title	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Reporting group description: Participants with HIV-1, orally received one 50 mg tablet per day of DTG in an open label setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.	

Primary: Percentage of participants with plasma HIV-1 ribonucleic acid (RNA) less than (<)50 copies per milliliter (c/mL) at Week 24

End point title	Percentage of participants with plasma HIV-1 ribonucleic acid (RNA) less than (<)50 copies per milliliter (c/mL) at Week 24
End point description: Percentage of participants with plasma HIV-1 RNA <50 c/mL at week 24 was assessed using the Food and Drug Administration (FDA) snapshot algorithm to demonstrate the antiviral activity of GSK3640254 given in combination with either ABC/3TC or FTC/TAF compared to the reference treatment of DTG+ABC/3TC or DTG+FTC/TAF. The other snapshot outcomes (HIV-1 RNA ≥50 c/mL and no virologic data at Week 24 Window) were classified as non-responders. Intent-to-Treat Exposed (ITT-E) population included all randomized participants who received at least one dose of study intervention.	
End point type	Primary
End point timeframe: At Week 24	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Percentage of participants				

number (confidence interval 95%)	82.5 (68.1 to 91.3)	90.7 (78.4 to 96.3)	76.2 (61.5 to 86.5)	91.7 (78.2 to 97.1)
----------------------------------	---------------------	---------------------	---------------------	---------------------

Statistical analyses

Statistical analysis title	Statistical Analysis 1
Statistical analysis description: Differences in percentage of participant = Percentage of participant in GSK3640254 100 mg+2NRTIs - Percentage of participant in DTG+2NRTIs	
Comparison groups	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF v Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Number of subjects included in analysis	76
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Differences in percentage of participant
Point estimate	-9.2
Confidence interval	
level	95 %
sides	2-sided
lower limit	-24
upper limit	5.7

Statistical analysis title	Statistical Analysis 3
Statistical analysis description: Differences in percentage of participant = Percentage of participant in GSK3640254 200 mg+2NRTIs - Percentage of participant in DTG+2NRTIs	
Comparison groups	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF v Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Number of subjects included in analysis	78
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Differences in percentage of participant
Point estimate	-15.5
Confidence interval	
level	95 %
sides	2-sided
lower limit	-31.2
upper limit	0.3

Statistical analysis title	Statistical Analysis 2
Statistical analysis description: Differences in percentage of participant = Percentage of participant in GSK3640254 150 mg+2NRTIs - Percentage of participant in DTG+2NRTIs	
Comparison groups	GSK3640254 150 mg + ABC/3TC or FTC/TAF v Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF

Number of subjects included in analysis	79
Analysis specification	Pre-specified
Analysis type	other
Parameter estimate	Differences in percentage of participant
Point estimate	-1
Confidence interval	
level	95 %
sides	2-sided
lower limit	-13.5
upper limit	11.6

Secondary: Percentage of participants with plasma HIV-1 RNA <50 c/mL at Week 48

End point title	Percentage of participants with plasma HIV-1 RNA <50 c/mL at Week 48
-----------------	--

End point description:

Percentage of participants with plasma HIV-1 RNA <50 c/mL at week 48 was assessed using the FDA snapshot algorithm to demonstrate the antiviral activity of GSK3640254 given in combination with either ABC/3TC or FTC/TAF compared to the reference treatment of DTG+ABC/3TC or DTG+FTC/TAF. The other snapshot outcomes (HIV-1 RNA ≥50 c/mL and no virologic data at Week 48 Window) were classified as non-responders. Intent-to-Treat Exposed (ITT-E) population.

End point type	Secondary
----------------	-----------

End point timeframe:

At Week 48

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Percentage of participants				
number (confidence interval 95%)	85.0 (70.9 to 92.9)	83.7 (70.0 to 91.9)	76.2 (61.5 to 86.5)	77.8 (61.9 to 88.3)

Statistical analyses

No statistical analyses for this end point

Secondary: Absolute values of HIV-1 RNA at Weeks 24 and 48

End point title	Absolute values of HIV-1 RNA at Weeks 24 and 48
-----------------	---

End point description:

Plasma samples were collected for quantitative analysis of HIV-1 RNA. Logarithm to base 10 (log10) values for plasma HIV-1 RNA have been presented. Baseline was defined as the latest pre-dose assessment with a non-missing value, including those from unscheduled visits. Intent-to-Treat Exposed (ITT-E) population.

End point type	Secondary
End point timeframe:	
Baseline (Day 1) and at Weeks 24 and 48	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Log 10 copies per milliliter				
arithmetic mean (standard deviation)				
Baseline (Day 1)	4.351 (± 0.5712)	4.353 (± 0.6705)	4.165 (± 0.6505)	4.247 (± 0.6765)
Week 24	1.619 (± 0.1260)	1.607 (± 0.0663)	1.610 (± 0.0679)	1.592 (± 0.0126)
Week 48	1.602 (± 0.0536)	1.605 (± 0.0647)	1.594 (± 0.0233)	1.590 (± 0.0000)

Statistical analyses

No statistical analyses for this end point

Secondary: Change from Baseline in plasma HIV-1 RNA at Weeks 24 and 48

End point title	Change from Baseline in plasma HIV-1 RNA at Weeks 24 and 48
End point description:	
Plasma samples were collected for quantitative analysis of HIV-1 RNA. log10 values for plasma HIV-1 RNA have been presented. Baseline was defined as the latest pre-dose assessment with a non-missing value, including those from unscheduled visits. Change from Baseline is defined as post-dose visit value minus Baseline value. Intent-to-Treat Exposed (ITT-E) population.	
End point type	Secondary
End point timeframe:	
Baseline (Day 1) and at Weeks 24 and 48 - ITT-E population	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Log 10 copies per milliliter				
arithmetic mean (standard deviation)				
Baseline (Day 1)	4.351 (± 0.5712)	4.353 (± 0.6705)	4.165 (± 0.6505)	4.247 (± 0.6765)

Week 24	-2.718 (± 0.5501)	-2.784 (± 0.6615)	-2.565 (± 0.6513)	-2.629 (± 0.6835)
Week 48	-2.675 (± 0.5640)	-2.762 (± 0.6784)	-2.580 (± 0.6941)	-2.717 (± 0.6672)

Statistical analyses

No statistical analyses for this end point

Secondary: Absolute values of Cluster of differentiation 4 plus (CD4+) cell counts at Weeks 24 and 48

End point title	Absolute values of Cluster of differentiation 4 plus (CD4+) cell counts at Weeks 24 and 48
-----------------	--

End point description:

Blood samples were collected and CD4+ cell count was assessed using flow cytometry. Baseline was defined as the latest pre-dose assessment with a non-missing value, including those from unscheduled visits. Intent-to-Treat Exposed (ITT-E) population.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Weeks 24 and 48

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Cells per cubic millimeter				
arithmetic mean (standard deviation)				
Baseline (Day 1)	480.3 (± 171.72)	509.7 (± 207.13)	478.7 (± 204.04)	514.1 (± 240.88)
Week 24	717.5 (± 222.91)	643.5 (± 202.27)	689.8 (± 316.64)	724.8 (± 403.99)
Week 48	749.9 (± 328.28)	702.6 (± 258.65)	747.3 (± 313.15)	705.2 (± 221.18)

Statistical analyses

No statistical analyses for this end point

Secondary: Change from Baseline in CD4+ cell counts at Weeks 24 and 48

End point title	Change from Baseline in CD4+ cell counts at Weeks 24 and 48
-----------------	---

End point description:

Blood samples were collected and CD4+ cell count was assessed using flow cytometry. Baseline was defined as the latest pre-dose assessment with a non-missing value, including those from unscheduled visits. Change from Baseline is defined as post-dose visit value minus Baseline value. Intent-to-Treat Exposed (ITT-E) population.

End point type	Secondary
End point timeframe:	
Baseline (Day 1) and at Weeks 24 and 48	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Cells per cubic millimeter				
arithmetic mean (standard deviation)				
Baseline (Day 1)	480.3 (± 171.72)	509.7 (± 207.13)	478.7 (± 204.04)	514.1 (± 240.88)
Week 24	241.3 (± 191.26)	129.3 (± 233.07)	202.3 (± 271.98)	198.5 (± 285.00)
Week 48	292.4 (± 264.16)	189.2 (± 216.10)	243.0 (± 233.24)	190.6 (± 180.19)

Statistical analyses

No statistical analyses for this end point

Secondary: Number of participants with serious adverse events (SAEs) and Deaths through Weeks 24 and 48

End point title	Number of participants with serious adverse events (SAEs) and Deaths through Weeks 24 and 48
-----------------	--

End point description:

An SAE is defined as any serious adverse event that, at any dose results in death, is life threatening, requires inpatient hospitalization or prolongation of existing hospitalization, results in persistent or significant disability/incapacity, is a congenital anomaly/birth defect, or any other situation according to medical or scientific judgment. Safety population included all randomized participants who were exposed to study intervention with the exception of any participants with documented evidence of not having consumed any amount of study intervention.

End point type	Secondary
End point timeframe:	
Up to Weeks 24 and 48	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Participants				

Week 24, SAEs	2	4	2	2
Week 48, SAEs	2	7	2	2
Week 24, Deaths	0	0	0	0
Week 48, Deaths	0	0	0	0

Statistical analyses

No statistical analyses for this end point

Secondary: Number of participants with adverse events (AEs) leading to treatment discontinuation through Weeks 24 and 48

End point title	Number of participants with adverse events (AEs) leading to treatment discontinuation through Weeks 24 and 48
-----------------	---

End point description:

An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of a study intervention, whether or not considered related to the study intervention. Number of participants who discontinued study treatment due to AEs are presented. Safety population.

End point type	Secondary
----------------	-----------

End point timeframe:

Up to Weeks 24 and 48

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Participants				
Week 24	2	3	4	2
Week 48	2	4	5	2

Statistical analyses

No statistical analyses for this end point

Secondary: Number of participants by severity of AEs through Weeks 24 and 48

End point title	Number of participants by severity of AEs through Weeks 24 and 48
-----------------	---

End point description:

An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of a study intervention, whether or not considered related to the study intervention. The severity of AEs was defined as per the Division of AIDS Table for Grading the Severity of Adult and Pediatric Adverse Events (DAIDS AE Grading Table) Version 2.1 and was categorized into grades as following: Grade 1 - mild, Grade 2 - moderate, Grade 3 - severe, Grade 4 - Potentially life threatening and Grade 5 - Fatal. Higher grade indicates more severe condition. There were 0 participants who experienced grade 5 severity up to weeks 24 and 48. Safety population.

End point type	Secondary
End point timeframe:	
Up to Weeks 24 and 48	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Participants				
Weeks 24, Grade 1	14	18	16	15
Weeks 48, Grade 1	12	14	17	15
Weeks 24, Grade 2	16	16	18	10
Weeks 48, Grade 2	20	17	17	10
Weeks 24, Grade 3	4	3	2	3
Weeks 48, Grade 3	4	5	3	3
Weeks 24, Grade 4	1	1	1	0
Weeks 48, Grade 4	1	2	1	0
Weeks 24, Grade 5	0	0	0	0
Weeks 48, Grade 5	0	0	0	0

Statistical analyses

No statistical analyses for this end point

Secondary: Number of participants with AEs in gastrointestinal (GI), nervous system, and psychiatric conditions through Weeks 24 and 48

End point title	Number of participants with AEs in gastrointestinal (GI), nervous system, and psychiatric conditions through Weeks 24 and 48
-----------------	--

End point description:

An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of a study intervention, whether or not considered related to the study intervention. Number of participants with AEs in gastrointestinal (GI), nervous system, and psychiatric conditions are presented. Safety population.

End point type	Secondary
End point timeframe:	
At Weeks 24 and 48	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	40	43	42	36
Units: Participants				
Week 24, Gastrointestinal disorders	13	18	12	11
Week 48, Gastrointestinal disorders	14	18	14	14
Week 24, Nervous system disorders	5	4	7	3
Week 48, Nervous system disorders	7	5	8	3
Week 24, Psychiatric disorders	1	3	2	2
Week 48, Psychiatric disorders	4	3	4	4

Statistical analyses

No statistical analyses for this end point

Secondary: Number of participants with genotypic resistance

End point title	Number of participants with genotypic resistance
End point description:	
Plasma samples were collected for resistance testing. Genotypic testing was conducted in participants meeting protocol-defined virologic failure (PVSF) criteria has been presented. Major resistance mutations to other classes (NRTI, non-NRTI and protease inhibitor (PI)) were defined by the International Antiviral Society-USA (IAS-USA). Baseline was defined as the latest pre-dose assessment with a non-missing value, including those from unscheduled visits. No participants were reported to have genotypic resistance through weeks 24 and 48. PVSF population included participant with A. virologic non-response (Decrease from Baseline in plasma HIV-1 RNA of 1.0 log ₁₀ c/mL unless plasma HIV-1 RNA is <200 c/mL by Week 12; confirmed plasma HIV-1 RNA levels ≥200 c/mL at or after Week 24; plasma HIV-1 RNA ≥50 c/mL on repeat testing of Week 24 results and prior to Week 28) and B. virologic rebound (confirmed plasma HIV-1 RNA ≥200 c/mL after confirmed consecutive plasma HIV-1 RNA <50 c/mL).	
End point type	Secondary
End point timeframe:	
Baseline (Day 1) and at Weeks 24 and 48	

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	3	2	5	1
Units: Participants				
Baseline (Day 1)	0	0	0	0
Weeks 24	0	0	0	0
Weeks 48	0	0	0	0

Statistical analyses

No statistical analyses for this end point

Secondary: Number of participants with phenotypic resistance

End point title	Number of participants with phenotypic resistance
-----------------	---

End point description:

Plasma samples were collected for resistance testing. Phenotypic testing was conducted in participants meeting PDVF criteria has been presented. Major resistance mutations to other classes (nucleoside reverse transcriptase inhibitor (NRTI), non-nucleoside reverse transcriptase inhibitors (NNRTI) and protease inhibitor (PI)) were defined by the International Antiviral Society-USA (IAS-USA). Baseline was defined as the latest pre-dose assessment with a non-missing value, including those from unscheduled visits. No participants were reported to have phenotypic resistance at week 48. Protocol-defined virologic failure (PDVF) population. Only those participants with data available at specified time points have been analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline (Day 1) and at Weeks 24 and 48

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	3	2	5	1
Units: Participants				
Baseline (Day 1), NNRTI	0	0	0	0
Weeks 24, NNRTI	0	0	1	0
Weeks 48, NNRTI	0	0	0	0

Statistical analyses

No statistical analyses for this end point

Secondary: Observed plasma concentration at the end of the dosing interval (Ctau) of GSK3640254 at steady state

End point title	Observed plasma concentration at the end of the dosing interval (Ctau) of GSK3640254 at steady state ^[1]
-----------------	---

End point description:

Blood samples were collected at indicated time points for pharmacokinetic (PK) analysis of GSK3640254. Observed plasma concentration at the end of the dosing interval was determined directly from the concentration-time data. Intensive pharmacokinetic (PK) population included all participants who

received at least one dose of GSK3640254 and have evaluable drug concentrations reported, where samples are collected according to the intensive PK sampling scheme. Only those participants who received GSK3640254 have been analyzed. Only those participants with data available at specified time points have been analyzed.

End point type	Secondary
End point timeframe:	
At Weeks 2, 24 and 48	

Notes:

[1] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Per objective of this endpoint, only analysis of GSK3640254 was planned to be presented

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	14	13	18	
Units: Nanogram/ milliliter (ng/mL)				
geometric mean (geometric coefficient of variation)				
Week 2	430.5361 (± 56.5)	604.8387 (± 44.2)	805.2209 (± 43.7)	
Week 24	396.5117 (± 62.5)	519.6041 (± 84.9)	922.9638 (± 51.6)	
Week 48	310.9877 (± 135.4)	568.0656 (± 13.4)	812.4745 (± 49.1)	

Statistical analyses

No statistical analyses for this end point

Secondary: Area under the plasma drug concentration-time curve from pre-dose to the end of the dosing interval (AUC [0-tau]) of GSK3640254 at steady state

End point title	Area under the plasma drug concentration-time curve from pre-dose to the end of the dosing interval (AUC [0-tau]) of GSK3640254 at steady state ^[2]
-----------------	--

End point description:

Blood samples were collected at indicated time points for PK analysis of GSK3640254. Intensive pharmacokinetic (PK) population. Only those participants who received GSK3640254 have been analyzed.

End point type	Secondary
End point timeframe:	
At Week 2	

Notes:

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Per objective of this endpoint, only analysis of GSK3640254 was planned to be presented

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	14	13	18	
Units: Hour*nanogram/ milliliter (h*ng/mL)				
geometric mean (geometric coefficient of variation)	14995.7576 (± 49.8)	21212.2480 (± 44.5)	30708.2546 (± 40.1)	

Statistical analyses

No statistical analyses for this end point

Secondary: Maximum observed concentration (Cmax) of GSK3640254 at steady state

End point title	Maximum observed concentration (Cmax) of GSK3640254 at steady state ^[3]
-----------------	--

End point description:

Blood samples were collected at indicated time points for PK analysis of GSK3640254. Intensive pharmacokinetic (PK) population. Only those participants who received GSK3640254 have been analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

At Week 2

Notes:

[3] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: Per objective of this endpoint, only analysis of GSK3640254 was planned to be presented

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	15	13	18	
Units: ng/mL				
geometric mean (geometric coefficient of variation)	929.8 (± 45.9)	1337.3 (± 47.7)	2094.5 (± 39.2)	

Statistical analyses

No statistical analyses for this end point

Secondary: Observed pre-dose plasma concentration (C0) of GSK3640254 at steady state

End point title	Observed pre-dose plasma concentration (C0) of GSK3640254
-----------------	---

End point description:

Blood samples were collected at indicated time points for PK analysis of GSK3640254. Observed pre-dose plasma concentration was determined directly from the concentration-time data. Intensive pharmacokinetic (PK) population. Only those participants who received GSK3640254 have been analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

At Week 2

Notes:

[4] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: Per objective of this endpoint, only analysis of GSK3640254 was planned to be presented

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	15	13	18	
Units: ng/mL				
geometric mean (geometric coefficient of variation)	435.9 (± 54.6)	603.2 (± 59.5)	865.2 (± 43.4)	

Statistical analyses

No statistical analyses for this end point

Secondary: Time to Cmax (Tmax) of GSK3640254 at steady state

End point title	Time to Cmax (Tmax) of GSK3640254 at steady state ^[5]
-----------------	--

End point description:

Blood samples were collected at indicated time points for PK analysis of GSK3640254. Tmax was determined directly from the concentration-time data. Intensive pharmacokinetic (PK) population. Only those participants who received GSK3640254 have been analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

At Week 2

Notes:

[5] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period. Justification: Per objective of this endpoint, only analysis of GSK3640254 was planned to be presented

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	15	13	18	
Units: Hour				
median (full range (min-max))	3.0000 (1.900 to 9.017)	3.4167 (1.000 to 9.000)	3.4583 (1.917 to 5.000)	

Statistical analyses

No statistical analyses for this end point

Secondary: Oral clearance (CL_{tau}/F) of GSK3640254 at steady state

End point title	Oral clearance (CL _{tau} /F) of GSK3640254 at steady state ^[6]
-----------------	--

End point description:

Blood samples were collected at indicated time points for PK analysis of GSK3640254. CL_{tau}/F was calculated as the dose of GSK3640254 divided by AUC [0-tau]. Intensive pharmacokinetic (PK) population. Only those participants who received GSK3640254 have been analyzed.

End point type	Secondary
----------------	-----------

End point timeframe:

At Week 2

Notes:

[6] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.
Justification: Per objective of this endpoint, only analysis of GSK3640254 was planned to be presented

End point values	GSK3640254 100 milligram(mg) + Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF	
Subject group type	Reporting group	Reporting group	Reporting group	
Number of subjects analysed	14	13	18	
Units: Liter/ hour				
geometric mean (geometric coefficient of variation)	6.6686 (± 49.8)	7.0714 (± 44.5)	6.5129 (± 40.1)	

Statistical analyses

No statistical analyses for this end point

Adverse events

Adverse events information

Timeframe for reporting adverse events:

All-cause mortality, non-serious adverse events (Non-SAEs) and serious adverse events (SAEs) were collected maximum up to 48 weeks

Adverse event reporting additional description:

Safety population included all randomized participants who were exposed to study intervention with the exception of any participants with documented evidence of not having consumed any amount of study intervention.

Assessment type	Systematic
-----------------	------------

Dictionary used

Dictionary name	V25.0
-----------------	-------

Dictionary version	25.0
--------------------	------

Reporting groups

Reporting group title	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF
-----------------------	---

Reporting group description:

Participants with human immunodeficiency virus type 1 (HIV-1), orally received one 100 mg tablet per day of GSK3640254 and two tablets per day of matching placebo in a blinded setting. Open label dual nucleoside reverse transcriptase inhibitors (NRTIs) background therapies were given as one tablet per day of 600 mg abacavir (ABC) / 300 mg lamivudine (3TC) OR 200 mg emtricitabine (FTC) / 25 mg tenofovir alafenamide (TAF) orally.

Reporting group title	GSK3640254 150 mg + ABC/3TC or FTC/TAF
-----------------------	--

Reporting group description:

Participants with HIV-1, orally received one 50 mg tablet per day of DTG in an open label setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Reporting group title	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
-----------------------	--

Reporting group description:

Participants with HIV-1, orally received two 100 mg tablets (200 mg) per day of GSK3640254 and one tablet per day of matching placebo in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Reporting group title	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF
-----------------------	---

Reporting group description:

Participants with HIV-1, orally received 150 mg (one 100 mg tablet + two 25 mg tablets per day) of GSK3640254 in a blinded setting. Open label dual NRTI background therapies were given as one tablet per day combination of 600 mg ABC / 300 mg 3TC OR 200 mg FTC / 25 mg TAF orally.

Serious adverse events	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
Total subjects affected by serious adverse events			
subjects affected / exposed	2 / 40 (5.00%)	2 / 36 (5.56%)	2 / 42 (4.76%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Anogenital warts			

subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Injury, poisoning and procedural complications			
Ankle fracture			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Limb fracture			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pregnancy, puerperium and perinatal conditions			
Abortion spontaneous			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	1 / 42 (2.38%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Gastrointestinal disorders			
Colitis			
subjects affected / exposed	0 / 40 (0.00%)	1 / 36 (2.78%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pancreatitis			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	1 / 42 (2.38%)
occurrences causally related to treatment / all	0 / 0	0 / 0	1 / 1
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Hepatobiliary disorders			
Cholecystitis acute			
subjects affected / exposed	1 / 40 (2.50%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Respiratory, thoracic and mediastinal disorders			
Pulmonary hypertension			

subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Pulmonary embolism			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Skin and subcutaneous tissue disorders			
Dermatitis allergic			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Psychiatric disorders			
Suicidal ideation			
subjects affected / exposed	1 / 40 (2.50%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Substance dependence			
subjects affected / exposed	0 / 40 (0.00%)	1 / 36 (2.78%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Infections and infestations			
Appendicitis			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Cellulitis			
subjects affected / exposed	0 / 40 (0.00%)	1 / 36 (2.78%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0
Necrotising fasciitis			
subjects affected / exposed	0 / 40 (0.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences causally related to treatment / all	0 / 0	0 / 0	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0	0 / 0

Serious adverse events	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF		
Total subjects affected by serious adverse events			
subjects affected / exposed	7 / 43 (16.28%)		
number of deaths (all causes)	0		
number of deaths resulting from adverse events			
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Anogenital warts			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Injury, poisoning and procedural complications			
Ankle fracture			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Limb fracture			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Pregnancy, puerperium and perinatal conditions			
Abortion spontaneous			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Gastrointestinal disorders			
Colitis			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Pancreatitis			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		

Hepatobiliary disorders			
Cholecystitis acute			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Respiratory, thoracic and mediastinal disorders			
Pulmonary hypertension			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Pulmonary embolism			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Skin and subcutaneous tissue disorders			
Dermatitis allergic			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	1 / 1		
deaths causally related to treatment / all	0 / 0		
Psychiatric disorders			
Suicidal ideation			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Substance dependence			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Infections and infestations			
Appendicitis			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		
Cellulitis			

subjects affected / exposed	0 / 43 (0.00%)		
occurrences causally related to treatment / all	0 / 0		
deaths causally related to treatment / all	0 / 0		
Necrotising fasciitis			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences causally related to treatment / all	0 / 1		
deaths causally related to treatment / all	0 / 0		

Frequency threshold for reporting non-serious adverse events: 2 %

Non-serious adverse events	GSK3640254 100 milligram(mg)+ Placebo+ ABC/3TC or FTC/TAF	GSK3640254 150 mg + ABC/3TC or FTC/TAF	GSK3640254 200 mg+ Placebo+ ABC/3TC or FTC/TAF
Total subjects affected by non-serious adverse events			
subjects affected / exposed	37 / 40 (92.50%)	27 / 36 (75.00%)	38 / 42 (90.48%)
Investigations			
Blood creatine phosphokinase increased			
subjects affected / exposed	3 / 40 (7.50%)	1 / 36 (2.78%)	1 / 42 (2.38%)
occurrences (all)	3	1	1
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Anogenital warts			
subjects affected / exposed	1 / 40 (2.50%)	0 / 36 (0.00%)	1 / 42 (2.38%)
occurrences (all)	1	0	1
Injury, poisoning and procedural complications			
Limb injury			
subjects affected / exposed	1 / 40 (2.50%)	1 / 36 (2.78%)	2 / 42 (4.76%)
occurrences (all)	1	1	2
Vascular disorders			
Hypertension			
subjects affected / exposed	2 / 40 (5.00%)	0 / 36 (0.00%)	1 / 42 (2.38%)
occurrences (all)	2	0	1
Nervous system disorders			
Headache			
subjects affected / exposed	6 / 40 (15.00%)	3 / 36 (8.33%)	7 / 42 (16.67%)
occurrences (all)	6	4	10
Dizziness			

subjects affected / exposed occurrences (all)	2 / 40 (5.00%) 2	1 / 36 (2.78%) 1	2 / 42 (4.76%) 3
General disorders and administration site conditions			
Pyrexia			
subjects affected / exposed	4 / 40 (10.00%)	1 / 36 (2.78%)	1 / 42 (2.38%)
occurrences (all)	5	1	1
Asthenia			
subjects affected / exposed	2 / 40 (5.00%)	1 / 36 (2.78%)	3 / 42 (7.14%)
occurrences (all)	2	2	3
Gastrointestinal disorders			
Abdominal pain			
subjects affected / exposed	2 / 40 (5.00%)	2 / 36 (5.56%)	3 / 42 (7.14%)
occurrences (all)	2	2	3
Abdominal pain upper			
subjects affected / exposed	1 / 40 (2.50%)	3 / 36 (8.33%)	3 / 42 (7.14%)
occurrences (all)	1	3	3
Diarrhoea			
subjects affected / exposed	5 / 40 (12.50%)	9 / 36 (25.00%)	6 / 42 (14.29%)
occurrences (all)	5	10	7
Nausea			
subjects affected / exposed	2 / 40 (5.00%)	1 / 36 (2.78%)	3 / 42 (7.14%)
occurrences (all)	2	1	3
Vomiting			
subjects affected / exposed	2 / 40 (5.00%)	0 / 36 (0.00%)	0 / 42 (0.00%)
occurrences (all)	2	0	0
Toothache			
subjects affected / exposed	0 / 40 (0.00%)	1 / 36 (2.78%)	2 / 42 (4.76%)
occurrences (all)	0	1	2
Dyspepsia			
subjects affected / exposed	1 / 40 (2.50%)	1 / 36 (2.78%)	2 / 42 (4.76%)
occurrences (all)	1	1	3
Food poisoning			
subjects affected / exposed	2 / 40 (5.00%)	1 / 36 (2.78%)	1 / 42 (2.38%)
occurrences (all)	2	1	1
Haemorrhoids			

subjects affected / exposed occurrences (all)	0 / 40 (0.00%) 0	2 / 36 (5.56%) 2	0 / 42 (0.00%) 0
Respiratory, thoracic and mediastinal disorders Rhinitis allergic subjects affected / exposed occurrences (all)	3 / 40 (7.50%) 3	1 / 36 (2.78%) 1	1 / 42 (2.38%) 1
Skin and subcutaneous tissue disorders Rash subjects affected / exposed occurrences (all)	2 / 40 (5.00%) 2	3 / 36 (8.33%) 4	2 / 42 (4.76%) 2
Psychiatric disorders Anxiety subjects affected / exposed occurrences (all)	1 / 40 (2.50%) 1	3 / 36 (8.33%) 3	1 / 42 (2.38%) 1
Infections and infestations Gonorrhoea subjects affected / exposed occurrences (all)	2 / 40 (5.00%) 2	0 / 36 (0.00%) 0	2 / 42 (4.76%) 2
Gastroenteritis subjects affected / exposed occurrences (all)	1 / 40 (2.50%) 1	0 / 36 (0.00%) 0	2 / 42 (4.76%) 2
Chlamydial infection subjects affected / exposed occurrences (all)	2 / 40 (5.00%) 2	1 / 36 (2.78%) 1	1 / 42 (2.38%) 2
COVID-19 subjects affected / exposed occurrences (all)	10 / 40 (25.00%) 12	10 / 36 (27.78%) 13	6 / 42 (14.29%) 7
Influenza subjects affected / exposed occurrences (all)	3 / 40 (7.50%) 3	4 / 36 (11.11%) 5	3 / 42 (7.14%) 3
Syphilis subjects affected / exposed occurrences (all)	3 / 40 (7.50%) 3	5 / 36 (13.89%) 6	3 / 42 (7.14%) 3
Respiratory tract infection subjects affected / exposed occurrences (all)	3 / 40 (7.50%) 3	0 / 36 (0.00%) 0	1 / 42 (2.38%) 1

Pharyngitis			
subjects affected / exposed	4 / 40 (10.00%)	3 / 36 (8.33%)	1 / 42 (2.38%)
occurrences (all)	4	3	1
Nasopharyngitis			
subjects affected / exposed	4 / 40 (10.00%)	3 / 36 (8.33%)	3 / 42 (7.14%)
occurrences (all)	5	6	5
Monkeypox			
subjects affected / exposed	1 / 40 (2.50%)	1 / 36 (2.78%)	2 / 42 (4.76%)
occurrences (all)	1	1	2
Upper respiratory tract infection			
subjects affected / exposed	4 / 40 (10.00%)	1 / 36 (2.78%)	3 / 42 (7.14%)
occurrences (all)	5	1	5
Urethritis			
subjects affected / exposed	1 / 40 (2.50%)	2 / 36 (5.56%)	0 / 42 (0.00%)
occurrences (all)	1	2	0

Non-serious adverse events	Dolutegravir (DTG) 50 mg + ABC/3TC or FTC/TAF		
Total subjects affected by non-serious adverse events			
subjects affected / exposed	37 / 43 (86.05%)		
Investigations			
Blood creatine phosphokinase increased			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences (all)	0		
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Anogenital warts			
subjects affected / exposed	2 / 43 (4.65%)		
occurrences (all)	2		
Injury, poisoning and procedural complications			
Limb injury			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences (all)	1		
Vascular disorders			
Hypertension			
subjects affected / exposed	2 / 43 (4.65%)		
occurrences (all)	2		
Nervous system disorders			

Headache subjects affected / exposed occurrences (all)	4 / 43 (9.30%) 4		
Dizziness subjects affected / exposed occurrences (all)	3 / 43 (6.98%) 3		
General disorders and administration site conditions Pyrexia subjects affected / exposed occurrences (all)	4 / 43 (9.30%) 4		
Asthenia subjects affected / exposed occurrences (all)	0 / 43 (0.00%) 0		
Gastrointestinal disorders Abdominal pain subjects affected / exposed occurrences (all)	1 / 43 (2.33%) 1		
Abdominal pain upper subjects affected / exposed occurrences (all)	3 / 43 (6.98%) 3		
Diarrhoea subjects affected / exposed occurrences (all)	7 / 43 (16.28%) 8		
Nausea subjects affected / exposed occurrences (all)	6 / 43 (13.95%) 8		
Vomiting subjects affected / exposed occurrences (all)	3 / 43 (6.98%) 3		
Toothache subjects affected / exposed occurrences (all)	2 / 43 (4.65%) 3		
Dyspepsia subjects affected / exposed occurrences (all)	1 / 43 (2.33%) 1		
Food poisoning			

subjects affected / exposed occurrences (all)	0 / 43 (0.00%) 0		
Haemorrhoids subjects affected / exposed occurrences (all)	2 / 43 (4.65%) 2		
Respiratory, thoracic and mediastinal disorders Rhinitis allergic subjects affected / exposed occurrences (all)	1 / 43 (2.33%) 1		
Skin and subcutaneous tissue disorders Rash subjects affected / exposed occurrences (all)	2 / 43 (4.65%) 3		
Psychiatric disorders Anxiety subjects affected / exposed occurrences (all)	1 / 43 (2.33%) 1		
Infections and infestations Gonorrhoea subjects affected / exposed occurrences (all)	0 / 43 (0.00%) 0		
Gastroenteritis subjects affected / exposed occurrences (all)	2 / 43 (4.65%) 2		
Chlamydial infection subjects affected / exposed occurrences (all)	0 / 43 (0.00%) 0		
COVID-19 subjects affected / exposed occurrences (all)	12 / 43 (27.91%) 12		
Influenza subjects affected / exposed occurrences (all)	4 / 43 (9.30%) 4		
Syphilis subjects affected / exposed occurrences (all)	1 / 43 (2.33%) 1		

Respiratory tract infection			
subjects affected / exposed	0 / 43 (0.00%)		
occurrences (all)	0		
Pharyngitis			
subjects affected / exposed	4 / 43 (9.30%)		
occurrences (all)	5		
Nasopharyngitis			
subjects affected / exposed	3 / 43 (6.98%)		
occurrences (all)	4		
Monkeypox			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences (all)	1		
Upper respiratory tract infection			
subjects affected / exposed	3 / 43 (6.98%)		
occurrences (all)	3		
Urethritis			
subjects affected / exposed	1 / 43 (2.33%)		
occurrences (all)	1		

More information

Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
25 September 2020	This is a global amendment and was created primarily to replace details of the internal Safety Review Committee to reflect the use instead of an Independent Data Monitoring Committee (IDMC). The decision to switch was based on feedback received from several health authorities and ethics committees, with advice and approval by VH guidance committee. This amendment also includes corrections, clarifications and minor administrative errors. In addition, details from 2 prior country-specific amendments (Canada amendment CAN-1 and Portugal amendment POR-1) were incorporated into this global amendment.
09 April 2021	This is a global amendment. This amendment was created primarily to make non-substantial changes generally related to inclusion/exclusion criteria and the screening period, esophagogastroduodenoscopy (EGD) biopsy-based stopping criteria for individual participants and the study overall. The other changes exist for further detail, clarity, and current information.
10 May 2021	Study 208379 is undergoing this substantial amendment to reduce the sample size from approximately 210 participants to approximately 150 participants. This reduction of 60 participants will occur equally among the 3 experimental arms. There is no change to the Dolutegravir (DTG) reference arm. Thus, the number of participants will be as follows (each arm in combination with open-label Nucleoside Reverse Transcriptase Inhibitor [NRTI] backbone as originally described):

Notes:

Interruptions (globally)

Were there any global interruptions to the trial? No

Limitations and caveats

None reported