

TABLE 14.2.3.20.1 Summary of Hs-CRP and change from baseline (Continued)
FULL ANALYSIS SET

	Reparixin N=182		Placebo N=88	
High-Sensitivity C Reactive Protein (mg/L)				
End of Treatment				
n	7		6	
Mean (SD)	104.94	(121.85)	60.30	(136.74)
Median	62.10		4.70	
Q1; Q3	1.40 ;	255.60	1.50 ;	10.50
Min; Max	1.2 ;	290.6	1.1 ;	339.3
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.432
CHANGE FROM BASELINE				
n	7		6	
Mean (SD)	37.50	(151.54)	-37.68	(76.26)
Median	16.90		-19.05	
Q1; Q3	-103.70 ;	194.70	-92.20 ;	0.70
Min; Max	-112.3 ;	275.7	-156.9 ;	60.4
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.432

Note 1: Comparison between treatment arms is performed by means of two-sample Mann-Whitney U test.

Note 2: Hs-CRP = High-sensitivity C-reactive protein.

Cross-reference: Listing 16.2.8.2

Program (Date Time): (19Jan2022 19:54) SAS Version: 9.4

Page 6 of 9

TABLE 14.2.3.20.1 Summary of Hs-CRP and change from baseline (Continued)
FULL ANALYSIS SET

	Reparixin N=182		Placebo N=88	
High-Sensitivity C Reactive Protein (mg/L)				
Day 28 (+/-2)				
n	4		2	
Mean (SD)	9.48	(10.82)	19.90	(22.34)
Median	4.90		19.90	
Q1; Q3	2.95 ;	16.00	4.10 ;	35.70
Min; Max	2.6 ;	25.5	4.1 ;	35.7
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.487
CHANGE FROM BASELINE				
n	4		2	
Mean (SD)	-42.60	(30.15)	9.65	(26.94)
Median	-36.90		9.65	
Q1; Q3	-61.35 ;	-23.85	-9.40 ;	28.70
Min; Max	-84.3 ;	-12.3	-9.4 ;	28.7
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.105

Note 1: Comparison between treatment arms is performed by means of two-sample Mann-Whitney U test.

Note 2: Hs-CRP = High-sensitivity C-reactive protein.

Cross-reference: Listing 16.2.8.2

Program (Date Time): (19Jan2022 19:54) SAS Version: 9.4

Page 7 of 9

TABLE 14.2.3.20.1 Summary of Hs-CRP and change from baseline (Continued)
FULL ANALYSIS SET

	Reparixin N=182		Placebo N=88	
High-Sensitivity C Reactive Protein (mg/L)				
Hospital Discharge				
n	5		3	
Mean (SD)	22.14	(28.44)	3.57	(3.93)
Median	3.20		1.50	
Q1; Q3	1.60 ;	42.60	1.10 ;	8.10
Min; Max	1.2 ;	62.1	1.1 ;	8.1
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.371
CHANGE FROM BASELINE				
n	5		3	
Mean (SD)	-54.46	(63.12)	-84.83	(76.02)
Median	-86.00		-92.20	
Q1; Q3	-103.70 ;	11.00	-156.90 ;	-5.40
Min; Max	-110.5 ;	16.9	-156.9 ;	-5.4
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.551

Note 1: Comparison between treatment arms is performed by means of two-sample Mann-Whitney U test.

Note 2: Hs-CRP = High-sensitivity C-reactive protein.

Cross-reference: Listing 16.2.8.2

Program (Date Time): (19Jan2022 19:54) SAS Version: 9.4

Page 8 of 9