

TABLE 14.2.3.19.1 Summary of PaO₂/FiO₂ ratio and change from baseline (Continued)
FULL ANALYSIS SET

	Reparixin N=182		Placebo N=88	
P/F ratio (mmHg)				
End of Treatment				
n	140		66	
Mean (SD)	334.468	(151.717)	304.273	(153.055)
Median	344.500		296.430	
Q1; Q3	219.500 ;	417.310	195.000 ;	380.000
Min; Max	53.00 ;	690.00	50.00 ;	690.00
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.152
CHANGE FROM BASELINE				
n	140		66	
Mean (SD)	140.575	(139.619)	106.332	(144.608)
Median	143.500		95.500	
Q1; Q3	25.500 ;	214.000	16.000 ;	204.000
Min; Max	-188.52 ;	564.00	-221.00 ;	445.00
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.133

Note 1: Comparison between treatment arms is performed by means of two-sample Mann-Whitney U test.

Note 2: PaO₂/FiO₂ ratio = Partial arteriolar oxygen pressure to fraction of inspiration O₂ ratio.

Cross-reference: Listing 16.2.6.1.8

Program (Date Time): (19Jan2022 19:54) SAS Version: 9.4

Page 6 of 10

TABLE 14.2.3.19.1 Summary of PaO₂/FiO₂ ratio and change from baseline (Continued)
FULL ANALYSIS SET

	Reparixin N=182		Placebo N=88	
P/F ratio (mmHg)				
Hospital Discharge				
n	96		48	
Mean (SD)	436.527	(121.855)	413.328	(107.256)
Median	409.000		376.595	
Q1; Q3	347.070 ;	525.430	340.810 ;	468.500
Min; Max	172.00 ;	861.43	144.00 ;	690.00
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.380
CHANGE FROM BASELINE				
n	96		48	
Mean (SD)	228.999	(119.337)	210.765	(112.418)
Median	216.500		203.000	
Q1; Q3	147.500 ;	286.560	135.000 ;	291.500
Min; Max	-9.00 ;	593.93	-8.38 ;	445.00
Comparison Between Treatments (p-value[1]):				
Reparixin vs Placebo				0.593

Note 1: Comparison between treatment arms is performed by means of two-sample Mann-Whitney U test.

Note 2: PaO₂/FiO₂ ratio = Partial arteriolar oxygen pressure to fraction of inspiration O₂ ratio.

Cross-reference: Listing 16.2.6.1.8

Program (Date Time): (19Jan2022 19:54) SAS Version: 9.4

Page 8 of 10