



## Clinical trial results:

### A Randomized, Phase 3, Double-Masked, Parallel Group, Multicenter Study to Compare the Efficacy and Safety of ALT-L9 Versus Eylea® in Patients With Neovascular Age-Related Macular Degeneration (ALTERA)

#### Summary

EudraCT number	2021-004530-11
Trial protocol	CZ HU SK LV AT ES BG EE LT
Global end of trial date	20 February 2024

#### Results information

Result version number	v1 (current)
This version publication date	07 March 2025
First version publication date	07 March 2025

#### Trial information

##### Trial identification

Sponsor protocol code	ALT-L9-03
-----------------------	-----------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	-
WHO universal trial number (UTN)	-

Notes:

##### Sponsors

Sponsor organisation name	Altos Biologics Inc.
Sponsor organisation address	8F, 15, Teheran-ro 84-gil, Gangnam-gu, Seoul, Korea, Republic of, 06179
Public contact	SoJin Lee, Altos Biologics Inc., +82 2 2039 9520, sjlee@altosbio.com
Scientific contact	SoJin Lee, Altos Biologics Inc., +82 2 2039 9520, sjlee@altosbio.com

Notes:

##### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	No

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	18 April 2024
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	20 February 2024
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

The primary objective of the study was to demonstrate that the biosimilar candidate ALT-L9 2 mg/50 microliter (mcL) was equivalent to Eylea® (aflibercept) in subjects with wet (neovascular) age-related macular degeneration (nAMD) in terms of best-corrected visual acuity (BCVA).

Protection of trial subjects:

This study was conducted in accordance with the accepted version of the Declaration of Helsinki and/or all relevant federal regulations in compliance with the International Council for Harmonization of Technical Requirements for Pharmaceuticals for Human Use good clinical practice (ICH GCP) guidelines, Japanese GCP and Korean GCP, and according to the appropriate regulatory requirements in the countries where the study was conducted.

Background therapy: -

Evidence for comparator: -

Actual start date of recruitment	02 June 2022
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	Austria: 7
Country: Number of subjects enrolled	Bulgaria: 16
Country: Number of subjects enrolled	Czechia: 56
Country: Number of subjects enrolled	Estonia: 6
Country: Number of subjects enrolled	Hungary: 97
Country: Number of subjects enrolled	Latvia: 35
Country: Number of subjects enrolled	Lithuania: 9
Country: Number of subjects enrolled	Poland: 92
Country: Number of subjects enrolled	Slovakia: 19
Country: Number of subjects enrolled	Spain: 22
Country: Number of subjects enrolled	Japan: 33
Country: Number of subjects enrolled	Korea, Republic of: 39
Worldwide total number of subjects	431
EEA total number of subjects	359

Notes:

---

**Subjects enrolled per age group**

---

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	0
Adolescents (12-17 years)	0
Adults (18-64 years)	49
From 65 to 84 years	345
85 years and over	37

---

## Subject disposition

### Recruitment

Recruitment details:

Subjects took part in the study at 79 investigative sites in Japan, Republic of Korea, Austria, Bulgaria, Czech Republic, Estonia, Hungary, Latvia, Lithuania, Poland, Slovakia and Spain from 02 June 2022 to 20 February 2024.

### Pre-assignment

Screening details:

A total of 642 subjects with Neovascular Age-Related Macular Degeneration (nAMD) were screened, of which 211 subjects were screen failures and 431 subjects were randomized to receive ALT-L9 or Eylea.

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator, Monitor, Data analyst, Assessor

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	ALT-L9

Arm description:

Subjects received single dose of ALT-L9 40 milligrams per milliliter (mg/mL) (2 mg/50 microliter [mcL] aflibercept), ophthalmic intravitreal (IVT) injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

Arm type	Experimental
Investigational medicinal product name	Aflibercept biosimilar
Investigational medicinal product code	ALT-L9
Other name	
Pharmaceutical forms	Solution for injection in vial
Routes of administration	Intravitreal use

Dosage and administration details:

ALT-L9 40 mg/mL (2 mg/50 mcL aflibercept), ophthalmic IVT injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48. The 40 mg/mL concentration of aflibercept was contained in a single vial.

<b>Arm title</b>	Eylea®
------------------	--------

Arm description:

Subjects received single dose of Eylea® 40 mg/mL (2 mg/50 mcL aflibercept), ophthalmic IVT injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

Arm type	Active comparator
Investigational medicinal product name	Eylea®
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection in pre-filled syringe
Routes of administration	Intravitreal use

Dosage and administration details:

Eylea® 40 mg/mL (2 mg/50 mcL aflibercept), ophthalmic IVT injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48. The 40 mg/mL concentration of aflibercept was contained in a PFS.

<b>Number of subjects in period 1</b>	ALT-L9	Eylea®
Started	216	215
Completed	204	201
Not completed	12	14
Consent withdrawn by subject	3	4
Physician decision	1	-
Adverse event, non-fatal	4	8
Death	1	-
Unspecified	2	1
Lost to follow-up	1	1

## Baseline characteristics

### Reporting groups

Reporting group title	ALT-L9
-----------------------	--------

Reporting group description:

Subjects received single dose of ALT-L9 40 milligrams per milliliter (mg/mL) (2 mg/50 microliter [mL] aflibercept), ophthalmic intravitreal (IVT) injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

Reporting group title	Eylea®
-----------------------	--------

Reporting group description:

Subjects received single dose of Eylea® 40 mg/mL (2 mg/50 mL aflibercept), ophthalmic IVT injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

Reporting group values	ALT-L9	Eylea®	Total
Number of subjects	216	215	431
Age categorical Units: Subjects			
Age continuous Units: years arithmetic mean standard deviation	74.4 ± 7.83	73.9 ± 7.97	-
Gender categorical Units: Subjects			
Female	132	126	258
Male	84	89	173

## End points

### End points reporting groups

Reporting group title	ALT-L9
-----------------------	--------

Reporting group description:

Subjects received single dose of ALT-L9 40 milligrams per milliliter (mg/mL) (2 mg/50 microliter [mcL] aflibercept), ophthalmic intravitreal (IVT) injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

Reporting group title	Eylea®
-----------------------	--------

Reporting group description:

Subjects received single dose of Eylea® 40 mg/mL (2 mg/50 mcL aflibercept), ophthalmic IVT injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

### Primary: Change from Baseline in Best-corrected Visual Acuity (BCVA) at Week 8 as Measured by the Early Treatment Diabetic Retinopathy Study (ETDRS) Letter Score

End point title	Change from Baseline in Best-corrected Visual Acuity (BCVA) at Week 8 as Measured by the Early Treatment Diabetic Retinopathy Study (ETDRS) Letter Score
-----------------	----------------------------------------------------------------------------------------------------------------------------------------------------------

End point description:

BCVA was measured on the Early Treatment Diabetic Retinopathy Study (ETDRS) chart. The BCVA letter score ranges from 0 to 100 (best score). A positive change indicates an improvement, and a negative change indicates worsening. The Intent-to-treat (ITT) set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation. Here, "n" refers to subjects who were evaluable for this endpoint and for specified estimands.

End point type	Primary
----------------	---------

End point timeframe:

Baseline and at Week 8

End point values	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: score on a scale				
least squares mean (standard error)				
Primary Estimand (n=216,215)	5.771 (± 0.5821)	7.863 (± 0.5888)		
Secondary Estimand (n=216,215)	6.272 (± 0.6240)	7.972 (± 0.6248)		
Tertiary Estimand (n=213,214)	5.782 (± 0.5870)	7.862 (± 0.5905)		

### Statistical analyses

Statistical analysis title	Primary Estimand: ALT-L9 vs Eylea
Comparison groups	ALT-L9 v Eylea®

Number of subjects included in analysis	431
Analysis specification	Pre-specified
Analysis type	equivalence <sup>[1]</sup>
Parameter estimate	Least-square Mean Difference
Point estimate	-2.092
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.431
upper limit	-0.753
Variability estimate	Standard error of the mean
Dispersion value	0.6834

Notes:

[1] - A predefined margin of  $\pm 3.49$  letters was used for analysis.

<b>Statistical analysis title</b>	Secondary Estimand: ALT-L9 vs Eylea
Comparison groups	ALT-L9 v Eylea®
Number of subjects included in analysis	431
Analysis specification	Pre-specified
Analysis type	equivalence <sup>[2]</sup>
Parameter estimate	Least-square Mean Difference
Point estimate	-1.7
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.113
upper limit	-0.287
Variability estimate	Standard error of the mean
Dispersion value	0.7209

Notes:

[2] - A predefined margin of  $\pm 3.49$  letters was used for analysis.

<b>Statistical analysis title</b>	Tertiary Estimand: ALT-L9 vs Eylea
Comparison groups	ALT-L9 v Eylea®
Number of subjects included in analysis	431
Analysis specification	Pre-specified
Analysis type	equivalence <sup>[3]</sup>
Parameter estimate	Least-square Mean Difference
Point estimate	-2.079
Confidence interval	
level	95 %
sides	2-sided
lower limit	-3.425
upper limit	-0.734
Variability estimate	Standard error of the mean
Dispersion value	0.6846

Notes:

[3] - A predefined margin of  $\pm 3.49$  letters was used for analysis.

## Secondary: Proportion of Subjects With a Loss of at Least 5, 10, or 15 Letters in

## BCVA Letter Score in the Study Eye Over Time up to Week 52, Compared With Baseline, Using the ETDRS Protocol

End point title	Proportion of Subjects With a Loss of at Least 5, 10, or 15 Letters in BCVA Letter Score in the Study Eye Over Time up to Week 52, Compared With Baseline, Using the ETDRS Protocol
-----------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

End point description:

BCVA was measured on the ETDRS chart. The BCVA letter score ranges from 0 to 100 (best score). A positive change indicates an improvement, and a negative change indicates worsening. The ITT set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to Week 52

End point values	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: percentage of subjects				
number (not applicable)				
Primary Estimand: $\geq 5$ letters	10.6	4.0		
Secondary Estimand: $\geq 5$ letters	8.1	3.7		
Tertiary Estimand: $\geq 5$ letters	10.3	4.0		
Primary Estimand: $\geq 10$ letters	6.4	2.0		
Secondary Estimand: $\geq 10$ letters	3.5	1.5		
Tertiary Estimand: $\geq 10$ letters	6.4	2.0		
Primary Estimand: $\geq 15$ letters	3.4	1.0		
Secondary Estimand: $\geq 15$ letters	2.0	0.6		
Tertiary Estimand: $\geq 15$ letters	3.4	1.0		

### Statistical analyses

No statistical analyses for this end point

## Secondary: Change From Baseline in Central Subfield Thickness (CST) in the Study Eye Up to Week 4 and Week 52 as Measured by Spectral Domain-optical Coherence Tomography (SD-OCT)

End point title	Change From Baseline in Central Subfield Thickness (CST) in the Study Eye Up to Week 4 and Week 52 as Measured by Spectral Domain-optical Coherence Tomography (SD-OCT)
-----------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------

End point description:

CST was measured in the study eye by SD-OCT. The ITT set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation. Here, "n" refers to subjects who were evaluable for specified estimands.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline, up to Weeks 4 and 52

<b>End point values</b>	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: microns				
least squares mean (standard error)				
Week 4: Primary Estimand (n=216,215)	-100.485 (± 6.2919)	-93.569 (± 6.3687)		
Week 4: Secondary Estimand (n=216,215)	-98.250 (± 6.7016)	-93.151 (± 6.8495)		
Week 4: Tertiary Estimand (n=213,213)	-101.102 (± 6.2981)	-93.734 (± 6.3697)		
Week 52: Primary Estimand (n=216,215)	-116.023 (± 7.4471)	-111.281 (± 7.5733)		
Week 52: Secondary Estimand (n=216,215)	-113.450 (± 8.1122)	-111.471 (± 8.2944)		
Week 52: Tertiary Estimand (n=203,195)	-119.154 (± 7.3191)	-108.528 (± 7.4820)		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Proportion of Subjects With a Gain of at least 5, 10, or 15 Letters in BCVA Letter Score in the Study Eye Over Time up to Week 52, Compared With Baseline, Using the ETDRS Protocol

End point title	Proportion of Subjects With a Gain of at least 5, 10, or 15 Letters in BCVA Letter Score in the Study Eye Over Time up to Week 52, Compared With Baseline, Using the ETDRS Protocol
End point description:	BCVA was measured on the ETDRS chart. The BCVA letter score ranges from 0 to 100 (best score). A positive change indicates an improvement, and a negative change indicates worsening. The ITT set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation.
End point type	Secondary
End point timeframe:	Baseline up to Week 52

<b>End point values</b>	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: percentage of subjects				
number (not applicable)				
Primary Estimand: >=5 letters	68.8	72.9		
Secondary Estimand: >=5 letters	71.0	74.2		
Tertiary Estimand: >=5 letters	69.6	73.1		
Primary Estimand: >=10 letters	41.5	44.6		

Secondary Estimand: >=10 letters	43.9	47.2		
Tertiary Estimand: >=10 letters	42.2	44.3		
Primary Estimand: >=15 letters	19.2	21.8		
Secondary Estimand: >=15 letters	21.5	25.0		
Tertiary Estimand: >=15 letters	19.6	21.4		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Proportion of Subjects With Existing Intraretinal or Subretinal Fluid in the Study Eye Over Time up to Week 4 and Week 52, Compared With Baseline, as Measured by SD-OCT

End point title	Proportion of Subjects With Existing Intraretinal or Subretinal Fluid in the Study Eye Over Time up to Week 4 and Week 52, Compared With Baseline, as Measured by SD-OCT
End point description:	Existing intraretinal or subretinal fluid was measured in the study eye by SD-OCT. The ITT set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation.
End point type	Secondary
End point timeframe:	Up to Weeks 4 and 52

End point values	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: percentage of subjects				
number (not applicable)				
Week 4: Primary Estimand	62.2	56.9		
Week 4: Secondary Estimand	61.7	56.2		
Week 4: Tertiary Estimand	62.1	56.8		
Week 52: Primary Estimand	39.7	39.1		
Week 52: Secondary Estimand	42.4	40.5		
Week 52: Tertiary Estimand	38.2	37.5		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change From Baseline in BCVA Letter Score in The Study Eye Over Time up to Week 52 Using the ETDRS Protocol

End point title	Change From Baseline in BCVA Letter Score in The Study Eye Over Time up to Week 52 Using the ETDRS Protocol
End point description:	BCVA was measured on the ETDRS chart at a starting distance of 4 meters. The BCVA letter score ranges from 0 to 100 (best score). A positive change indicates an improvement, and a negative change

indicates worsening. The ITT set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation. Here, "n" refers to subjects who were evaluable for specified estimands.

End point type	Secondary
End point timeframe:	
Baseline up to Week 52	

End point values	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: score on a scale				
least squares mean (standard error)				
Primary Estimand (n=216, 215)	7.315 (± 0.7871)	9.259 (± 0.8018)		
Secondary Estimand (n=216, 215)	8.171 (± 0.9405)	9.551 (± 0.8940)		
Tertiary Estimand (n=204, 201)	7.415 (± 0.7872)	9.228 (± 0.8014)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Change From Baseline in the Total Size of the Active Choroidal Neovascularization (CNV) Area in the Study Eye Over Time up to Week 52, as Measured by Fluorescein Angiography (FA)

End point title	Change From Baseline in the Total Size of the Active Choroidal Neovascularization (CNV) Area in the Study Eye Over Time up to Week 52, as Measured by Fluorescein Angiography (FA)
-----------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

End point description:

Active CNV leakage of the study eye was assessed with fluorescein angiography. The active CNV leakage was defined as result of  $>0$  (mm<sup>2</sup>). ITT set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation. Here, "n" refers to subjects who were evaluable for this endpoint and for specified estimands.

End point type	Secondary
End point timeframe:	
Baseline up to Week 52	

End point values	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: square millimeter (mm <sup>2</sup> )				
least squares mean (standard error)				
Primary Estimand (n=216,215)	-1.890 (± 0.4174)	-2.123 (± 0.4188)		

Secondary Estimand (n=216,215)	-1.992 (± 0.5344)	-2.130 (± 0.5201)		
Tertiary Estimand (n=199,194)	-1.857 (± 0.4205)	-2.100 (± 0.4242)		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Number of Subjects with Treatment-related Treatment-emergent Adverse Events (TEAEs) and Serious Adverse Events (SAEs) for Systemic and Ocular Category

End point title	Number of Subjects with Treatment-related Treatment-emergent Adverse Events (TEAEs) and Serious Adverse Events (SAEs) for Systemic and Ocular Category
-----------------	--------------------------------------------------------------------------------------------------------------------------------------------------------

End point description:

A treatment-related TEAEs and SAEs were clinical event with plausible time relationship to study drug administration, and that cannot be explained by concurrent disease or other drugs or chemicals. The SAF included all randomized subjects who receive at least 1 administration (full or partial) of the study drug.

End point type	Secondary
----------------	-----------

End point timeframe:

Baseline up to Week 52

End point values	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: subjects				
Systemic Category: Related TEAEs	0	2		
Systemic Category: Related SAEs	0	2		
Ocular Category: Related TEAEs	3	3		
Ocular Category: Related SAEs	0	0		

## Statistical analyses

No statistical analyses for this end point

### Secondary: Number of Subjects With Treatment-emergent AEs (TEAEs) and Serious AEs (SAEs) for Systemic and Ocular Category

End point title	Number of Subjects With Treatment-emergent AEs (TEAEs) and Serious AEs (SAEs) for Systemic and Ocular Category
-----------------	----------------------------------------------------------------------------------------------------------------

End point description:

A TEAE was defined as any event not present before the initiation of the study treatment or any event already present that worsens in either intensity or frequency following exposure to the treatments. An SAE is any untoward medical occurrence, in the view of either the investigator or Sponsor, that meets one or more of the following criteria: results in death, is life-threatening, results in inpatient hospitalization or prolongation of existing hospitalization, results in persistent or significant disability/incapacity, is a congenital anomaly/birth defect. The SAF included of all randomized subjects

who receive at least 1 administration (full or partial) of the study drug.

End point type	Secondary
End point timeframe:	
Baseline up to Week 52	

<b>End point values</b>	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: subjects				
Systemic Category: TEAEs	116	112		
Systemic Category: SAEs	25	25		
Ocular Category: TEAEs	90	75		
Ocular Category: SAEs	1	3		

### Statistical analyses

No statistical analyses for this end point

### Secondary: Proportion of Subjects With Active CNV Leakage in the Study Eye Over Time up to Week 52, Compared With Baseline, as Measured by FA

End point title	Proportion of Subjects With Active CNV Leakage in the Study Eye Over Time up to Week 52, Compared With Baseline, as Measured by FA
-----------------	------------------------------------------------------------------------------------------------------------------------------------

End point description:

Active CNV leakage of the study eye was assessed with fluorescein angiography. The active CNV leakage was defined as result of  $>0$  (mm<sup>2</sup>). ITT set included all randomized subjects, irrespective of any deviation from the protocol or premature discontinuation.

End point type	Secondary
End point timeframe:	
Up to Week 52	

<b>End point values</b>	ALT-L9	Eylea®		
Subject group type	Reporting group	Reporting group		
Number of subjects analysed	216	215		
Units: percentage of subjects				
number (not applicable)				
Primary Estimand	61.7	55.1		
Secodary Estimand	64.4	56.6		
Tertiary Estimand	62.3	54.9		

### Statistical analyses

---

No statistical analyses for this end point

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

Baseline up to Week 52

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	26.1
--------------------	------

### Reporting groups

Reporting group title	ALT-L9
-----------------------	--------

Reporting group description:

Subjects received single dose of ALT-L9 40 mg/mL (2 mg/50 mcL aflibercept), ophthalmic IVT injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

Reporting group title	Eylea®
-----------------------	--------

Reporting group description:

Subjects received single dose of Eylea® 40 mg/mL (2 mg/50 mcL aflibercept), ophthalmic IVT injection, in study eye, once every 4 weeks, for up to Week 8 and once every 8 weeks up to Week 48.

<b>Serious adverse events</b>	ALT-L9	Eylea®	
Total subjects affected by serious adverse events			
subjects affected / exposed	26 / 216 (12.04%)	28 / 215 (13.02%)	
number of deaths (all causes)	1	0	
number of deaths resulting from adverse events	0	0	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Acute myeloid leukaemia			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Bladder transitional cell carcinoma			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Follicular thyroid cancer			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Glioma			

subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 1	0 / 0	
Hormone-dependent prostate cancer			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lung adenocarcinoma			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lung cancer metastatic			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Lung neoplasm malignant			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Oesophageal carcinoma			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Polycythaemia vera			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Prostate cancer			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Salivary gland neoplasm			

subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Vascular disorders</b>			
Arteriosclerosis			
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Orthostatic hypotension			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Reproductive system and breast disorders</b>			
Cystocele			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Uterovaginal prolapse			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Respiratory, thoracic and mediastinal disorders</b>			
Chronic obstructive pulmonary disease			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Psychiatric disorders</b>			
Anxiety disorder			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Injury, poisoning and procedural complications</b>			
Accidental overdose			

subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Humerus fracture</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Rib fracture</b>			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Cardiac disorders</b>			
<b>Acute myocardial infarction</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Arrhythmia</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Atrial fibrillation</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Cardiac failure</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Coronary artery disease</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Myocardial ischaemia</b>			

subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Nervous system disorders</b>			
<b>Ischaemic stroke</b>			
subjects affected / exposed	1 / 216 (0.46%)	3 / 215 (1.40%)	
occurrences causally related to treatment / all	0 / 1	1 / 3	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Epilepsy</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Haemorrhage intracranial</b>			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Intracranial aneurysm</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Thrombotic cerebral infarction</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	1 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Trigeminal palsy</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Blood and lymphatic system disorders</b>			
<b>Anaemia</b>			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Blood loss anaemia</b>			

subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Ear and labyrinth disorders</b>			
Vertigo positional			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Eye disorders</b>			
Blindness			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
<b>Gastrointestinal disorders</b>			
Gastric ulcer haemorrhage			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Haemorrhoids			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Ileus			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Inguinal hernia			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Intestinal haemorrhage			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	

Pancreatitis acute			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Hepatobiliary disorders			
Cholelithiasis			
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 1	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Skin and subcutaneous tissue disorders			
Pemphigoid			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Renal and urinary disorders			
Renal colic			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Urethral stenosis			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Musculoskeletal and connective tissue disorders			
Osteoarthritis			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences causally related to treatment / all	0 / 1	0 / 0	
deaths causally related to treatment / all	0 / 0	0 / 0	
Spinal stenosis			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences causally related to treatment / all	0 / 0	0 / 1	
deaths causally related to treatment / all	0 / 0	0 / 0	
Infections and infestations			
COVID-19			

subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences causally related to treatment / all	0 / 2	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Pneumonia		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Pneumonia aspiration		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Pyelonephritis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Urethritis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Vulval abscess		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences causally related to treatment / all	0 / 1	0 / 0
deaths causally related to treatment / all	0 / 0	0 / 0
Endophthalmitis		
subjects affected / exposed	1 / 216 (0.46%)	2 / 215 (0.93%)
occurrences causally related to treatment / all	0 / 1	0 / 2
deaths causally related to treatment / all	0 / 0	0 / 0

Frequency threshold for reporting non-serious adverse events: 0 %

<b>Non-serious adverse events</b>	ALT-L9	Eylea®	
Total subjects affected by non-serious adverse events			
subjects affected / exposed	204 / 216 (94.44%)	180 / 215 (83.72%)	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)			
Basal cell carcinoma			
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)	
occurrences (all)	2	0	
Salivary gland neoplasm			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	2	0	
Adenoma benign			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Bowen's disease			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Glioma			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Laryngeal cancer			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Metastases to bone			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Neoplasm			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Oesophageal carcinoma			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Polycythaemia vera			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Vascular disorders			

Hypertension			
subjects affected / exposed	11 / 216 (5.09%)	10 / 215 (4.65%)	
occurrences (all)	13	11	
Arteriosclerosis			
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)	
occurrences (all)	1	1	
Blood pressure fluctuation			
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)	
occurrences (all)	1	1	
Hot flush			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Hypertensive crisis			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Subclavian artery stenosis			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Thrombophlebitis			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Venous thrombosis limb			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
White coat hypertension			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
General disorders and administration site conditions			
Asthenia			
subjects affected / exposed	1 / 216 (0.46%)	3 / 215 (1.40%)	
occurrences (all)	1	3	
Peripheral swelling			
subjects affected / exposed	2 / 216 (0.93%)	1 / 215 (0.47%)	
occurrences (all)	2	1	
Pyrexia			

subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	0 / 215 (0.00%) 0	
Chest pain subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Hernia subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Immune system disorders Drug hypersensitivity subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	1 / 215 (0.47%) 1	
Contrast media allergy subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Reproductive system and breast disorders Benign prostatic hyperplasia subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	1 / 215 (0.47%) 1	
Uterine polyp subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Cystocele subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Endometrial hyperplasia subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Uterine prolapse subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Uterovaginal prolapse subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Respiratory, thoracic and mediastinal disorders			

Cough			
subjects affected / exposed	6 / 216 (2.78%)	1 / 215 (0.47%)	
occurrences (all)	6	1	
Upper respiratory tract inflammation			
subjects affected / exposed	1 / 216 (0.46%)	2 / 215 (0.93%)	
occurrences (all)	1	2	
Rhinorrhoea			
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)	
occurrences (all)	1	1	
Lung opacity			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Oropharyngeal pain			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Paranasal sinus inflammation			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Rhinitis allergic			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Psychiatric disorders			
Insomnia			
subjects affected / exposed	0 / 216 (0.00%)	3 / 215 (1.40%)	
occurrences (all)	0	3	
Anxiety			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Anxiety disorder			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Body dysmorphic disorder			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Delirium			

subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Depression subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Sleep disorder subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Product issues Device dislocation subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Investigations Blood pressure increased subjects affected / exposed occurrences (all)	4 / 216 (1.85%) 4	1 / 215 (0.47%) 1	
Gamma-glutamyltransferase increased subjects affected / exposed occurrences (all)	3 / 216 (1.39%) 3	1 / 215 (0.47%) 1	
Blood triglycerides increased subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	3 / 215 (1.40%) 3	
Blood urea increased subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	2 / 215 (0.93%) 2	
Aspartate aminotransferase increased subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Blood creatinine increased subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Blood glucose increased subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	0 / 215 (0.00%) 0	
Alanine aminotransferase increased			

subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Anti-thyroid antibody increased		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Blood alkaline phosphatase increased		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Blood creatine phosphokinase increased		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Blood glucose decreased		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Blood lactate dehydrogenase increased		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Blood phosphorus decreased		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Blood thyroid stimulating hormone increased		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Body temperature increased		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Fibrin D dimer increased		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Glomerular filtration rate decreased		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Influenza A virus test positive		

subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Platelet count increased subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Ultrasound liver abnormal subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Vitamin B12 decreased subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Intraocular pressure increased subjects affected / exposed occurrences (all)	4 / 216 (1.85%) 7	3 / 215 (1.40%) 5	
Optical coherence tomography abnormal subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Injury, poisoning and procedural complications			
Fall subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Tooth fracture subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	2 / 215 (0.93%) 2	
Rib fracture subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 2	0 / 215 (0.00%) 0	
Bone contusion subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Contusion subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Hand fracture			

subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Head injury		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Humerus fracture		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Limb injury		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Patella fracture		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Pelvic fracture		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Post-traumatic pain		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Skin wound		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Synovial rupture		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Ulna fracture		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Upper limb fracture		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Wrist fracture		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Corneal abrasion		

subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Craniofacial fracture subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Injury of conjunctiva subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
<b>Congenital, familial and genetic disorders</b>			
Atrial septal defect subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Type IIa hyperlipidaemia subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
<b>Cardiac disorders</b>			
Cardiac failure chronic subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	2 / 215 (0.93%) 2	
Angina pectoris subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	2 / 215 (0.93%) 2	
Arrhythmia subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	0 / 215 (0.00%) 0	
Myocardial ischaemia subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	2 / 215 (0.93%) 2	
Palpitations subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Atrial fibrillation subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Cardiac asthma			

subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Chronic coronary syndrome subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Coronary artery disease subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Tricuspid valve incompetence subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
<b>Nervous system disorders</b>			
Headache subjects affected / exposed occurrences (all)	7 / 216 (3.24%) 8	4 / 215 (1.86%) 5	
Intracranial aneurysm subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 2	
Lumbar radiculopathy subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 2	
Cognitive disorder subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	0 / 215 (0.00%) 0	
Dizziness subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Memory impairment subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	0 / 215 (0.00%) 0	
Polyneuropathy subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	2 / 215 (0.93%) 2	
Trigeminal palsy subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 2	

Balance disorder		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Carotid artery stenosis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Encephalomalacia		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Hypoaesthesia		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Loss of consciousness		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Lumbosacral radiculopathy		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Migraine		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Neuropathy peripheral		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Paresis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Peripheral sensorimotor neuropathy		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Sciatica		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Spinal cord herniation		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1

Thrombotic cerebral infarction subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Vascular dementia subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
<b>Blood and lymphatic system disorders</b>			
Anaemia subjects affected / exposed occurrences (all)	3 / 216 (1.39%) 3	0 / 215 (0.00%) 0	
Blood loss anaemia subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Leukocytosis subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Leukopenia subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
<b>Ear and labyrinth disorders</b>			
Vertigo subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	0 / 215 (0.00%) 0	
Vertigo positional subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	2 / 215 (0.93%) 2	
Cerumen impaction subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Ear pain subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Tinnitus subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Vestibular disorder			

subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Eye disorders			
Neovascular age-related macular degeneration			
subjects affected / exposed	23 / 216 (10.65%)	13 / 215 (6.05%)	
occurrences (all)	24	13	
Visual acuity reduced			
subjects affected / exposed	12 / 216 (5.56%)	10 / 215 (4.65%)	
occurrences (all)	12	11	
Cataract			
subjects affected / exposed	8 / 216 (3.70%)	4 / 215 (1.86%)	
occurrences (all)	8	5	
Subretinal fluid			
subjects affected / exposed	6 / 216 (2.78%)	4 / 215 (1.86%)	
occurrences (all)	6	4	
Conjunctival haemorrhage			
subjects affected / exposed	2 / 216 (0.93%)	6 / 215 (2.79%)	
occurrences (all)	2	6	
Posterior capsule opacification			
subjects affected / exposed	3 / 216 (1.39%)	5 / 215 (2.33%)	
occurrences (all)	3	5	
Vitreous floaters			
subjects affected / exposed	4 / 216 (1.85%)	3 / 215 (1.40%)	
occurrences (all)	4	3	
Retinal oedema			
subjects affected / exposed	3 / 216 (1.39%)	3 / 215 (1.40%)	
occurrences (all)	3	3	
Retinal pigment epithelial tear			
subjects affected / exposed	3 / 216 (1.39%)	3 / 215 (1.40%)	
occurrences (all)	3	3	
Macular fibrosis			
subjects affected / exposed	2 / 216 (0.93%)	3 / 215 (1.40%)	
occurrences (all)	2	3	
Retinal haemorrhage			

subjects affected / exposed	2 / 216 (0.93%)	3 / 215 (1.40%)
occurrences (all)	2	3
Vitreous detachment		
subjects affected / exposed	3 / 216 (1.39%)	2 / 215 (0.93%)
occurrences (all)	3	2
Macular degeneration		
subjects affected / exposed	3 / 216 (1.39%)	1 / 215 (0.47%)
occurrences (all)	3	1
Visual impairment		
subjects affected / exposed	1 / 216 (0.46%)	3 / 215 (1.40%)
occurrences (all)	1	3
Eye pain		
subjects affected / exposed	3 / 216 (1.39%)	0 / 215 (0.00%)
occurrences (all)	4	0
Cataract subcapsular		
subjects affected / exposed	1 / 216 (0.46%)	2 / 215 (0.93%)
occurrences (all)	1	2
Choroidal neovascularisation		
subjects affected / exposed	3 / 216 (1.39%)	0 / 215 (0.00%)
occurrences (all)	3	0
Conjunctival hyperaemia		
subjects affected / exposed	2 / 216 (0.93%)	1 / 215 (0.47%)
occurrences (all)	2	1
Cystoid macular oedema		
subjects affected / exposed	1 / 216 (0.46%)	2 / 215 (0.93%)
occurrences (all)	1	2
Eye irritation		
subjects affected / exposed	2 / 216 (0.93%)	1 / 215 (0.47%)
occurrences (all)	2	1
Optic disc haemorrhage		
subjects affected / exposed	3 / 216 (1.39%)	0 / 215 (0.00%)
occurrences (all)	3	0
Blepharitis		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Cataract cortical		

subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences (all)	2	0
Chalazion		
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences (all)	2	0
Corneal erosion		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Detachment of retinal pigment epithelium		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Lacrimation increased		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Retinal exudates		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Vision blurred		
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences (all)	2	0
Vitreous haemorrhage		
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences (all)	2	0
Age-related macular degeneration		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Anterior chamber cell		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Arteriosclerotic retinopathy		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Asthenopia		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1

Conjunctival suffusion		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Cornea verticillata		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Corneal oedema		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Corneal scar		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Dacryostenosis acquired		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Detachment of macular retinal pigment epithelium		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Diplopia		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Dry eye		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Epiretinal membrane		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Eye discharge		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Eye inflammation		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Eye pruritus		

subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Eyelid pain		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Foreign body sensation in eyes		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Macular hole		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Macular oedema		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Macular pseudohole		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Macular scar		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Metamorphopsia		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Optic disc traction syndrome		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Optic nerve sheath haemorrhage		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Photophobia		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Photopsia		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Punctate keratitis		

subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Retinal cyst subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Retinal depigmentation subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Retinal detachment subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Retinal drusen subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Retinal pigment epitheliopathy subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Retinal vein occlusion subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Retinoschisis subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Subretinal fibrosis subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Visual field defect subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Gastrointestinal disorders			
Diarrhoea subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	3 / 215 (1.40%) 3	
Abdominal pain subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	

Abdominal pain upper		
subjects affected / exposed	0 / 216 (0.00%)	2 / 215 (0.93%)
occurrences (all)	0	2
Constipation		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Nausea		
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences (all)	2	0
Abdominal discomfort		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Dental caries		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Diverticulum intestinal		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Gastritis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Gastrointestinal disorder		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Gastrooesophageal reflux disease		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Haemorrhoids		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Inguinal hernia		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Noninfective gingivitis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1

Oesophagitis subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Hepatobiliary disorders			
Cholelithiasis subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	2 / 215 (0.93%) 2	
Cholecystitis chronic subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Cholestasis subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Gallbladder polyp subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Hepatic function abnormal subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Hepatic steatosis subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Skin and subcutaneous tissue disorders			
Pruritus subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	2 / 215 (0.93%) 2	
Rash subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	2 / 215 (0.93%) 2	
Pemphigoid subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 3	
Actinic keratosis subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 2	0 / 215 (0.00%) 0	
Dermatitis			

subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
<b>Dermatitis allergic</b> subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
<b>Eczema</b> subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
<b>Pain of skin</b> subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
<b>Rosacea</b> subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
<b>Scab</b> subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 0	0 / 215 (0.00%) 0	
<b>Skin ulcer</b> subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
<b>Urticaria</b> subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
<b>Petechiae</b> subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
<b>Renal and urinary disorders</b>			
Chronic kidney disease subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	1 / 215 (0.47%) 1	
Renal cyst subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	0 / 215 (0.00%) 0	
Renal failure subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	

Cystitis noninfective		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Haematuria		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Hypertensive nephropathy		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Leukocyturia		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Nephritis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Nephrolithiasis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Nephropathy		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Nephropathy toxic		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Renal impairment		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Stress urinary incontinence		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Urethral pain		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Urethral stenosis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0

Urinary bladder herniation subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Urinary tract discomfort subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Endocrine disorders			
Hypothyroidism subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Hyperthyroidism subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 2	
Musculoskeletal and connective tissue disorders			
Back pain subjects affected / exposed occurrences (all)	4 / 216 (1.85%) 6	5 / 215 (2.33%) 5	
Osteoarthritis subjects affected / exposed occurrences (all)	3 / 216 (1.39%) 3	2 / 215 (0.93%) 2	
Pain in extremity subjects affected / exposed occurrences (all)	3 / 216 (1.39%) 3	2 / 215 (0.93%) 2	
Spinal osteoarthritis subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	4 / 215 (1.86%) 7	
Spinal stenosis subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 3	2 / 215 (0.93%) 2	
Arthralgia subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	2 / 215 (0.93%) 2	
Muscle spasms subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	1 / 215 (0.47%) 1	
Myalgia			

subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Periarthritis		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Intervertebral disc protrusion		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	3	0
Bone pain		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Greater trochanteric pain syndrome		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Intervertebral disc disorder		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Lumbar spinal stenosis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Musculoskeletal stiffness		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Osteoporotic fracture		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Polymyalgia rheumatica		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Rotator cuff syndrome		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Sacral pain		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Spinal pain		

subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Spondylolisthesis subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Sympathetic posterior cervical syndrome subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Tendon disorder subjects affected / exposed occurrences (all)	1 / 216 (0.46%) 1	0 / 215 (0.00%) 0	
Tenosynovitis subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Vertebral foraminal stenosis subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	1 / 215 (0.47%) 1	
Infections and infestations			
Nasopharyngitis subjects affected / exposed occurrences (all)	11 / 216 (5.09%) 11	21 / 215 (9.77%) 23	
COVID-19 subjects affected / exposed occurrences (all)	11 / 216 (5.09%) 12	3 / 215 (1.40%) 3	
Influenza subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	4 / 215 (1.86%) 4	
Urinary tract infection subjects affected / exposed occurrences (all)	3 / 216 (1.39%) 5	1 / 215 (0.47%) 1	
Asymptomatic bacteriuria subjects affected / exposed occurrences (all)	2 / 216 (0.93%) 2	2 / 215 (0.93%) 2	
Cystitis			

subjects affected / exposed	2 / 216 (0.93%)	2 / 215 (0.93%)
occurrences (all)	2	2
Upper respiratory tract infection		
subjects affected / exposed	1 / 216 (0.46%)	3 / 215 (1.40%)
occurrences (all)	1	3
Viral infection		
subjects affected / exposed	2 / 216 (0.93%)	2 / 215 (0.93%)
occurrences (all)	2	2
Otitis externa		
subjects affected / exposed	1 / 216 (0.46%)	2 / 215 (0.93%)
occurrences (all)	1	2
Rhinitis		
subjects affected / exposed	1 / 216 (0.46%)	2 / 215 (0.93%)
occurrences (all)	1	2
Enteritis infectious		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Gastroenteritis viral		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Herpes zoster		
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)
occurrences (all)	1	1
Otitis media		
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences (all)	2	0
Sinusitis		
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)
occurrences (all)	2	0
Erysipelas		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	2	0
Laryngitis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	2	0
Vulval abscess		

subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	2	0
Bronchitis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Hepatitis C		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Lower respiratory tract infection		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Lyme disease		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Onychomycosis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Periodontitis		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Pneumonia aspiration		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Pyelonephritis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Skin infection		
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)
occurrences (all)	0	1
Tonsillitis		
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)
occurrences (all)	1	0
Conjunctivitis		
subjects affected / exposed	6 / 216 (2.78%)	2 / 215 (0.93%)
occurrences (all)	7	2
Hordeolum		

subjects affected / exposed occurrences (all)	0 / 216 (0.00%) 0	2 / 215 (0.93%) 2	
Metabolism and nutrition disorders			
Hypercholesterolaemia			
subjects affected / exposed	2 / 216 (0.93%)	5 / 215 (2.33%)	
occurrences (all)	2	5	
Hypertriglyceridaemia			
subjects affected / exposed	1 / 216 (0.46%)	4 / 215 (1.86%)	
occurrences (all)	1	4	
Diabetes mellitus			
subjects affected / exposed	2 / 216 (0.93%)	1 / 215 (0.47%)	
occurrences (all)	2	1	
Dyslipidaemia			
subjects affected / exposed	2 / 216 (0.93%)	1 / 215 (0.47%)	
occurrences (all)	2	1	
Hypophosphataemia			
subjects affected / exposed	0 / 216 (0.00%)	2 / 215 (0.93%)	
occurrences (all)	0	3	
Hyperkalaemia			
subjects affected / exposed	2 / 216 (0.93%)	0 / 215 (0.00%)	
occurrences (all)	2	0	
Hyperuricaemia			
subjects affected / exposed	1 / 216 (0.46%)	1 / 215 (0.47%)	
occurrences (all)	1	1	
Glucose tolerance impaired			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Gout			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Hyperglycaemia			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Hyperlipidaemia			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	

Hyponatraemia			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Insulin resistance			
subjects affected / exposed	0 / 216 (0.00%)	1 / 215 (0.47%)	
occurrences (all)	0	1	
Iron deficiency			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Lipid metabolism disorder			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	
Type 2 diabetes mellitus			
subjects affected / exposed	1 / 216 (0.46%)	0 / 215 (0.00%)	
occurrences (all)	1	0	

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? Yes

Date	Amendment
28 September 2022	Protocol Amendment 1: The primary change of this amendment is the removal of re-randomization at Week 32, which led to adjustments in the sample size and schedule of assessments.

Notes:

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported