

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
1. - I prefer to utilize a ready to use product as compared to a product to be reconstituted	Strongly agree	n (%)	57 (57.6%)
		95% CI	(0.478, 0.673)
	Agree	n (%)	23 (23.2%)
		95% CI	(0.149, 0.316)
	Neither agree nor disagree	n (%)	19 (19.2%)
		95% CI	(0.114, 0.269)
	Disagree	n (%)	0
		95% CI	(NA, NA)
	Strongly disagree	n (%)	0
		95% CI	(NA, NA)
2. - I could dedicate more time to explain the treatment procedure in using a ready to use product as compared to a product to be reconstituted	Strongly agree	n (%)	49 (49.5%)
		95% CI	(0.396, 0.593)
	Agree	n (%)	47 (47.5%)
		95% CI	(0.376, 0.573)
	Neither agree nor disagree	n (%)	2 (2.0%)
		95% CI	(0.000, 0.048)
	Disagree	n (%)	1 (1.0%)
		95% CI	(0.000, 0.030)
	Strongly disagree	n (%)	0
		95% CI	(NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
3. - I could save time on the reconstitution and injection procedure to do something else	Strongly agree	n (%)	56 (56.6%)
		95% CI	(0.468, 0.663)
	Agree	n (%)	36 (36.4%)
		95% CI	(0.269, 0.458)
	Neither agree nor disagree	n (%)	7 (7.1%)
		95% CI	(0.020, 0.121)
	Disagree	n (%)	0
		95% CI	(NA, NA)
	Strongly disagree	n (%)	0
		95% CI	(NA, NA)
4. - I felt more relaxed/less stressed in not having to reconstitute the product to be injected	Strongly agree	n (%)	52 (52.5%)
		95% CI	(0.427, 0.624)
	Agree	n (%)	16 (16.2%)
		95% CI	(0.089, 0.234)
	Neither agree nor disagree	n (%)	24 (24.2%)
		95% CI	(0.158, 0.327)
	Disagree	n (%)	2 (2.0%)
		95% CI	(0.000, 0.048)
	Strongly disagree	n (%)	5 (5.1%)
		95% CI	(0.007, 0.094)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
5. - I feel more secure when I have a pre-diluted solution and don't have to reconstitute the product myself	Strongly agree	n (%)	53 (53.5%)
		95% CI	(0.437, 0.634)
	Agree	n (%)	11 (11.1%)
		95% CI	(0.049, 0.173)
	Neither agree nor disagree	n (%)	25 (25.3%)
		95% CI	(0.167, 0.338)
	Disagree	n (%)	4 (4.0%)
		95% CI	(0.002, 0.079)
	Strongly disagree	n (%)	6 (6.1%)
		95% CI	(0.014, 0.108)
6. - I have a good feeling about injecting a modern and innovative product as compared to the reconstituted product	Strongly agree	n (%)	42 (42.4%)
		95% CI	(0.327, 0.522)
	Agree	n (%)	39 (39.4%)
		95% CI	(0.298, 0.490)
	Neither agree nor disagree	n (%)	17 (17.2%)
		95% CI	(0.097, 0.246)
	Disagree	n (%)	1 (1.0%)
		95% CI	(0.000, 0.030)
	Strongly disagree	n (%)	0
		95% CI	(NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
7. - I feel I am more precise in my injection with a ready to use product compared to a product to bereconstituted	Strongly agree	n (%)	50 (50.5%)
		95% CI	(0.407, 0.604)
	Agree	n (%)	25 (25.3%)
		95% CI	(0.167, 0.338)
	Neither agree nor disagree	n (%)	12 (12.1%)
		95% CI	(0.057, 0.186)
	Disagree	n (%)	8 (8.1%)
		95% CI	(0.027, 0.134)
	Strongly disagree	n (%)	4 (4.0%)
		95% CI	(0.002, 0.079)
8. - I like to propose new innovative treatment with liquid ready to use product to my patient	Strongly agree	n (%)	43 (43.4%)
		95% CI	(0.337, 0.532)
	Agree	n (%)	43 (43.4%)
		95% CI	(0.337, 0.532)
	Neither agree nor disagree	n (%)	11 (11.1%)
		95% CI	(0.049, 0.173)
	Disagree	n (%)	2 (2.0%)
		95% CI	(0.000, 0.048)
	Strongly disagree	n (%)	0
		95% CI	(NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question	Statistics	Alluzience (N=99)	
9. - I produced substantially less waste of non-toxin material (syringes, needles, saline etc.) using ready to use product as compared to the reconstituted product	Strongly agree	n (%) 75 (75.8%) 95% CI (0.673, 0.842)	
	Agree	n (%) 21 (21.2%) 95% CI (0.132, 0.293)	
	Neither agree nor disagree	n (%) 3 (3.0%) 95% CI (0.000, 0.064)	
	Disagree	n (%) 0 95% CI (NA, NA)	
	Strongly disagree	n (%) 0 95% CI (NA, NA)	
	10. - When I use a ready to use toxin I spend less materials for injection	Strongly agree	n (%) 84 (84.8%) 95% CI (0.778, 0.919)
		Agree	n (%) 14 (14.1%) 95% CI (0.073, 0.210)
		Neither agree nor disagree	n (%) 1 (1.0%) 95% CI (0.000, 0.030)
		Disagree	n (%) 0 95% CI (NA, NA)
		Strongly disagree	n (%) 0 95% CI (NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
11. - I believe that the use of ready to use toxin is better for the environment as I produce less waste	Strongly agree	n (%)	78 (78.8%)
		95% CI	(0.707, 0.868)
	Agree	n (%)	21 (21.2%)
		95% CI	(0.132, 0.293)
	Neither agree nor disagree	n (%)	0
		95% CI	(NA, NA)
	Disagree	n (%)	0
		95% CI	(NA, NA)
	Strongly disagree	n (%)	0
		95% CI	(NA, NA)
12. - I feel more confident when injecting a product free from animal and human excipients	Strongly agree	n (%)	67 (67.7%)
		95% CI	(0.585, 0.769)
	Agree	n (%)	15 (15.2%)
		95% CI	(0.081, 0.222)
	Neither agree nor disagree	n (%)	17 (17.2%)
		95% CI	(0.097, 0.246)
	Disagree	n (%)	0
		95% CI	(NA, NA)
	Strongly disagree	n (%)	0
		95% CI	(NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
1. - I prefer to utilize a ready to use product as compared to a product to be reconstituted	Strongly agree/Agree	n (%)	80 (80.8%)
		95% CI	(0.731, 0.886)
	Neither agree nor disagree	n (%)	19 (19.2%)
		95% CI	(0.114, 0.269)
	Disagree/Strongly disagree	n (%)	0
		95% CI	(NA, NA)
2. - I could dedicate more time to explain the treatment procedure in using a ready to use product as compared to a product to be reconstituted	Strongly agree/Agree	n (%)	96 (97.0%)
		95% CI	(0.936, 1.000)
	Neither agree nor disagree	n (%)	2 (2.0%)
		95% CI	(0.000, 0.048)
	Disagree/Strongly disagree	n (%)	1 (1.0%)
		95% CI	(0.000, 0.030)
3. - I could save time on the reconstitution and injection procedure to do something else	Strongly agree/Agree	n (%)	92 (92.9%)
		95% CI	(0.879, 0.980)
	Neither agree nor disagree	n (%)	7 (7.1%)
		95% CI	(0.020, 0.121)
	Disagree/Strongly disagree	n (%)	0
		95% CI	(NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
4. - I felt more relaxed/less stressed in not having to reconstitute the product to be injected	Strongly agree/Agree	n (%)	68 (68.7%)
		95% CI	(0.596, 0.778)
	Neither agree nor disagree	n (%)	24 (24.2%)
		95% CI	(0.158, 0.327)
	Disagree/Strongly disagree	n (%)	7 (7.1%)
		95% CI	(0.020, 0.121)
5. - I feel more secure when I have a pre-diluted solution and don't have to reconstitute the product myself	Strongly agree/Agree	n (%)	64 (64.6%)
		95% CI	(0.552, 0.741)
	Neither agree nor disagree	n (%)	25 (25.3%)
		95% CI	(0.167, 0.338)
	Disagree/Strongly disagree	n (%)	10 (10.1%)
		95% CI	(0.042, 0.160)
6. - I have a good feeling about injecting a modern and innovative product as compared to the reconstituted product	Strongly agree/Agree	n (%)	81 (81.8%)
		95% CI	(0.742, 0.894)
	Neither agree nor disagree	n (%)	17 (17.2%)
		95% CI	(0.097, 0.246)
	Disagree/Strongly disagree	n (%)	1 (1.0%)
		95% CI	(0.000, 0.030)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question	Statistics	Alluzience (N=99)
7. - I feel I am more precise in my injection with a ready to use product compared to a product to bereconstituted	Strongly agree/Agree	n (%)
		75 (75.8%)
		95% CI
		(0.673, 0.842)
	Neither agree nor disagree	n (%)
		12 (12.1%)
8. - I like to propose new innovative treatment with liquid ready to use product to my patient		95% CI
		(0.057, 0.186)
	Disagree/Strongly disagree	n (%)
		12 (12.1%)
		95% CI
		(0.057, 0.186)
9. - I produced substantially less waste of non-toxin material (syringes, needles, saline etc.) using ready to use product as compared to the reconstituted product	Strongly agree/Agree	n (%)
		86 (86.9%)
		95% CI
		(0.802, 0.935)
	Neither agree nor disagree	n (%)
		11 (11.1%)
9. - I produced substantially less waste of non-toxin material (syringes, needles, saline etc.) using ready to use product as compared to the reconstituted product		95% CI
		(0.049, 0.173)
	Disagree/Strongly disagree	n (%)
		2 (2.0%)
		95% CI
		(0.000, 0.048)
9. - I produced substantially less waste of non-toxin material (syringes, needles, saline etc.) using ready to use product as compared to the reconstituted product	Strongly agree/Agree	n (%)
		96 (97.0%)
		95% CI
		(0.936, 1.000)
	Neither agree nor disagree	n (%)
		3 (3.0%)
9. - I produced substantially less waste of non-toxin material (syringes, needles, saline etc.) using ready to use product as compared to the reconstituted product		95% CI
		(0.000, 0.064)
	Disagree/Strongly disagree	n (%)
		0
		95% CI
		(NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.

Table 14.2.6.1
 Investigator Treatment Session Questionnaire (Alluzience Only)
 FAS Population

Question		Statistics	Alluzience (N=99)
10. - When I use a ready to use toxin I spend less materials for injection	Strongly agree/Agree	n (%)	98 (99.0%)
		95% CI	(0.970, 1.000)
	Neither agree nor disagree	n (%)	1 (1.0%)
		95% CI	(0.000, 0.030)
	Disagree/Strongly disagree	n (%)	0
		95% CI	(NA, NA)
11. - I believe that the use of ready to use toxin is better for the environment as I produce less waste	Strongly agree/Agree	n (%)	99 (100%)
		95% CI	(1.000, 1.000)
	Neither agree nor disagree	n (%)	0
		95% CI	(NA, NA)
	Disagree/Strongly disagree	n (%)	0
		95% CI	(NA, NA)
12. - I feel more confident when injecting a product free from animal and human excipients	Strongly agree/Agree	n (%)	82 (82.8%)
		95% CI	(0.754, 0.903)
	Neither agree nor disagree	n (%)	17 (17.2%)
		95% CI	(0.097, 0.246)
	Disagree/Strongly disagree	n (%)	0
		95% CI	(NA, NA)

Abbreviation: FAS=Full Analysis Set;
 Notes: Percentages are based on the FAS population.
 The FAS population is defined as all randomized and treated subjects.