



## Clinical trial results:

### A Randomized, Double-Blind, Placebo-Controlled, Phase 3 Trial to Investigate the Safety and Immunogenicity of a Dengue Tetravalent Vaccine (Live, Attenuated) (TDV) Administered Subcutaneously to Healthy Subjects Aged 4 to 60 Years in India

#### Summary

EudraCT number	2023-000134-15
Trial protocol	Outside EU/EEA
Global end of trial date	05 May 2025

#### Results information

Result version number	v2 (current)
This version publication date	20 May 2026
First version publication date	21 November 2025
Version creation reason	

#### Trial information

##### Trial identification

Sponsor protocol code	DEN-302
-----------------------	---------

##### Additional study identifiers

ISRCTN number	-
ClinicalTrials.gov id (NCT number)	NCT06060067
WHO universal trial number (UTN)	-

Notes:

#### Sponsors

Sponsor organisation name	Takeda Vaccines, Inc.
Sponsor organisation address	40 Landsdowne Street, Cambridge, MA, United States, 02139
Public contact	Study Director, Takeda, TrialDisclosures@takeda.com
Scientific contact	Study Director, Takeda, TrialDisclosures@takeda.com

Notes:

#### Paediatric regulatory details

Is trial part of an agreed paediatric investigation plan (PIP)	No
Does article 45 of REGULATION (EC) No 1901/2006 apply to this trial?	No
Does article 46 of REGULATION (EC) No 1901/2006 apply to this trial?	Yes

Notes:

## Results analysis stage

Analysis stage	Final
Date of interim/final analysis	05 May 2025
Is this the analysis of the primary completion data?	No
Global end of trial reached?	Yes
Global end of trial date	05 May 2025
Was the trial ended prematurely?	No

Notes:

## General information about the trial

Main objective of the trial:

The main objective of the trial was to evaluate the immunogenicity and safety of TDV in the Indian population.

Protection of trial subjects:

Each participant or their legally authorized representative signed an informed consent form (ICF).

Background therapy:

NA

Evidence for comparator:

NA

Actual start date of recruitment	29 March 2024
Long term follow-up planned	No
Independent data monitoring committee (IDMC) involvement?	Yes

Notes:

## Population of trial subjects

### Subjects enrolled per country

Country: Number of subjects enrolled	India: 480
Worldwide total number of subjects	480
EEA total number of subjects	0

Notes:

### Subjects enrolled per age group

In utero	0
Preterm newborn - gestational age < 37 wk	0
Newborns (0-27 days)	0
Infants and toddlers (28 days-23 months)	0
Children (2-11 years)	141
Adolescents (12-17 years)	99
Adults (18-64 years)	240
From 65 to 84 years	0
85 years and over	0

## Subject disposition

### Recruitment

Recruitment details:

Participants took part in the study at 10 investigative sites in India from 29 March 2024 to 05 May 2025.

### Pre-assignment

Screening details:

Healthy children and adult participants were enrolled in Cohort 1 ( $\geq 18$  to  $\leq 60$  Age Group) and Cohort 2 ( $\geq 4$  to  $< 18$  Age Group) to receive either Dengue Tetravalent Vaccine (TDV) or placebo in a 3:1 ratio.

### Period 1

Period 1 title	Overall Study (overall period)
Is this the baseline period?	Yes
Allocation method	Randomised - controlled
Blinding used	Double blind
Roles blinded	Subject, Investigator, Monitor, Data analyst

### Arms

Are arms mutually exclusive?	Yes
<b>Arm title</b>	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: TDV

Arm description:

Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Arm type	Experimental
Investigational medicinal product name	Dengue Tetravalent Vaccine (Live, Attenuated)
Investigational medicinal product code	
Other name	TAK-003
Pharmaceutical forms	Powder and solvent for solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

TDV SC injection on Day 1 and Day 90 of the study.

<b>Arm title</b>	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: Placebo
------------------	---

Arm description:

Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Arm type	Placebo
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Normal Saline (0.9% NaCl) SC injection on Day 1 and Day 90 of the study.

<b>Arm title</b>	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: TDV
------------------	---

Arm description:

Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Arm type	Experimental
----------	--------------

Investigational medicinal product name	Dengue Tetravalent Vaccine (Live, Attenuated)
Investigational medicinal product code	
Other name	TAK-003
Pharmaceutical forms	Powder and solvent for solution for injection
Routes of administration	Subcutaneous use
Dosage and administration details: TDV SC injection on Day 1 and Day 90 of the study.	
<b>Arm title</b>	Cohort 2: ≥4 to <18 Years Age Group: Placebo

Arm description:

Participants received placebo-matching TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Arm type	Placebo
Investigational medicinal product name	Placebo
Investigational medicinal product code	
Other name	
Pharmaceutical forms	Solution for injection
Routes of administration	Subcutaneous use

Dosage and administration details:

Normal Saline (0.9% NaCl) SC injection on Day 1 and Day 90 of the study.

<b>Number of subjects in period 1</b>	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV
Started	180	60	180
Completed	175	59	178
Not completed	5	1	2
Consent withdrawn by subject	2	-	-
Adverse event, non-fatal	1	-	-
Reason Not Specified	1	-	1
Lost to follow-up	1	1	1

<b>Number of subjects in period 1</b>	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Started	60
Completed	58
Not completed	2
Consent withdrawn by subject	1
Adverse event, non-fatal	-
Reason Not Specified	-
Lost to follow-up	1

## Baseline characteristics

### Reporting groups

Reporting group title	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: TDV
Reporting group description: Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Reporting group title	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: Placebo
Reporting group description: Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Reporting group title	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: TDV
Reporting group description: Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Reporting group title	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: Placebo
Reporting group description: Participants received placebo-matching TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	

Reporting group values	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: TDV	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: Placebo	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: TDV
Number of subjects	180	60	180
Age Categorical Units: Subjects			

Age continuous Units: years			
arithmetic mean	37.2	36.5	9.7
standard deviation	$\pm 11.66$	$\pm 11.12$	$\pm 4.05$
Gender categorical Units: Subjects			
Female	54	14	79
Male	126	46	101
Race (NIH/OMB) Units: Subjects			
American Indian or Alaska Native	0	0	0
Asian	180	60	180
Native Hawaiian or Other Pacific Islander	0	0	0
Black or African American	0	0	0
White	0	0	0
More than one race	0	0	0
Unknown or Not Reported	0	0	0
Ethnicity (NIH/OMB) Units: Subjects			
Hispanic or Latino	0	0	0
Not Hispanic or Latino	180	60	180
Unknown or Not Reported	0	0	0

Reporting group values	Cohort 2: $\geq 4$ to $< 18$ Years Age Group:	Total	
------------------------	---	-------	--

Placebo

Number of subjects	60	480	
Age Categorical Units: Subjects			
Age continuous Units: years arithmetic mean standard deviation	10.3 ± 4.35	-	
Gender categorical Units: Subjects			
Female	29	176	
Male	31	304	
Race (NIH/OMB) Units: Subjects			
American Indian or Alaska Native	0	0	
Asian	59	479	
Native Hawaiian or Other Pacific Islander	0	0	
Black or African American	0	0	
White	0	0	
More than one race	0	0	
Unknown or Not Reported	1	1	
Ethnicity (NIH/OMB) Units: Subjects			
Hispanic or Latino	0	0	
Not Hispanic or Latino	59	479	
Unknown or Not Reported	1	1	

## End points

### End points reporting groups

Reporting group title	Cohort 1: ≥18 to ≤60 Years Age Group: TDV
Reporting group description: Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Reporting group title	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo
Reporting group description: Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Reporting group title	Cohort 2: ≥4 to <18 Years Age Group: TDV
Reporting group description: Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Reporting group title	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Reporting group description: Participants received placebo-matching TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Subject analysis set title	Cohort 2: ≥4 to <6 Years Age group: TDV
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Subject analysis set title	Cohort 2: ≥4 to <6 Years Age group: Placebo
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Subject analysis set title	Cohort 2: ≥6 to <18 Years Age group: TDV
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Subject analysis set title	Cohort 2: ≥6 to <18 Years Age group: Placebo
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Subject analysis set title	Cohort 1 and 2 Combined: TDV
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	
Subject analysis set title	Cohort 1 and 2 Combined: Placebo
Subject analysis set type	Sub-group analysis
Subject analysis set description: Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.	

### Primary: Number of Participants with Solicited Local (Injection Site) Adverse Events (AEs) by Severity Within 7 Days Post-vaccination at Day 1 for Age Group 6 Years and Over

End point title	Number of Participants with Solicited Local (Injection Site) Adverse Events (AEs) by Severity Within 7 Days Post-vaccination at Day 1 for Age Group 6 Years and Over <sup>[1][2]</sup>
End point description: Solicited local AEs at injection site were defined as injection site pain, injection site erythema, and	

injection site swelling. The AEs were graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe based on solicited safety parameters defined for child/adolescent/adult ( $\geq 6$  years of age). Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited local AEs were graded separately for the subgroups of age  $< 6$  years old and age  $\geq 6$  years old to  $< 18$  years old. Data for subgroup of age  $\geq 6$  years old to  $< 18$  years old are reported in this outcome measure along with the data for  $\geq 18$  to  $\leq 60$  age group. Subjects analysed are the number of participants with data available for analysis.

End point type	Primary
----------------	---------

End point timeframe:

Within 7 days postvaccination at Day 1

Notes:

[1] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

[2] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Descriptive analysis for this end point was planned only for the arm groups where it is reported.

End point values	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: TDV	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: Placebo	Cohort 2: $\geq 6$ to $< 18$ Years Age group: TDV	Cohort 2: $\geq 6$ to $< 18$ Years Age group: Placebo
Subject group type	Reporting group	Reporting group	Subject analysis set	Subject analysis set
Number of subjects analysed	179	60	133	46
Units: participants				
number (not applicable)				
Pain: None	152	53	119	42
Pain: Mild	25	7	13	4
Pain: Moderate	2	0	1	0
Pain: Severe	0	0	0	0
Erythema: None	177	60	133	46
Erythema: Mild	2	0	0	0
Erythema: Moderate	0	0	0	0
Erythema: Severe	0	0	0	0
Swelling: None	179	60	132	46
Swelling: Mild	0	0	1	0
Swelling: Moderate	0	0	0	0
Swelling: Severe	0	0	0	0

## Statistical analyses

No statistical analyses for this end point

## Primary: Number of Participants with Solicited Local (Injection Site) Adverse Events (AEs) by Severity Within 7 Days Post-vaccination at Day 1 for Age Group Less Than 6 Years

End point title	Number of Participants with Solicited Local (Injection Site) Adverse Events (AEs) by Severity Within 7 Days Post-vaccination at Day 1 for Age Group Less Than 6 Years <sup>[3]</sup>
-----------------	--

End point description:

Solicited local AEs at injection site were defined as injection site pain, injection site erythema, and injection site swelling. The AEs were graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe based on solicited safety parameters defined for infant/toddler/child ( $< 6$  years of age). Safety Analysis Set included all randomized participants who received at least 1 dose of



IP. As prespecified, solicited local AEs were graded separately for the subgroups of age <6 years old and age ≥6 years old to <18 years old. Data for subgroup of age <6 years old are reported in this outcome measure.

End point type	Primary
End point timeframe:	
Within 7 days post-vaccination at Day 1	

Notes:

[3] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 2: ≥4 to <6 Years Age group: TDV	Cohort 2: ≥4 to <6 Years Age group: Placebo		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	46	14		
Units: participants				
Pain: None	42	10		
Pain: Mild	4	4		
Pain: Moderate	0	0		
Pain: Severe	0	0		
Erythema: None	46	14		
Erythema: Mild	0	0		
Erythema: Moderate	0	0		
Erythema: Severe	0	0		
Swelling: None	45	14		
Swelling: Mild	1	0		
Swelling: Moderate	0	0		
Swelling: Severe	0	0		

## Statistical analyses

No statistical analyses for this end point

### Primary: Number of Participants With Solicited Local (Injection Site) AEs by Severity Within 7 Days Post-vaccination at Day 90 for Age Group Less Than 6 Years

End point title	Number of Participants With Solicited Local (Injection Site) AEs by Severity Within 7 Days Post-vaccination at Day 90 for Age Group Less Than 6 Years <sup>[4]</sup>
-----------------	--

End point description:

Solicited local AEs at injection site were defined as injection site pain, injection site erythema, and injection site swelling. The AEs were graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe based on solicited safety parameters defined for infant/toddler/child (<6 years of age). Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited local AEs were graded separately for the subgroups of age <6 years old and age ≥6 years old to <18 years old. Data for subgroup of age <6 years old are reported in this outcome measure.

End point type	Primary
End point timeframe:	
Within 7 days post-vaccination at Day 90	

Notes:

[4] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 2: ≥4 to <6 Years Age group: TDV	Cohort 2: ≥4 to <6 Years Age group: Placebo		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	46	14		
Units: participants				
Pain: None	44	14		
Pain: Mild	2	0		
Pain: Moderate	0	0		
Pain: Severe	0	0		
Erythema: None	46	14		
Erythema: Mild	0	0		
Erythema: Moderate	0	0		
Erythema: Severe	0	0		
Swelling: None	46	14		
Swelling: Mild	0	0		
Swelling: Moderate	0	0		
Swelling: Severe	0	0		

## Statistical analyses

No statistical analyses for this end point

## Primary: Number of Participants with Solicited Local (Injection Site) AEs by Severity Within 7 Days Post-vaccination at Day 90 for Age Group 6 Years and Over

End point title	Number of Participants with Solicited Local (Injection Site) AEs by Severity Within 7 Days Post-vaccination at Day 90 for Age Group 6 Years and Over <sup>[5][6]</sup>
-----------------	--

End point description:

Solicited local AEs at injection site were defined as injection site pain, injection site erythema, and injection site swelling. The AEs were graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe based on solicited safety parameters defined for child/adolescent/adult (≥6 years of age). Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited local AEs were graded separately for the subgroups of age <6 years old and age ≥6 years old to <18 years old. Data for subgroup of age ≥6 years old to <18 years old are reported in this outcome measure along with the data for ≥18 to ≤60 age group. Subjects analysed are the number of participants with data available for analysis.

End point type	Primary
----------------	---------

End point timeframe:

Within 7 days postvaccination at Day 90

Notes:

[5] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

[6] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Descriptive analysis for this end point was planned only for the arm groups where it is reported.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥6 to <18 Years Age group: TDV	Cohort 2: ≥6 to <18 Years Age group: Placebo
Subject group type	Reporting group	Reporting group	Subject analysis set	Subject analysis set
Number of subjects analysed	176	59	132	45
Units: participants				
number (not applicable)				
Pain: None	163	55	117	41
Pain: Mild	12	3	14	4
Pain: Moderate	1	1	1	0
Pain: Severe	0	0	0	0
Erythema: None	176	59	132	45
Erythema: Mild	0	0	0	0
Erythema: Moderate	0	0	0	0
Erythema: Severe	0	0	0	0
Swelling: None	176	59	132	45
Swelling: Mild	0	0	0	0
Swelling: Moderate	0	0	0	0
Swelling: Severe	0	0	0	0

## Statistical analyses

No statistical analyses for this end point

## Primary: Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 1 for Age Group Less Than 6 Years

End point title	Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 1 for Age Group Less Than 6 Years <sup>[7]</sup>
-----------------	--

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Solicited systemic AEs assessments for children <6 years old comprised: irritability/fussiness, drowsiness, loss of appetite (graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe) and fever in °C (collected by temperature ranges as indicated). Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited systemic AEs were graded separately for the subgroups of age <6 years old and age ≥6 years old to <18 years old. Data for subgroup of age <6 years old are reported in this outcome measure.

End point type	Primary
----------------	---------

End point timeframe:

Within 14 days postvaccination at Day 1

Notes:

[7] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 2: ≥4 to <6 Years Age group: TDV	Cohort 2: ≥4 to <6 Years Age group: Placebo		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	46	14		
Units: participants				
number (not applicable)				
Irritability or Fussiness: None	45	13		
Irritability or Fussiness: Mild	1	1		
Irritability or Fussiness: Moderate	0	0		
Irritability or Fussiness: Severe	0	0		
Drowsiness: None	45	13		
Drowsiness: Mild	1	1		
Drowsiness: Moderate	0	0		
Drowsiness: Severe	0	0		
Loss of Appetite: None	46	14		
Loss of Appetite: Mild	0	0		
Loss of Appetite: Moderate	0	0		
Loss of Appetite: Severe	0	0		
Fever: None	46	14		
Fever: 38.0-<38.5 °C	0	0		
Fever: 38.5-<39.0 °C	0	0		
Fever: 39.0-<39.5 °C	0	0		
Fever: 39.5-<40.0 °C	0	0		
Fever: 40.0-<40.5 °C	0	0		
Fever: 40.5-<41.0 °C	0	0		
Fever: ≥41.0 °C	0	0		

## Statistical analyses

No statistical analyses for this end point

## Primary: Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 1 for Age Group 6 Years and Over

End point title	Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 1 for Age Group 6 Years and Over <sup>[8][9]</sup>
-----------------	--

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Solicited systemic AEs assessments for children ≥ 6 years old/adolescents/adults comprised: headache, asthenia, malaise, myalgia (graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe) and fever in °C (collected by temperature ranges as indicated). Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited systemic AEs were graded separately for the subgroups of age <6 years old and age ≥ 6 years old to < 18 years old. Data for subgroup of age ≥ 6 years old to < 18 years old are reported in this outcome measure along with the data for ≥18 to ≤60 age group. Subjects analysed are the number of participants with data available for analysis.

End point type	Primary
----------------	---------

End point timeframe:

Within 14 days postvaccination at Day 1

Notes:

[8] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

[9] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Descriptive analysis for this end point was planned only for the arm groups where it is reported.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥6 to <18 Years Age group: TDV	Cohort 2: ≥6 to <18 Years Age group: Placebo
Subject group type	Reporting group	Reporting group	Subject analysis set	Subject analysis set
Number of subjects analysed	179	60	133	46
Units: participants				
number (not applicable)				
Headache: None	165	54	127	45
Headache: Mild	11	6	6	1
Headache: Moderate	1	0	0	0
Headache: Severe	2	0	0	0
Asthenia: None	168	56	126	44
Asthenia: Mild	11	4	6	2
Asthenia: Moderate	0	0	1	0
Asthenia: Severe	0	0	0	0
Malaise: None	174	57	130	46
Malaise: Mild	5	3	2	0
Malaise: Moderate	0	0	1	0
Malaise: Severe	0	0	0	0
Myalgia: None	168	56	128	45
Myalgia: Mild	10	3	4	1
Myalgia: Moderate	1	1	1	0
Myalgia: Severe	0	0	0	0
Fever: None	178	58	131	45
Fever: 38.0-<38.5 °C	0	2	1	0
Fever: 38.5-<39.0 °C	0	0	0	0
Fever: 39.0-<39.5 °C	0	0	0	1
Fever: 39.5-<40.0 °C	0	0	0	0
Fever: 40.0-<40.5 °C	1	0	1	0
Fever: 40.5-<41.0 °C	0	0	0	0
Fever: ≥41.0 °C	0	0	0	0

## Statistical analyses

No statistical analyses for this end point

## Primary: Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 90 for Age Group Less Than 6 Years

End point title	Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 90 for Age Group Less Than 6 Years <sup>[10]</sup>
-----------------	--

**End point description:**

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Solicited systemic AEs assessments for children <6 years old comprised: drowsiness, irritability/fussiness, loss of appetite (graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe). Data for drowsiness, irritability/fussiness and loss of appetite has been reported in this outcome measure. Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited systemic AEs were graded separately for the subgroups of age <6 years old and age ≥6 years old to <18 years old. Data for subgroup of age <6 years old are reported in this outcome measure. Subjects analysed are the number of participants with data available for analysis for this outcome measure.

End point type	Primary
----------------	---------

**End point timeframe:**

Within 14 days post-vaccination at Day 90

**Notes:**

[10] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 2: ≥4 to <6 Years Age group: TDV	Cohort 2: ≥4 to <6 Years Age group: Placebo		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	38	14		
Units: participants				
number (not applicable)				
Irritability or Fussiness: None	38	14		
Irritability or Fussiness: Mild	0	0		
Irritability or Fussiness: Moderate	0	0		
Irritability or Fussiness: Severe	0	0		
Drowsiness: None	37	13		
Drowsiness: Mild	1	1		
Drowsiness: Moderate	0	0		
Drowsiness: Severe	0	0		
Loss of Appetite: None	38	13		
Loss of Appetite: Mild	0	1		
Loss of Appetite: Moderate	0	0		
Loss of Appetite: Severe	0	0		

**Statistical analyses**

No statistical analyses for this end point

**Primary: Number of Participants With Solicited Systemic AEs of Fever by Severity Within 14 Days Post-vaccination at Day 90 for Age Group Less Than 6 Years**

End point title	Number of Participants With Solicited Systemic AEs of Fever by Severity Within 14 Days Post-vaccination at Day 90 for Age Group Less Than 6 Years <sup>[11]</sup>
-----------------	---

**End point description:**

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Solicited systemic AEs assessments for children <6 years old also comprised of fever in °C (collected by temperature ranges as indicated). Data for fever has been reported in this outcome measure. Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited

systemic AEs were graded separately for the subgroups of age <6 years old and age ≥6 years old to <18 years old. Data for subgroup of age <6 years old are reported in this outcome measure.

End point type	Primary
----------------	---------

End point timeframe:

Within 14 days post-vaccination at Day 90

Notes:

[11] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 2: ≥4 to <6 Years Age group: TDV	Cohort 2: ≥4 to <6 Years Age group: Placebo		
Subject group type	Subject analysis set	Subject analysis set		
Number of subjects analysed	46	14		
Units: participants				
number (not applicable)				
Fever: None	46	14		
Fever: 38.0-<38.5 °C	0	0		
Fever: 38.5-<39.0 °C	0	0		
Fever: 39.0-<39.5 °C	0	0		
Fever: 39.5-<40.0 °C	0	0		
Fever: 40.0-<40.5 °C	0	0		
Fever: 40.5-<41.0 °C	0	0		
Fever: ≥41.0 °C	0	0		

## Statistical analyses

No statistical analyses for this end point

## Primary: Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 90 for Age Group 6 Years and Over

End point title	Number of Participants with Solicited Systemic AEs by Severity Within 14 Days Post-vaccination at Day 90 for Age Group 6 Years and Over <sup>[12][13]</sup>
-----------------	---

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Solicited systemic AEs assessments for children ≥ 6 years old/adolescents/adults comprised: headache, asthenia, malaise, myalgia (graded by investigator as Grade 0: none, Grade 1: mild, Grade 2: moderate and Grade 3: severe) and fever in °C (collected by temperature ranges as indicated). Safety Analysis Set included all randomized participants who received at least 1 dose of IP. As prespecified, solicited systemic AEs were graded separately for the subgroups of age <6 years old and age ≥6 years old to <18 years old. Data for subgroup of age ≥6 years old to <18 years old are reported in this outcome measure along with the data for ≥18 to ≤60 age group. Subjects analysed are the number of participants with data available for analysis.

End point type	Primary
----------------	---------

End point timeframe:

Within 14 days postvaccination at Day 90

Notes:

[12] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

[13] - The end point is not reporting statistics for all the arms in the baseline period. It is expected all the baseline period arms will be reported on when providing values for an end point on the baseline period.

Justification: Descriptive analysis for this end point was planned only for the arm groups where it is reported.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥6 to <18 Years Age group: TDV	Cohort 2: ≥6 to <18 Years Age group: Placebo
Subject group type	Reporting group	Reporting group	Subject analysis set	Subject analysis set
Number of subjects analysed	176	59	132	45
Units: participants				
number (not applicable)				
Headache: None	169	59	129	45
Headache: Mild	6	0	3	0
Headache: Moderate	1	0	0	0
Headache: Severe	0	0	0	0
Asthenia: None	172	58	129	44
Asthenia: Mild	3	1	3	1
Asthenia: Moderate	1	0	0	0
Asthenia: Severe	0	0	0	0
Malaise: None	173	59	132	45
Malaise: Mild	3	0	0	0
Malaise: Moderate	0	0	0	0
Malaise: Severe	0	0	0	0
Myalgia: None	173	59	129	44
Myalgia: Mild	1	0	3	1
Myalgia: Moderate	2	0	0	0
Myalgia: Severe	0	0	0	0
Fever: None	172	57	129	44
Fever: 38.0-<38.5 °C	3	1	2	0
Fever: 38.5-<39.0 °C	1	1	0	1
Fever: 39.0-<39.5 °C	0	0	0	0
Fever: 39.5-<40.0 °C	0	0	0	0
Fever: 40.0-<40.5 °C	0	0	1	0
Fever: 40.5-<41.0 °C	0	0	0	0
Fever: ≥41.0 °C	0	0	0	0

## Statistical analyses

No statistical analyses for this end point

## Primary: Percentage of Participants with Any Unsolicited AEs Within 28 Days Post-vaccination at Day 1

End point title	Percentage of Participants with Any Unsolicited AEs Within 28 Days Post-vaccination at Day 1 <sup>[14]</sup>
-----------------	--

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Safety Analysis Set included all randomized participants who received at least 1 dose of IP.

End point type	Primary
----------------	---------



End point timeframe:

Within 28 days postvaccination at Day 1

Notes:

[14] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	180	60	180	60
Units: percentage of participants				
number (not applicable)	2.2	1.7	2.2	1.7

## Statistical analyses

No statistical analyses for this end point

## Primary: Percentage of Participants with Any Unsolicited AEs Within 28 Days Post-vaccination at Day 90

End point title	Percentage of Participants with Any Unsolicited AEs Within 28 Days Post-vaccination at Day 90 <sup>[15]</sup>
-----------------	---

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Safety Analysis Set included all randomized participants who received at least 1 dose of IP.

End point type	Primary
----------------	---------

End point timeframe:

Within 28 days postvaccination at Day 90

Notes:

[15] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	180	60	180	60
Units: percentage of participants				
number (not applicable)	1.7	3.4	4.5	3.4

## Statistical analyses

No statistical analyses for this end point

## Primary: Percentage of Participants with an AE Leading to Participant Withdrawal

**from Trial**

End point title	Percentage of Participants with an AE Leading to Participant Withdrawal from Trial <sup>[16]</sup>
-----------------	--

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Safety Analysis Set included all randomized participants who received at least 1 dose of IP.

End point type	Primary
----------------	---------

End point timeframe:

From first vaccination on Day 1 through the end of trial (up to Day 270)

Notes:

[16] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	180	60	180	60
Units: percentage of participants				
number (not applicable)	0.6	0	0	0

**Statistical analyses**

No statistical analyses for this end point

**Primary: Percentage of Participants with an AE Leading to TDV or Placebo Discontinuation**

End point title	Percentage of Participants with an AE Leading to TDV or Placebo Discontinuation <sup>[17]</sup>
-----------------	---

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. Safety Analysis Set included all randomized participants who received at least 1 dose of IP.

End point type	Primary
----------------	---------

End point timeframe:

From first vaccination on Day 1 through the end of trial (up to Day 270)

Notes:

[17] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	180	60	180	60
Units: percentage of participants				
number (not applicable)	0.6	0	0	0

## Statistical analyses

No statistical analyses for this end point

### Primary: Percentage of Participants with a Medically-attended AE (MAAE)

End point title	Percentage of Participants with a Medically-attended AE
-----------------	---

End point description:

MAAEs are defined as AEs leading to an unscheduled visit to or by a healthcare professional including visits to an emergency department but not fulfilling seriousness criteria. Safety Analysis Set included all randomized participants who received at least 1 dose of IP.

End point type	Primary
----------------	---------

End point timeframe:

From first vaccination on Day 1 through the end of trial (up to Day 270)

Notes:

[18] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	180	60	180	60
Units: percentage of participants				
number (not applicable)	5.0	3.3	10.0	13.3

## Statistical analyses

No statistical analyses for this end point

### Primary: Percentage of Participants with a Serious Adverse Event (SAE)

End point title	Percentage of Participants with a Serious Adverse Event
-----------------	---

End point description:

An AE is defined as any untoward medical occurrence in a clinical investigation participant administered a drug; it does not necessarily have to have a causal relationship with this treatment. An SAE is defined as any untoward medical occurrence that: results in death, is life-threatening, requires inpatient hospitalization or prolongation of existing hospitalization, results in persistent or significant disability/incapacity, leads to a congenital anomaly/birth defect in the offspring of a participant, is an important medical event. Safety Analysis Set included all randomized participants who received at least 1 dose of IP.

End point type	Primary
----------------	---------

End point timeframe:

From first vaccination on Day 1 through the end of trial (up to Day 270)

Notes:

[19] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	180	60	180	60
Units: percentage of participants				
number (not applicable)	0.0	0.0	0.0	0.0

## Statistical analyses

No statistical analyses for this end point

## Primary: GMTs of Neutralizing Antibodies by Microneutralization Test (MNT50) for Each of the 4 Dengue Virus Serotypes at Day 120

End point title	GMTs of Neutralizing Antibodies by Microneutralization Test (MNT50) for Each of the 4 Dengue Virus Serotypes at Day 120 <sup>[20]</sup>
-----------------	---

End point description:

GMTs of neutralizing antibodies were measured by microneutralization test 50% (MNT50) for each of the 4 dengue serotypes for all participants. The 4 dengue virus serotypes (DENV) are DENV-1, DENV-2, DENV-3 and DENV-4. FAS included all randomized participants who received at least 1 dose of TDV or placebo, and for whom a valid pre dose measurement, and at least 1 valid post dose measurement were received for immunogenicity assessments. Subjects analysed is the number of participants with data available for analysis. Number analyzed (n) is the number of participants with data available for analysis for the specified categories.

End point type	Primary
----------------	---------

End point timeframe:

Day 120 (Month 4)

Notes:

[20] - No statistical analyses have been specified for this primary end point. It is expected there is at least one statistical analysis for each primary end point.

Justification: Only descriptive analyses were planned for this endpoint.

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	176	59	178	59
Units: titers				
geometric mean (standard deviation)				
DENV-1 (n=176,59,178,59)	3196.9 (± 5.36)	676.6 (± 13.70)	2968.0 (± 4.50)	267.9 (± 14.31)
DENV-2 (n=176,59,178,59)	2547.3 (± 4.34)	624.2 (± 12.63)	2869.1 (± 3.78)	316.6 (± 15.66)
DENV-3 (n=176,59,177,59)	1616.0 (± 5.13)	452.3 (± 10.42)	1596.7 (± 4.82)	196.8 (± 14.67)
DENV-4 (n=176,59,178,59)	999.0 (± 4.65)	265.9 (± 8.42)	1018.4 (± 4.15)	135.8 (± 10.97)

## Statistical analyses

No statistical analyses for this end point

### Secondary: GMTs by MNT50 Against Each of the 4 Dengue Virus Serotypes at Day 1 and Day 270

End point title	GMTs by MNT50 Against Each of the 4 Dengue Virus Serotypes at Day 1 and Day 270
-----------------	---

End point description:

GMTs of neutralizing antibodies were measured by MNT50 for each of the 4 dengue serotypes for all participants. The 4 dengue virus serotypes are DENV-1, DENV-2, DENV-3 and DENV-4. FAS included all randomized participants who received at least 1 dose of TDV or placebo, and for whom a valid pre dose measurement, and at least 1 valid post dose measurement were received for immunogenicity assessments. Subjects analysed is the number of participants with data available for analysis. Number of subjects analysed is the number of participants with data available for analysis for the specified categories at the specified timepoints and was variable for each category.

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1 and 270

End point values	Cohort 1: ≥18 to ≤60 Years Age Group: TDV	Cohort 1: ≥18 to ≤60 Years Age Group: Placebo	Cohort 2: ≥4 to <18 Years Age Group: TDV	Cohort 2: ≥4 to <18 Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	176	59	178	59
Units: titers				
geometric mean (standard deviation)				
Day 1: DENV-1	1038.5 (± 10.63)	699.8 (± 12.42)	362.0 (± 14.27)	223.1 (± 16.76)
Day 1: DENV-2	797.9 (± 9.95)	460.1 (± 13.63)	418.4 (± 12.92)	314.1 (± 19.00)
Day 1: DENV-3	618.9 (± 9.23)	463.9 (± 10.91)	238.2 (± 11.60)	202.4 (± 17.11)
Day 1: DENV-4	392.4 (± 8.61)	275.3 (± 10.61)	166.8 (± 10.05)	147.3 (± 17.29)
Day 270: DENV-1	2557.9 (± 4.80)	703.3 (± 10.97)	1531.5 (± 6.27)	393.9 (± 12.85)
Day 270: DENV-2	2454.6 (± 3.69)	548.9 (± 12.17)	1740.7 (± 5.61)	441.7 (± 16.56)
Day 270: DENV-3	1067.8 (± 5.19)	402.9 (± 10.99)	765.2 (± 7.63)	287.8 (± 13.11)
Day 270: DENV-4	708.7 (± 4.94)	188.9 (± 9.79)	452.0 (± 6.61)	172.6 (± 11.28)

## Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of Participants With Seroconversion for Each of the 4 Dengue Virus Serotypes

End point title	Percentage of Participants With Seroconversion for Each of the 4 Dengue Virus Serotypes
-----------------	---

End point description:

Seroconversion rate, defined as the percentage of participants seropositive, is derived from the titers of dengue-neutralizing antibodies. Seropositive is defined as a reciprocal neutralizing titer  $\geq 10$ . The 4 dengue virus serotypes are DENV-1, DENV-2, DENV-3 and DENV-4. FAS included all randomized participants who received at least 1 dose of TDV or placebo, and for whom a valid pre dose measurement, and at least 1 valid post dose measurement were received for immunogenicity assessments. Subjects analysed is the number of participants with data available for analysis. Number of subjects analysed is the number of participants with data available for analysis for the specified categories at the specified timepoints and was variable for each category.

End point type	Secondary
----------------	-----------

End point timeframe:

Day 1, Day 120 and Day 270

End point values	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: TDV	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: Placebo	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: TDV	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	176	59	178	59
Units: percentage of participants				
number (not applicable)				
Day 1: DENV-1	90.9	86.4	82.6	76.3
Day 1: DENV-2	91.5	83.1	87.1	83.1
Day 1: DENV-3	88.6	86.4	79.2	72.9
Day 1: DENV-4	85.8	79.7	78.1	69.5
Day 120: DENV-1	98.3	83.1	100	83.1
Day 120: DENV-2	98.3	86.4	100	78.0
Day 120: DENV-3	98.9	86.4	100	71.2
Day 120: DENV-4	97.7	84.7	100	72.9
Day 270: DENV-1	98.9	89.8	99.4	86.2
Day 270: DENV-2	99.4	89.8	98.3	81.0
Day 270: DENV-3	98.3	83.1	96.6	81.0
Day 270: DENV-4	95.4	78.0	92.7	75.9

## Statistical analyses

No statistical analyses for this end point

### Secondary: Percentage of Participants With Seroconversion for Multiple (2, 3, or 4) Dengue Virus Serotypes

End point title	Percentage of Participants With Seroconversion for Multiple (2, 3, or 4) Dengue Virus Serotypes
-----------------	---

---

**End point description:**

Seroconversion rate, defined as the percentage of participants seropositive, is derived from the titers of dengue-neutralizing antibodies. Seropositive is defined as a reciprocal neutralizing titer  $\geq 10$ . The 4 dengue virus serotypes are DENV-1, DENV-2, DENV-3 and DENV-4. FAS included all randomized participants who received at least 1 dose of TDV or placebo, and for whom a valid pre dose measurement, and at least 1 valid post dose measurement were received for immunogenicity assessments. Subjects analysed is the number of participants with data available for analysis. Number of subjects analysed is the number of participants with data available for analysis for the specified categories at the specified timepoints and was variable for each category.

---

End point type	Secondary
----------------	-----------

---

**End point timeframe:**

Day 1, Day 120 and Day 270

---

End point values	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: TDV	Cohort 1: $\geq 18$ to $\leq 60$ Years Age Group: Placebo	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: TDV	Cohort 2: $\geq 4$ to $< 18$ Years Age Group: Placebo
Subject group type	Reporting group	Reporting group	Reporting group	Reporting group
Number of subjects analysed	176	59	178	59
Units: percentage of participants				
number (not applicable)				
Day 1: Monovalent	5.7	5.1	8.4	8.5
Day 1: Bivalent	2.3	1.7	4.5	5.1
Day 1: Trivalent	1.1	5.1	0.6	1.7
Day 1: Tetravalent	85.8	78.0	77.0	69.5
Day 120: Monovalent	0.0	1.7	0.0	13.6
Day 120: Bivalent	0.6	3.4	0.0	3.4
Day 120: Trivalent	1.1	0.0	0.6	0
Day 120: Tetravalent	97.2	83.1	99.4	71.2
Day 270: Monovalent	0.6	13.6	0.6	10.3
Day 270: Bivalent	0.6	0.0	1.7	5.2
Day 270: Trivalent	4.0	5.1	5.6	0
Day 270: Tetravalent	94.3	78.0	91.6	75.9

---

**Statistical analyses**

No statistical analyses for this end point

---

## Adverse events

### Adverse events information

Timeframe for reporting adverse events:

All cause mortality and SAE: From Day 1 through end of trial on Day 270; Non-serious adverse events: Up to 28 days (day of vaccination + 27 days) following administration of TDV or placebo

Adverse event reporting additional description:

The Safety Analysis Set included all randomized participants who received at least 1 dose of IP.

Assessment type	Systematic
-----------------	------------

### Dictionary used

Dictionary name	MedDRA
-----------------	--------

Dictionary version	28.0
--------------------	------

### Reporting groups

Reporting group title	Cohort 1: $\geq 18$ to $\leq 60$ Age Group: TDV
-----------------------	---

Reporting group description:

Participants randomized to receive TDV subcutaneous (SC) injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Reporting group title	Cohort 1: $\geq 18$ to $\leq 60$ Age Group: Placebo
-----------------------	---

Reporting group description:

Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Reporting group title	Cohort 2: $\geq 4$ to $< 18$ Age Group: TDV
-----------------------	---

Reporting group description:

Participants received TDV SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Reporting group title	Cohort 2: $\geq 4$ to $< 18$ Age Group: Placebo
-----------------------	---

Reporting group description:

Participants received TDV-matching placebo SC injection on Day 1 (Month 0) and Day 90 (Month 3) of the study.

Serious adverse events	Cohort 1: $\geq 18$ to $\leq 60$ Age Group: TDV	Cohort 1: $\geq 18$ to $\leq 60$ Age Group: Placebo	Cohort 2: $\geq 4$ to $< 18$ Age Group: TDV
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 180 (0.00%)	0 / 60 (0.00%)	0 / 180 (0.00%)
number of deaths (all causes)	0	0	0
number of deaths resulting from adverse events	0	0	0

Serious adverse events	Cohort 2: $\geq 4$ to $< 18$ Age Group: Placebo		
Total subjects affected by serious adverse events			
subjects affected / exposed	0 / 60 (0.00%)		
number of deaths (all causes)	0		
number of deaths resulting from adverse events	0		



Frequency threshold for reporting non-serious adverse events: 5 %

<b>Non-serious adverse events</b>	Cohort 1: >=18 to <=60 Age Group: TDV	Cohort 1: >=18 to <=60 Age Group: Placebo	Cohort 2: >=4 to <18 Age Group: TDV
Total subjects affected by non-serious adverse events subjects affected / exposed	53 / 180 (29.44%)	19 / 60 (31.67%)	52 / 180 (28.89%)
Investigations Body temperature subjects affected / exposed occurrences (all)	5 / 180 (2.78%) 10	4 / 60 (6.67%) 7	5 / 180 (2.78%) 6
Nervous system disorders Headache subjects affected / exposed occurrences (all)	20 / 180 (11.11%) 42	7 / 60 (11.67%) 18	8 / 180 (4.44%) 18
General disorders and administration site conditions Asthenia subjects affected / exposed occurrences (all)  Pain subjects affected / exposed occurrences (all)	15 / 180 (8.33%) 37  38 / 180 (21.11%) 89	5 / 60 (8.33%) 13  8 / 60 (13.33%) 17	10 / 180 (5.56%) 26  41 / 180 (22.78%) 74
Musculoskeletal and connective tissue disorders Myalgia subjects affected / exposed occurrences (all)	14 / 180 (7.78%) 35	4 / 60 (6.67%) 8	7 / 180 (3.89%) 17

<b>Non-serious adverse events</b>	Cohort 2: >=4 to <18 Age Group: Placebo		
Total subjects affected by non-serious adverse events subjects affected / exposed	15 / 60 (25.00%)		
Investigations Body temperature subjects affected / exposed occurrences (all)	1 / 60 (1.67%) 6		
Nervous system disorders Headache			

subjects affected / exposed occurrences (all)	1 / 60 (1.67%) 3		
General disorders and administration site conditions Asthenia subjects affected / exposed occurrences (all)  Pain subjects affected / exposed occurrences (all)	3 / 60 (5.00%) 11  15 / 60 (25.00%) 23		
Musculoskeletal and connective tissue disorders Myalgia subjects affected / exposed occurrences (all)	2 / 60 (3.33%) 6		

## More information

### Substantial protocol amendments (globally)

Were there any global substantial amendments to the protocol? No

---

### Interruptions (globally)

Were there any global interruptions to the trial? No

### Limitations and caveats

None reported